ZooKeys 628: 65–246 (2016)
doi: 10.3897/zookeys.628.9799
http://zookeys.pensoft.net

CATALOGUE

A peer-reviewed open-access journal
ZooKeys
Launched to accelerate biodiversity research

Revised classification and catalogue of global Nepticulidae and Opostegidae (Lepidoptera, Nepticuloidea)

Erik J. van Nieukerken[1], Camiel Doorenweerd[1], Robert J. B. Hoare[2], Donald R. Davis[3]

1 *Naturalis Biodiversity Center, PO Box 9517, NL-2300 RA Leiden, The Netherlands* **2** *Landcare Research Ltd., Private Bag 92–170, Auckland, New Zealand* **3** *Department of Entomology, National Museum of Natural History, MRC 105, Smithsonian Institution, PO Box 37012, Washington, DC 20013–7012, USA*

Corresponding author: *Erik J. van Nieukerken* (nieukerken@naturalis.nl)

Academic editor: *T. Simonsen* | Received 8 July 2016 | Accepted 25 August 2016 | Published 31 October 2016

http://zoobank.org/2D256553-0AFA-45C8-97EA-B3A006CFF3F7

Citation: Nieukerken EJ van, Doorenweerd C, Hoare RJB, Davis DR (2016) Revised classification and catalogue of global Nepticulidae and Opostegidae (Lepidoptera, Nepticuloidea). ZooKeys 628: 65–246. doi: 10.3897/zookeys.628.9799

Abstract

A catalogue of all named Nepticulidae and Opostegidae is presented, including fossil species. The catalogue is simultaneously published online in the scratchpad http://nepticuloidea.info/ and in Catalogue of Life (http://www.catalogueoflife.org/col/details/database/id/172). We provide a historical overview of taxonomic research on Nepticuloidea and a brief 'state of the art'. A DNA barcode dataset with 3205 barcodes is made public at the same time, providing DNA barcodes of ca. 779 species, of which 2563 are identified as belonging to 444 validly published species. We recognise 862 extant and 18 fossil species of Nepticulidae in 22 extant genera and the fossil form genus *Stigmellites*. We count 192 valid Opostegidae species in 7 genera, without fossils. We also list seven dubious Nepticulidae names that cannot be placed due to absent type material and poor descriptions, 18 unavailable names in Nepticulidae that cannot be placed and we also list the 33 names (including four fossils) that once were placed as Nepticulidae or Opostegidae but are now excluded. All synonyms and previous combinations are listed. The generic classification follows the Molecular phylogeny that is published almost simultaneously. Subfamilies and tribes are not recognised, Trifurculinae Scoble, 1983 is synonymised with Nepticulidae Stainton, 1854 and Opostegoidinae Kozlov, 1987 is synonymised with Opostegidae Meyrick, 1893. The status of *Casanovula* Hoare, 2013, *Etainia* Beirne, 1945, *Fomoria* Beirne, 1945, *Glaucolepis* Braun, 1917, *Menurella* Hoare, 2013, *Muhabbetana* Koçak & Kemal, 2007 and *Zimmermannia* Hering, 1940 is changed from subgenus to full genus, whereas two

Copyright *Erik J. van Nieukerken et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

genera are considered synonyms again: *Manoneura* Davis, 1979, a synonym of *Enteucha* Meyrick, 1915 and *Levarchama* Beirne, 1945, a synonym of *Trifurcula* Zeller, 1848. We propose 87 new combinations in Nepticulidae and 10 in Opostegidae, largely due to the new classification, and re-examination of some species. We propose the following 37 new synonymies for species (35 in Nepticulidae, 2 in Opostegidae):

Stigmella acerifoliella Dovnar-Zapolski, 1969 (unavailable, = *S. acerna* Puplesis, 1988), *Stigmella nakamurai* Kemperman & Wilkinson, 1985 (= *S. palionisi* Puplesis, 1984), *Nepticula amseli* Skala, 1941 (unavailable = *S. birgittae* Gustafsson, 1985), *Stigmella cathepostis* Kemperman & Wilkinson, 1985 (= *S. microtheriella* (Stainton, 1854)), *Stigmella populnea* Kemperman & Wilkinson, 1985 (= *S. nivenburgensis* (Preissecker, 1942)), *Nepticula obscurella* Braun, 1912 (revised synonymy, = *S. myricafoliella* (Busck, 1900)), *Nepticula mandingella* Gustafsson, 1972 (= *S. wollofella* (Gustafsson, 1972)), *Stigmella rosaefoliella pectocatena* Wilkinson & Scoble, 1979 (= *S. centifoliella* (Zeller, 1848)), *Micropteryx pomivorella* Packard, 1870 (= *S. oxyacanthella* (Stainton, 1854)), *Stigmella crataegivora* Puplesis, 1985 (= *S. micromelis* Puplesis, 1985), *Stigmella scinanella* Wilkinson & Scoble, 1979 (= *S. purpuratella* (Braun, 1917)), *Stigmella palmatae* Puplesis, 1984 (= *S. filipendulae* (Wocke, 1871)), *Stigmella sesplicata* Kemperman & Wilkinson, 1985 (= *S. lediella* (Schleich, 1867)), *Stigmella rhododendrifolia* Dovnar-Zapolski & Tomilova, 1978 (unavailable, = *S. lediella* (Schleich, 1867)), *Stigmella oa* Kemperman & Wilkinson, 1985 (= *S. spiculifera* Kemperman & Wilkinson, 1985), *Stigmella gracilipae* Hirano, 2014 (= *S. monticulella* Puplesis, 1984), *Nepticula chaoniella* Herrich-Schäffer, 1863 (= *S. samiatella* (Zeller, 1839)), *Bohemannia piotra* Puplesis, 1984 (= *B. pulverosella* (Stainton, 1849)), *Bohemannia nipponicella* Hirano, 2010 (= *B. manschurella* Puplesis, 1984), *Sinopticula sinica* Yang, 1989 (= *Glaucolepis oishiella* (Matsumura, 1931)), *Trifurcula collinella* Nel, 2012 (= *Glaucolepis magna* (A. Laštuvka & Z. Laštuvka, 1997)), *Obrussa tigrinella* Puplesis, 1985 (= *Etainia trifasciata* (Matsumura, 1931)), *Microcalyptris vittatus* Puplesis, 1984 and *M. arenosus* Falkovitsh, 1986 (both = *Acalyptris falkovitshi* (Puplesis, 1984)), *Ectoedemia castaneae* Busck, 1913, *E. heinrichi* Busck, 1914 and *E. helenella* Wilkinson, 1981 (all three = *Zimmermannia bosquella* (Chambers, 1878)), *Ectoedemia chloranthis* Meyrick, 1928 and *E. acanthella* Wilkinson & Newton, 1981 (both = *Zimmermannia grandisella* (Chambers, 1880)), *Ectoedemia coruscella* Wilkinson, 1981 (= *Zimmermannia mesoloba* (Davis, 1978)), *Ectoedemia piperella* Wilkinson & Newton, 1981 and *E. reneella* Wilkinson, 1981 (both = *Zimmermannia obrutella* (Zeller, 1873)), *Ectoedemia similigena* Puplesis, 1994 (= *E. turbidella* (Zeller, 1848)), *Ectoedemia andrella* Wilkinson, 1981 (= *E. ulmella* (Braun, 1912)), *Nepticula canadensis* Braun, 1917 (= *E. minimella* (Zetterstedt, 1839)), *Opostega rezniki* Kozlov, 1985 (= *O. cretatella* Chrétien, 1915), *Pseudopostega cyrneochalcopepla* Nel & Varenne, 2012 (= *P. chalcopepla* (Walsingham, 1908)). *Stigmella caryaefoliella* (Clemens, 1861) and *Zimmermannia bosquella* (Chambers, 1878) are taken out of synonymy and re-instated as full species. Lectotypes are designated for *Trifurcula obrutella* Zeller, 1873 and *Nepticula grandisella* Chambers, 1880.

Keywords

Taxonomy, leaf miners, checklist, history, new synonyms, new combinations

Table of contents

Introduction 68
History of taxonomic research on Nepticulidae 69
Taxonomic history of Opostegidae 82
Material and methods 84
Taxonomic practice 84

Species 84
Subspecies 84
Higher categories 85
Species groups 86
Order of the list 87
Distribution 87
DNA barcodes and specimen data 87
Host plants 88
Nomenclatorial practice 89
References 89
Combinations 90
Synonyms 90
Types 91
Abbreviations and symbols used 91
Global Nepticuloidea: the state of the art 92
Catalogue 102
Family Nepticulidae 102
Enteucha 102
Varius 103
Simplimorpha 103
Stigmella 104
Non-core *Stigmella* 104
Core *Stigmella* 112
Ozadelpha 130
Roscidotoga 130
Casanovula 130
Menurella 130
Pectinivalva 131
Neotrifurcula 131
Hesperolyra 132
Bohemannia 132
Areticulata 133
Glaucolepis 133
Trifurcula 136
Fomoria 138
Muhabbetana 141
Parafomoria 144
Etainia 145
Acalyptris 147
Zimmermannia 151
Ectoedemia 153
Stigmellites 161
Nomina dubia et oblita 161

Unplaced unavailable names 161
Family Opostegidae 162
Notiopostega 162
Eosopostega 162
Neopostega 162
Paralopostega 162
Opostegoides 163
Opostega 164
"*Opostega*" 164
Pseudopostega 165
Taxa excluded from Nepticuloidea 169
Notes 171
Acknowledgements 201
References 202
Supplementary material 1 246
Supplementary material 2 246

Introduction

Names of organisms are the key to the biological literature, but even in our digital age it is often a challenge to find and apply the correct names. Taxonomic internet databases, spawned from global projects such as GBIF (Gbif Secretariat 2016) and Species 2000 (Catalogue of Life: Roskov et al. 2015) are growing in number and are impressively linked together, but in many cases lack informative and reliable content. Catalogue of Life is especially poor for Lepidoptera, being based on an online version of the Cardindex of the Natural History Museum in London (LepIndex: Beccaloni et al. 2005). This is a wonderful resource for taxonomists, but not an authoritative catalogue with modern classification and names. The plans for an update depend on external funding and may never materialize with the current taxonomic funding climate (I. Kitching personal communication).

Any online only publication of a catalogue has the disadvantage that nomenclatorial changes are unavailable and still need to be published in a unchangeable format on paper or as pdf file following the amendments made to the Code (International Commission on Zoological Nomenclature 2012). It is understandable that many taxonomists prefer paper publications, evident from several recent catalogues for Lepidoptera (Pterophoridae, Coleophoridae, Psychidae, Notodontidae, Yponomeutoidea) that cannot yet be found in an online database (Gielis 2003; Baldizzone et al. 2006; Sobczyk 2011; Schintlmeister 2013; Lewis and Sohn 2015). Fortunately other authors followed a printed catalogue soon with an online database version, e.g. for Gracillariidae (De Prins and De Prins 2005; De Prins and De Prins 2016) and Tortricidae (Brown et al. 2003; Gilligan et al. 2014) and for other groups there are online-only catalogues that are actively maintained, e.g. Pyraloidea (Nuss et al. 2003–2015) and part of the butterflies (Häuser et al. 2012). Still, for the largest part of Lepidoptera there are still

no global catalogues available. With our contribution we hope to fill a very small gap of the megadiverse order Lepidoptera.

For some years we have been preparing an online catalogue of Nepticuloidea in a so-called scratchpad (van Nieukerken 2016), which has several advantages, but also the cited disadvantage and the lack of an easy overview. The most recent Nepticuloidea catalogue was published in a book (Diškus and Puplesis 2003), using a somewhat different generic classification. Our phylogenetic study, to be published simultaneously with this catalogue (Doorenweerd et al. 2016a), resulted in the need for a new classification, which has consequences for the names of many taxa. We therefore decided to publish this static catalogue to fix the state of the art, make the necessary nomenclatorial changes, whereas at the same time the Nepticuloidea scratchpad version 2.0 (http://nepticuloidea.info/) is released with additional information and illustrations, and will be updated continuously. We are also happy to announce that at the same time this catalogue is made available to the Catalogue of Life (http://www.catalogueoflife.org/col/details/database/id/172) and GBIF. This catalogue is not only the taxonomic summary of the published knowledge on these families, it also contains many original data, such as new synonymies and taxonomic placement of species, based on our many years of research of these insects worldwide; some of these results are formalised here, but will be detailed elsewhere. In the material and methods section we discuss our various choices for this catalogue, such as species concepts, use of DNA barcodes, and nomenclatorial issues.

To place the list in a broader context we give an account of the taxonomic history and history of research on Nepticulidae and Opostegidae. We begin the results section with a review of the "state of the art" of Nepticuloidea in the various biogeographic regions.

History of taxonomic research on Nepticulidae

The first species of Nepticulidae that was described, was *Stigmella anomalella* (Goeze, 1783) of which Degeer (1752) described and illustrated the larva in detail, while using the term "miner" ("mineuse") for the first time (van Nieukerken 2008). Later, Degeer (1771) described the adult that he reared from these mines. Degeer's descriptions are still interesting reading these days, and were fully copied and translated by Stainton et al. (1855). Since Degeer did not use binominal nomenclature in these works, the species was only validly named later by Goeze (1783) as *Phalaena anomalella*. This makes *Ectoedemia occultella* (Linnaeus, 1767) the first formally named nepticulid (as *Phalaena (Tinea) occultella*), although Linnaeus himself had yet no notion of the life history of the small moth that he found on his windows (Robinson and Nielsen 1983). Five more names were given to Nepticulidae in the 18th century, all named in *Tinea*, but just two of these are still considered valid: *Stigmella aurella* and *S. hybnerella*. Even though Schrank already in 1802 gave the generic name *Stigmella* to his rose leafminer *Tinea rosella* (a junior synonym of *S. anomalella*), this was overlooked by most 19th century authors, and in the first decades of the 19th century, species were still placed in *Tinea* (e.g. by Haworth 1828), the first author to describe a number of species from the London area, nine in total) or in a number of other "tineid" genera such as *Microsetia* (Stephens 1834; Be-

Figures 1–9. Portraits of Lepidopterists who described Nepticuloidea. **1** Philipp Christoph Zeller (Stainton 1883) **2** Henry Tibats Stainton from Douglas and Mclachlan (1893) **3** Gottlieb August Wilhelm Herrich-Schäffer, ca 1870 (Kraatz 1875) **4** Maximilian Ferdinand Wocke (Dittrich 1907) **5** Edward Meyrick (Clarke 1955) **6** Lord Walsingham (Durrant 1920) **7** Annette Francis Braun, in 1973 in her home in Cincinnati, Ohio (photo Mignon Davis) **8** Pierre Chrétien, Digne, May 1903 (Oberthür 1915) **9** Erich Martin Hering. Figs 1–6 and 8 from Biodiversity Heritage Library, 9 from Zobodat.

dell 1848), *Oecophora* and *Elachista* (Kollar 1832; Zetterstedt 1838–1840; Duponchel [1842–1845]) and *Caloptilia* (Hübner 1816–1826), now all belonging in different families, but together these authors named only about ten species of Nepticulidae.

Towards the second half of the 19th century, things changed rapidly, in Europe attributable to four naturalists: in Germany Philipp Christoph Zeller (1808–1883) (Fig. 1) and Gottlieb August Wilhelm Herrich-Schäffer (1799–1874, Regensburg) (Fig. 3), in Great Britain Henry Tibbats Stainton (1822–1892) (Fig. 2) and in Switzerland Heinrich Frey (1822–1890). All four were lepidopterists with a broad interest, studying not only all Lepidoptera from their own country, but also from many exotic countries.

The genus *Nepticula* was formally erected in a meeting report by von Heyden (1843), but he did not publish much on this genus, and it was Zeller (1848) who made a detailed description of the genus and several species. Some years earlier Zeller (1839) had described several Nepticulidae (in his "Versuch einer naturgemässen Eintheilung der Schaben") together with other leafminers in the genus *Lyonetia*, subgenus *Bucculatrix*, in which he placed ten nepticulid species together with six that we still consider as *Bucculatrix* (Bucculatricidae). All these descriptions were based on adults that were collected in the field without knowledge of the life history (except DeGeer's work!). Stainton and his British followers drastically changed this situation, by eagerly rearing leafminers and many other microlepidoptera, resulting in a proliferation of new species discoveries. Following a few initial species descriptions, his book "Insecta Brittanica. Lepidoptera: Tineina" (Stainton 1854) contained the first extensive treatment of Nepticulidae – at that point recognised as a family with two genera (*Nepticula* and *Trifurcula*) – with detailed information on leafmines and biology of 32 species in total (see Table 1). Stainton continued publishing discoveries the next decades, partly using his own periodicals "The Entomologist's Weekly Intelligencer" and "The Entomologist's Annual", and later in the still running journal "Entomologist's Monthly Magazine" that he founded with entomologist friends (Emmet 1992). His magnum opus is the 13 volume work "The natural history of the Tineina", that he wrote collaboratively with Zeller, Frey and J.W. Douglas. Nepticulidae are treated in volumes 1 and 7 (Stainton et al. 1855; Stainton et al. 1862). These books are best known for the exquisite hand coloured plates with details of all life stages of many Microlepidoptera. During his life Stainton described 39 species of Nepticulidae, of which currently 30 are still considered valid names. With that record he is the most productive 19th century nepticulid taxonomist.

Soon after Stainton's Insecta Britannica, Frey wrote two books dealing with respectively Swiss and European Nepticulidae (Frey 1856; 1857), that he also partly had reared himself. In the same period, Herrich-Schäffer's large multivolume work "Systematische Bearbeitung der Schmetterlinge von Europa" (1843–1855) was concluded by volumes 5 and 6, dealing with Nepticulidae in volume 5, heft 67 (Herrich-Schäffer 1855a) (the book was issued in parts, the plates often earlier than the text, and the bibliography is very complicated, see Hemming 1937). He apparently did not rear nepticulids himself, but included information on rearing from others, such as Frey and Heyden. He also named several species that Frey intended to name (as he did later, in 1856), but changed the endings of the names deliberately into "–ella". This explains pairs of names such as: *Nepticula aeneofasciella* Herrich-Schäffer, 1855 and *Nepticula*

Table 1. Number of Nepticulidae species (valid and invalid) described per first author for authors who described at least ten valid species, including extant and fossil species.

First author	Valid spp	Invalid spp	Total
Puplesis (Stonis)	176	22	198
Scoble	73	1	74
Meyrick	67	6	73
van Nieukerken	55		55
Klimesch	38	10	48
Stainton	30	9	39
Braun	26	11	37
Remeikis	24		24
Kemperman	23	13	36
Laštuvka, A.	21		21
Vári	20		20
Laštuvka, Z.	17		17
Diškus	15		15
Donner	14		14
Clemens	13	7	20
Zeller	12	5	17
Chambers	12	12	24
Hoare	12		12
Herrich-Schäffer	11	10	21
Frey	10	8	18
Hirano	10	2	12
96 authors	201		
Total	**880**		

aeneofasciata Frey, 1856. The practice to use consistent endings for larger groups of Lepidoptera dated from Linnaeus, who used –ella for what he called *Tinea*, -ana for *Tortrix*, -alis for *Pyralis* etc. (Emmet 1991). In the 19th century there were no hard nomenclatorial rules yet, and specialists often changed names deliberately following this practice, see e.g. also several listed synonyms (incorrect subsequent spellings) by Doubleday (1859). Herrich-Schäffer named several nepticulids hidden in catalogues and a report on a collecting trip to Engadin (Herrich-Schäffer 1863b; c), names that have completely been forgotten and were never listed in catalogues, but only cited in two other papers (Snellen 1873; Segerer 1997). As far as these names are valid, they can be considered nomina oblita once their identities have been established, since they were never used as valid names since 1899 (ICZN art. 23.9.1.1), and do not compete with junior synonyms (reversal of precedence, ICZN art. 23).

By 1860 the number of species (all from Europe) had been tripled since 1848: from 24 to 77 (see Fig. 24B). In Europe the work on describing Nepticulidae continued both in Britain and in Central Europe, particularly with Herman von Heinemann (1812–1871) in Braunschweig (Heinemann 1861; 1862a; b) and Maximilian Ferdinand Wocke (1820–1906) (Fig. 4) in Breslau, who contributed many shorter papers (Wocke 1862;

1865; 1877). Jointly they wrote a larger fauna: Die Schmetterlinge Deutschlands und der Schweiz, partly finished after Heinemann's death (Heinemann and Wocke [1876]). After *ca* 1880 taxonomic work on Nepticulidae in Europe slowed down considerably.

Meanwhile, in the United States, James Brackenridge Clemens (1825–1867) in Easton, Pennsylvania, had started studies on microlepidoptera. He collected the leafmines actively and described them in a few papers, but in contrast to Stainton he did not wait for successful rearing results and described many new species solely on the basis of leafmines (Clemens 1861; 1862a; Clemens 1862b; Clemens 1865), often resulting in a conundrum for future taxonomists. There are no types of the mines left, and more than once multiple species are known to feed on the host plant from which he described a single species. Such species have to be interpreted carefully and sometimes Neotypes need to be selected (Busck 1903; Wilkinson and Scoble 1979). A second North American pioneer was Vactor Tousey Chambers (1830–1883) from Covington, Kentucky, who added more species on the basis of mines and larvae, but also many on the basis of adults (Chambers 1873; 1875b; 1878a). There are more types left of his species (in the Museum of Comparative Zoology, Cambridge, Mass.), but also many missing and reconstructing identities can still be cumbersome (Busck 1903; Miller and Hodges 1990). Here we resolve three of his names.

The first Nepticulidae described from other regions than Europe or the USA, were two species described from Colombia by Zeller (1877), although Francis Walker's (Walker 1864) *Stigmella maoriella* from New Zealand already predated these. However, Walker, often criticised for his numerous useless descriptions, was totally unaware he was naming a nepticulid and placed it in *Tinea*.

In the late 19th century Edward Meyrick (1854–1939) (Fig. 5) started his long career of describing numerous exotic Microlepidoptera, totalling approximately 20,000 species, including 68 Nepticulidae and 54 Opostegidae. The first Nepticulidae he described were three species from New Zealand (Meyrick 1889) where he lived in 1882–1883 (Hudson 1938). Up to 1935 Meyrick described many Nepticulidae from southern Africa, India, Australia, New Zealand and South America. His descriptions are typically very short, without illustrations, and only some of the species were reared (by other collectors who sent the specimens to him). When studying genitalia became more common practice, Meyrick refused to see the necessity and continued in a similar manner, relying almost completely on venation and colour pattern (Clarke 1955). As a result almost none of the species he described can be recognised from the description alone and the study of types is essential. Fortunately, most of those are kept in the Natural History Museum in London, and some in other collections, including the Ditsong Museum of Natural History in Pretoria. Those types that still exist have been studied by Malcolm Scoble, Erik van Nieukerken, Robert Hoare and co-authors and Rimantas Puplesis (now Jonas Stonis). Meyrick named only two nepticulid genera: *Acalyptris* and *Enteucha*, the latter he placed in Opostegidae (Davis 1985).

Quite a different approach was followed in the early 20th century in North America, where August Busck (1870–1944), a Danish immigrant, started his studies at the Division of Entomology of the U.S. Department of Agriculture and somewhat later Annette Frances Braun (1884–1978) (Fig. 7), who became the leading microlepidop-

terists of North America. Both made many careful descriptions of often reared species, but most of these before genitalia studies became fashionable (Busck 1900; 1913; 1914a; Braun 1912; 1914; 1917; 1923; 1925b).

In Europe, in addition to *Nepticula*, only two additional small genera were recognised on the basis of the venation: *Trifurcula* Zeller, 1848 and *Bohemannia* Stainton, 1859 (Zeller 1848; Stainton 1859a). Although many treatments of the family were published in Europe, the classification did not change for almost a century (Heinemann 1862a; b; Frey 1880; Snellen 1882; Meyrick 1895; Tutt 1899; Spuler and Meess 1910; Heinemann and Wocke [1876]). Walsingham (1908a) had shown that the senior name *Stigmella* Schrank, 1802 should be used instead of *Nepticula*, but few authors followed him until this case was settled by Wilkinson (1978) in favour of the senior name *Stigmella*. Between them, Busck and Braun erected four more genera on the basis of venation: *Ectoedemia* Busck, 1907 (also based on the galling habit), *Obrussa* Braun, 1915, *Glaucolepis* Braun, 1917 and *Microcalyptris* Braun, 1925 (Busck 1907; Braun 1915; 1917; 1925b) (see Table 2). It is strange that these authors who relied so much on venation in describing genera, did not scrutinize the many *Nepticula* species, amongst which were still many species with venation different from *Stigmella*, and more similar to their new genera. Also the name *Trifurcula* was several times misapplied to species with completely different venations, now belonging in *Zimmermannia* (*atrifrontella* Stainton, *obrutella* Zeller), *Bohemannia* (*pulverosella* Stainton) or *Acalyptris* (*minimella* Rebel).

In Europe, during the first decades of the 20th century, some more southern areas were explored by Lord Walsingham (Thomas de Grey, 6th Baron Walsingham, 1843–1919) (Fig. 6), who collected in France, Spain, the Canary Islands, Morocco and Algeria (Walsingham 1891; 1904; 1908a; b; 1911) and reared several of these species from their mines. Also Pierre Chrétien (1846–1934) (Fig. 8) reared several species from hitherto unexpected host plants in France and Algeria (*Bupleurum* in Apiaceae, *Linum* in Linaceae, *Launaea* [as *Zollikofferia*] in Asteraceae); these species are now all placed in *Glaucolepis* (Chrétien 1904; 1907; 1914).

In the 20th century the study of genitalia gradually became standard in Lepidoptera and the first study on nepticulids was by Petersen (1930b), who divided *Nepticula* into a number of species groups based on the male genitalia. The first following him was Hering (Hering 1942; 1943), but his first genitalia drawings were published by Klimesch (Klimesch 1940a; b; c; d), who in the last of these papers also shows his first own genitalia drawing (of *Acalyptris platani*). The single contribution by Bryan P. Beirne (1918–1998) was particularly important. He finished the British series of Lepidoptera genitalia started by Pierce and Metcalfe (1935) – who excluded most Nepticulidae, apart from *Trifurcula*, in their 'tineid' volume, referring to Petersen's work – , and described the male genitalia of the British species (Beirne 1945). While doing so he also erected five new genera with names based on Irish mythology (*Dechtiria*, *Levarchama*, *Fedalmia*, *Etainia* and *Fomoria*) and split *Stigmella* into *Stigmella* and *Nepticula*. From then on describing a new species without illustrating the male genitalia seldom occurred anymore, and the late 1940's showed several examples of this (Doets 1947; Janse 1948; Klimesch 1948a; b; Hartig 1949; Ford 1950a). Josef Wilhelm Klimesch (1902–1997) (Fig. 10) was one of the most productive authors of this period, and again in the 1970's after his

Table 2. Number of genera of Nepticuloidea named per author, including replacement names.

Author	# Valid genera	# Invalid genera
Davis (1 with Stonis)	4	3
Scoble	4	1
Hoare	3	
van Nieukerken	3	1
Beirne	2	3
Kozlov	2	
Meyrick	2	
Zeller	2	
Borkowski	1	1
Braun	1	2
Busck	1	
Hering	1	
Koçak	1	1
Schrank	1	
Stainton	1	
Müller-Rutz		2
Börner		1
Heyden		1
Puplesis		1
Strand		1
Yang		1

retirement (Gusenleitner 1988; Deschka 1998; Aspöck 2003). He not only carefully illustrated the genitalia, he also paid considerable attention to the leafmines and biology, with detailed illustrations and descriptions. Possibly because of the greater demands that including genitalia imposed on authors, the descriptions of new species decreased considerably during the 1950's and 1960's, only to rise again from 1978 on. Between 1950 and 1978 only 60 new species were named, most by Josef Klimesch in Europe and Lajos Vári in South Africa (Klimesch 1951b; 1953a; b; 1975c; Vári 1955; 1963).

Quite a reverse development occurred in the first half of the 20th century that involved an increased interest in the mine form and the host plant, especially promoted by Erich Martin Hering (1893–1967) (Fig. 9), an entomologist in Berlin. Although he was also a very active and experienced taxonomist in Lepidoptera and Diptera, who introduced the use of genitalia characters, he also gave often undue weight to characters of leafmines, for example in the *Stigmella ulmivora* group, where he recognised three species on the basis of leafmine form and frass pattern, that are in fact ecological forms of one species. However, with his many publications, Hering's work was an outstanding contribution to the study of leafminers, and his identification keys are still a much used reference (Hering 1957), as is his general book on the biology of leafminers (Hering 1951). Also Victor Hugo Otto Skala (1875–1952) studied leafminers intensively, but he also named many species on the basis of the mines only (which are unavailable after 1930, ICZN art. 13.6.2), provided with very poor and minute illustrations and brief descriptions (example: Skala 1939a; b;

Figures 10–16. Specialists on Nepticuloidea in 20th century. **10** Joseph Klimesch, in his house, Linz, October 1983 **11** Donald R. Davis in his office, December 2006, USNM, Washington DC **12** Christopher Wilkinson, April 1982, Cambridge, European Congress of Lepidopterology **13** Arthur Maitland Emmet, with C Wilkinson, Essex, UK, October 1979 **14** Meeting for preparation of the Nepticulidae volumes for Fauna Entomologica Scandinavica, Copenhagen, October 1980: from left: Ebbe S. Nielsen, Roland Johansson, Chris Wilkinson, Bert Gustafsson, Niels Peder Kristensen **15** Rimantas Puplesis, standing (now Jonas R. Stonis), Erik J. van Nieukerken, in Zoological Institute, Leningrad, September 1985 **16** Get together of leafminer enthusiasts at house of Aleš Laštůvka, Prostejov, Czech Rep., September 1994, from left: Steven Whitebread, Zdeněk Laštůvka, Roland Johansson, Aleš Laštůvka. Scans from transparencies by Erik J. van Nieukerken.

c; d; e; f; g; h), and only few of the names he introduced, and based on adults, remain valid today (*Ectoedemia klimeschi* (Skala, 1933), *Fomoria groschkei* (Skala, 1943)).

During the 1970's the study of nepticulids suddenly received considerable increase in support from several sources, following a general trend for greater investment in taxonomy and science at large and an increase of amateur interest. In Sweden amateur Roland Johansson (Figs 14, 16) revised the confusing oak feeding *Stigmella ruficapitella* group (Johansson 1971). In the same paper he provided an updated checklist of Scandinavian and British species, within a new framework of two genera: *Nepticula* and *Trifurcula*, the latter embracing all of Beirne's genera. He also introduced the use of species groups. At about the same time Alfred Borkowksi wrote a number of papers on the faunistics and taxonomy of Nepticulidae in Poland (Borkowski 1969; 1970a; b; 1972a; b). In one of these, he introduced a new generic classification, following partly Beirne (1945), using male genitalia, and combined this with the study of venation (Borkowski 1972a). In Britain Arthur Maitland Emmet (1908–2001) (Harley 2001) (Fig. 13) started to work on this group and many other Microlepidoptera for the new series "The Moths and Butterflies of Great Britain and Ireland" (MBGBI). He followed upon Johansson's and Borkowski's work and treated the Nepticulidae in the first volume (Emmet 1976), but his many shorter and longer papers in The Entomologist's Record, often made in preparation of the book, illustrated his in-depth knowledge, and are also enjoyable reading (for example: Emmet 1973a; b; c; 1974a; d; e; f; g). Also in the 1970's, Josef Klimesch, now retired, started working on the material he collected in southern Europe and the Canary Islands, and described a number of new species from that area (Klimesch 1975c; 1977; 1978b).

Before 1970, most work on Nepticulidae was carried out independently by either amateur entomologists or isolated professional biologists, but during the 1970's scientific interest in taxonomy was boosted by funding and for the first time University groups started to work on the taxonomy and phylogeny of the family. Christopher Wilkinson (1936–2010) (Figs 12, 14) developed an interest in the Nepticulidae during a sabbatical at the Canadian National Collection of Insects, Arachnids and Nematodes in Ottawa in the early 1970's. Bryan P. Beirne (see above), then an applied entomologist in Burnaby, British Columbia, encouraged Wilkinson to revise this group for Canada and the USA. Wilkinson brought this research back to Portsmouth Polytechnic University, where he had worked since 1965, and two of his graduate students (Philip J. Newton and Malcolm J. Scoble) worked jointly with him on North American Nepticulidae, resulting in five papers dealing with all known species and several new ones (Wilkinson 1979; Wilkinson and Scoble 1979; Wilkinson 1981; Wilkinson and Newton 1981; Newton and Wilkinson 1982). Later, Malcolm Scoble (Fig. 17) obtained an academic position at the Transvaal Museum in Pretoria and continued working on the Nepticulidae of South Africa in a global context, gaining his PhD in 1983 (Scoble 1978a; b; 1979; 1980a; b; 1982; 1983). He presented the first cladistic analysis and classification of the family, dividing the family into two subfamilies, Pectinivalvinae and Nepticulinae, and the latter into two tribes, Nepticulini and Trifurculini. He also introduced the use of subgenera in the genus *Ectoedemia* and he named no fewer than 73 species. After these

studies, Scoble turned his attention to other microlepidopterans, larger Lepidoptera and management while working, respectively, in the Oxford University Museum of Natural History and the Natural History Museum, London.

Christopher Wilkinson became professor of Animal Systematics and Zoogeography at the Free University of Amsterdam in 1977, and employed Erik J. van Nieukerken (Figs 15, 20) as PhD student and Georgina Bryan, Steph B. J. Menken and later Jacobus J. (Koos) Boomsma as postdocs, all to work on various aspects of the systematics and evolution of Nepticulidae. In Amsterdam, several MSc students worked with Wilkinson on the taxonomy of *Stigmella* species groups in Europe, and the taxonomy of *Stigmella* in Japan and New Zealand (Kemperman et al. 1985; Schoorl et al. 1985; Schoorl and Wilkinson 1986; Donner and Wilkinson 1989). The nepticulid research group (Wilkinson 1982) also published a joint paper on the *Ectoedemia angulifasciella* complex (Wilkinson et al. 1983). Unfortunately, the department was short lived being closed down after budget cuts in 1985, cuts that were particularly detrimental to taxonomy. EvN finished his degree in 1986 after the group had been disbanded (van Nieukerken 1983b; 1985b; 1986b; van Nieukerken and Dop 1987). His thesis built upon Scoble's results, but mainly based on the Holarctic fauna. He largely kept Scoble's classification, added two more subgenera to *Ectoedemia*, and included larval and detailed new adult characters in the character matrix. Together with his MSc student, Henk Dop, he also extensively studied antennal ultrastructure. Van Nieukerken and Dop discovered a novel sensillum, which they named sensillum vesiculocladum (van Nieukerken and Dop 1987). The work led to a first checklist of the Western Palearctic species (van Nieukerken 1986a). Steph Menken's research adopted new methods in the systematics of Nepticulidae by using allozyme characters, not only for species discrimination, but also for studies of population structure and parthenogenesis. Long before the advent of DNA barcoding, studies of allozymes facilitated species identification (Menken and Brouwer 1984; Menken and Wiebosch Steeman 1988; Menken 1990). After these studies, Steph Menken continued working on insect host relationships, particularly on Yponomeutidae as professor at the University of Amsterdam and Koos Boomsma continued his ecological studies, working as a professor in Copenhagen, particularly on social insects. The collaboration started by Menken, Boomsma and van Nieukerken resulted in a review paper published in 2010 on diet breadth in Lepidoptera, which also included important information on the host plants of Nepticulidae (Menken et al. 2010).

Wilkinson had also initiated collaboration to study Chinese Nepticulidae with professor Liu Youqiao from Beijing (Academia Sinica), resulting in a joint collecting trip by Erik van Nieukerken and Hans van Driel to China with Chinese counterparts in 1984 (van Driel and van Nieukerken 1985). After closing of the Amsterdam department, this research was unfortunately not continued. Only much later, van Nieukerken picked up the collaboration with Liu Youqiao and completed one joint paper on Chinese *Stigmella* (van Nieukerken and Liu 2000). Also plans by Wilkinson and Donald R. Davis (Fig. 11) for a volume in the series MONA (Moths of North America north of Mexico) never materialized, but checklists for the North American (Davis and Wilkinson 1983) and Neotropical Nepticulidae (Davis 1984) were completed.

At the same time that the Amsterdam group was active, Rimantas Puplesis (Fig. 15) started his PhD at the Zoological Institute of the Academy of Sciences in Leningrad,

Figures 17–23. Specialists on Nepticuloidea in 20th and 21st century. **17** Malcolm Scoble, April 1992, Helsinki, Congress SEL **18** Robert J. B. Hoare, collecting *Menurella quintiniae* at type locality, Lamington Park, Queensland, August 2004 **19, 20** Camiel Doorenweerd and Erik J. van Nieukerken collecting leafminers on *Fagus hayatae*, Taiwan, Taipingshan, October 2012 **21** Nagao Hirano and Toshiya Hirowatari, at Hirano's house, Matsumoto, Japan, September 2014 **22** Arūnas Diškus **23** Andrius Remeikis. Photos by Erik J. van Nieukerken (17, 18, 21) and Shipher Wu (19, 20).

Soviet Union, and this resulted in many papers describing new species of Nepticulidae that he and colleagues had collected in Far East Russia (Primorskij Kraj) and Central Asia (Puplesis 1984a; b; c; e; 1985b; c; Puplesis and Ivinskis 1985). He also produced a phylogeny, although not strictly following Hennigian cladistics. From 1985, Rimantas

continued his work in Vilnius (Lithuania), Pedagogical University and while his earlier papers were all in the Russian language, from 1988 he published mostly in English, and an important summary of all his earlier work was a book dealing with the Nepticulidae of the former Soviet Union, including 21 new species (Puplesis 1994). He continued his work first in Central Asia, where he had travelled extensively, usually together with his colleague Arūnas Diškus (Fig. 22), who received his PhD in 2005. From 2000 onwards their main interest shifted to the Neotropical fauna, and they travelled and collected particularly in Belize and Ecuador, and studied also the very interesting material earlier collected by Ebbe Nielsen and Ole Karsholt in Argentina and Chile (Nielsen 1980; Nielsen 1985). Puplesis made the first revisions of the Neotropic fauna in collaboration with Gaden S. Robinson (1949–2009) (Beccaloni et al. 2009) from the Natural History Museum in London (Puplesis and Robinson 2000; Puplesis et al. 2002a; 2002b). In 2003, Diškus and Puplesis published a book (Puplesis and Diškus 2003b) – partly in Lithuanian, partly in English – with several revisions of Nepticulidae from Asia and Europe and a first comprehensive global catalogue including both the Nepticuloidea and Tischerioidea (Diškus and Puplesis 2003). In 1996 Rimantas had become professor in Vilnius, and from 2007 onwards published under the name Jonas Rimantas Stonis. With a group of co-workers, including Arūnas Diškus (Fig. 22), Asta Navickaite, Agnė Rocienė and Andrius Remeikis (Fig. 23) he continued working on the Neotropical fauna and collecting, now also in Guatemala, Colombia, Mexico, Peru, Bolivia and Chile (Stonis et al. 2013c; 2013d; 2014; 2016b; Remeikis and Stonis 2015; Stonis and Remeikis 2015), but also revised the earlier described Far East Russian Nepticulidae (Rocienė and Stonis 2013; Stonis and Rocienė 2013) and worked on the Lithuanian and Crimean fauna (Navickaitė et al. 2011; 2014). Puplesis-Stonis and his co-authors have contributed by far the largest number of new species in Nepticulidae, by July 2016 in total ca 225 valid names.

In 1986 Erik van Nieukerken obtained a position at the Rijksmuseum van Natuurlijke Historie in Leiden (now Naturalis Biodiversity Center), but his initial responsibilities left him little opportunity to continue taxonomic work before he got a position as curator for microlepidoptera and Arachnida in 1999. Still he was able to finish some studies that he had started in Amsterdam, and continued collaborations with especially Roland Johansson and other Scandinavians: Ebbe Schmidt Nielsen (1950–2001) (Fig. 14), Bert Gustafsson (Fig. 14) and Ole Karsholt. This collaboration led to the two volume set in the series Fauna Entomologica Scandinavica (Johansson et al. 1990), very beautifully illustrated with water colours and line art by Roland Johansson (for examples see Figs 25–40) and also containing the first treatment of all nepticulid larvae of a single fauna (Gustafsson and van Nieukerken 1990). Nielsen, who had moved from Denmark to Canberra, Australia, had been the driving force for completion of this multi-authored work, and he was that again for the completion of the study of the previously described Australian Nepticulidae by him, Johansson and van Nieukerken. It was speeded up when Nielsen had been able to hire in 1994 Robert J. B. Hoare (Fig. 18) as PhD student to work on Australian Nepticulidae (Hoare et al. 1997). Hoare finished his study of Australian Nepticulidae in an unpublished PhD thesis (Hoare 1998), of which most relevant parts have since been published, including

the description of the new genus *Roscidotoga* and a revision of *Pectinivalva*, including two subgenera (Hoare 2000a; Hoare 2000b; Hoare and van Nieukerken 2013), that are raised here to genus. After moving to Auckland, New Zealand, Hoare shifted his attention to the Lepidoptera of New Zealand in general.

In the late 1980's and early 1990's the Czech brothers Aleš and Zdeněk Laštůvka (Fig. 16) developed an interest in leafmining Lepidoptera, particularly Nepticulidae and Gracillariidae, and the opening of the Iron Curtain provided them with the opportunity to start extensive collecting in the Mediterranean region: Spain, Portugal, France, Italy, Croatia and Greece. They published a general guide for Central European species (Laštůvka and Laštůvka 1997) and many descriptions of new species (e.g. Laštůvka and Laštůvka 1990; 1994; 1998; 2000b; 2007). They also extensively collaborate with van Nieukerken on this fauna, resulting in several of the papers cited below. Aleš Laštůvka also started to illustrate the moths with water colours, as can be seen in several of their papers.

Moving into the 21st century, Roland Johansson and Erik van Nieukerken renewed their collaboration, resulting in a second revision of the oak mining *Stigmella* (van Nieukerken and Johansson 2003). The initiation of a molecular laboratory at Naturalis prompted van Nieukerken to include sampling material for DNA analysis during his extensive collecting in many parts of the world, including Vietnam, Australia, Borneo and from 2010 onwards in the USA, Taiwan, Korea and Japan. Early molecular results in 2004 were shown at the International Entomological Congress in Brisbane (van Nieukerken et al. 2004a), but still considered insufficient for a full publication. Most publications by him and co-authors in the early 2000's concern detailed European faunistics (van Nieukerken et al. 2004b; 2004c; 2006; van Nieukerken 2006) and two small and one large revision of West Palearctic *Acalyptris*, the subgenus *Trifurcula* (*Levarchama*) and *Ectoedemia*, subgenera *Ectoedemia* and *Zimmermannia* (van Nieukerken 2007a; b; van Nieukerken et al. 2010). Meanwhile in Naturalis the DNA barcoding project was adopted, and extracting DNA for barcoding (Ratnasingham and Hebert 2007) became a routine during dissection. Several papers provided barcodes as support for the taxonomy (van Nieukerken 2007a; 2010; Ivinskis et al. 2012; Laštůvka et al. 2013), and with MSc student Camiel Doorenweerd (Fig. 19), they published two larger papers concentrating on DNA barcodes and how to use them in Nepticulidae (van Nieukerken et al. 2012a; 2012b). Camiel Doorenweerd continued as PhD student from 2012 and analysed more genes. This resulted in several phylogeny papers (Doorenweerd et al. 2015a; Doorenweerd et al. 2016a), and some that are still in preparation. In order to improve calibration points for molecular phylogenies, all Nepticulidae fossils were re-assessed and catalogued (Doorenweerd et al. 2015b), and the named ones with formal descriptions are also included in the present catalogue.

After a paper on *Stigmella* from Japan (Kemperman et al. 1985), a collaboration between the Free University of Amsterdam group of Wilkinson and two Japanese leafminer specialists: Hiroshi Kuroko and Tosio Kumata, work on that fauna had slowed down, but a few species were named or discussed (Kuroko 1989; 1990; Kumata and Nakatani 1995; Kuroko 1999; Kuroko 2004; van Nieukerken and

Kuroko 2005). The collaboration with Japan is now picked up again by van Nieukerken and Doorenweerd with Toshiya Hirowatari (Fig. 21), first in Osaka University, now professor in Kyushu University in Fukuoka and is expected to soon yield new publications. The study received further impetus from an amateur, Nagao Hirano (Fig. 21), who had collected and reared Nepticulidae extensively and described 12 new species (Hirano 2010; 2014); he also wrote the Nepticulidae texts for the book series "The standard of moths in Japan", written in Japanese (Hirano 2013). At the same time some small-scale collaborations with China (prof. Li Houhun) and Korea (Bong Woo Lee) resulted in new material for study and will hopefully result in new publications.

Also work on North American Nepticulidae is now being continued collaboratively by van Nieukerken, Doorenweerd, in collaboration with Davis, David Wagner, Charley Eiseman, Greg Pohl and others, and this includes extensive DNA barcoding studies. A checklist of Lepidoptera of Canada with our contribution is now in preparation (Greg Pohl and Jean-François Landry, editors) and some of our data are presented in this catalogue.

Taxonomic history of Opostegidae

The first opostegids to be named were *Tinea auritella* Hübner, 1813 (now *Pseudopostega auritella*) and *Elachista salaciella* Treitschke, 1833 (now *Opostega salaciella*). Zeller (1839) erected the genus *Opostega* for these two species and his new *O. crepusculella*, but he also included other small white species, that are now included in respectively *Phyllocnistis* Zeller, 1848 (Gracillariidae) and *Leucoptera* Hübner, 1825 (Lyonetiidae). Zeller did not select a type species, only much later Walsingham (1914) selected *salaciella* as such. In 1848 Zeller narrowed *Opostega* and only included species now still regarded as Opostegidae, by removing the other species to his new genera *Phyllocnistis* and *Cemiostoma* Zeller, 1848 (a junior synonym of *Leucoptera*). In 1855 five out of the six known European species had been named and two species from North America followed in the next 20 years (Clemens 1862a; Chambers 1875b).

Most early authors placed *Opostega* in the family Lyonetiidae (Stainton 1854; Frey 1856; Stainton 1859a), but Heinemann and Wocke ([1876]) already recognised the similarity with *Nepticula* and placed them in the Nepticulidae. The family name was first used by Meyrick (1893) as Opostegides, but Meyrick considered it to be a subfamily of Tineidae.

After the early descriptions, for a long period new species, mostly from other continents, were almost all named by Edward Meyrick and Lord Walsingham (66 in all) (Table 3). Just single species are attributed to Busck, Chrétien, Eyer, and Kuroko. A. Jefferis Turner (1923) described some species from Australia and O.H. Swezey (1921) discovered that the opostegids in Hawaii (now *Paralopostega*) make leafmines on *Melicope* (Rutaceae) (then named *Pelea*), whereas the life history of most Opostegidae remained a mystery. Only that of *Opostegoides scioterma* had been extensively described by Grossenbacher (1910), under the incorrect name *Opostega nonstrigella* (Davis and

Table 3. Number of Opostegidae species (valid and invalid) described per first author for authors who described at least six valid species.

First Author	# spp
Davis	74
Meyrick	53
Puplesis (Stonis)	21
Walsingham	11
Turner	6
17 authors	27
Total	**192**

Stonis 2007). This species is a cambium miner of *Ribes* (Grossulariaceae), and another species of the same genus, *Opostegoides minodensis* (Kuroko, 1982) was later discovered to be a cambium miner of *Betula*.

In the 20th century there was still no agreement about placement of the single genus *Opostega*, some textbooks placed it in the Nepticulidae, sometimes as subfamily (Spuler and Meess 1910; Hering 1932a), others kept it in Lyonetiidae, including the book first describing the male genitalia of three British species (Pierce and Metcalfe 1935). The first paper that recognised a grouping of families very similar to what we know nowadays was Busck (1914b), who considered the Opostegidae as family, close to Nepticulidae and Tischeriidae, in the Lepidoptera "Aculeates", rather similar to what we now call lower Heteroneura. This placement was confirmed by larval studies a few years later (Heinrich 1918).

When studying the Opostegidae from the Asian parts of the Soviet Union, Kozlov (1985) realised that the differences in the genitalia of the single recognised genus *Opostega* were large and he erected a new genus, *Opostegoides* Kozlov, and split *Opostega* in two subgenera: *Opostega* and *Pseudopostega* Kozlov. Soon thereafter Davis (1989) completed a generic revision of the family and a phylogenetic analysis, finally resulting in the recognition of three new genera and raising *Pseudopostega* to full genus. The most peculiar new genus was *Notiopostega*, with the single species *N. atrata* Davis, 1989, with a wingspan of 13–18 mm by far the largest opostegid and largest species of the superfamily, of which the larvae make extremely long mines of up to seven meters in the cambium of *Nothofagus* trees (Carey et al. 1978; Davis 1989).

The four Northern European Opostegidae were reviewed and keyed by van Nieukerken (1990a), and a key for all six European species was presented by van Nieukerken et al. (2004b). Erik van Nieukerken also discovered and confirmed host plants for the northern European *Pseudopostega* species: *Lycopus europaeus* (Lamiaceae) for *P. auritella* and *Mentha aquatica* (Lamiaceae) for *P. crepusculella.* Both make long linear mines in the bark and gradually go deeper in the stem (van Nieukerken 1990a; Regier et al. 2015).

Rimantas Puplesis also worked extensively on Opostegidae, and revised the Oriental species together with Gaden Robinson (Puplesis and Robinson 1999) and together with Donald Davis they revised the New World species, including 70 new species and one new genus (Davis and Stonis 2007).

Material and methods

Taxonomic practice

Species

Ultimately, a catalogue or checklist is a list of species, arranged in a linear classification framework. Few taxonomists describe the methodology and philosophy they follow to recognise and delimit species. They often use the simple species concept: "a species is what a taxonomist calls a species" (Kalkman 1987). There are many species concepts, see eg Mallet (2001) for an overview. If we have to choose, we probably adhere most to the phylogenetic or diagnostic species concept (Mallet 2001; Isaac et al. 2004), but agree with Mallet's notion that "agreement on a unified species-level taxonomy is possible, but will be forthcoming only if we accept that species lack a single, interpretable biological reality over their geographic range and across geological time". In the taxonomic literature on Nepticuloidea, only Scoble (1983) discussed how he recognised species, but also he concluded that the practice is rather different from the theory. With the explosion of genetic data we have much more information nowadays than just two decades ago, but that does not mean that it has become much easier to recognise species. DNA barcodes are helpful, but can also complicate species discrimination, especially for allopatric populations (see below). In practice we recognise species by a combination of morphological characters, where genitalia are important, but not the only characters, and of the biology: host plant and mine morphology and of DNA data, particularly DNA barcodes and also of distribution data.

All species recognised here are in fact just hypotheses of species, open to further testing by more data and subject to ever ongoing evolution. We change the status of previously recognised species whenever our own research or our interpretation of published research has given reason to do so. This is particularly the case for several North American and Asian species that we (EvN and CD) have been studying the last years, and for which several publications are in preparation.

Subspecies

We do not recognise any subspecies. The systematic category in itself is problematic, and is particularly a part of the polytypic biological species concept, that can only be used for species where distributions are known in detail and the amount of hybridisation in border areas has been studied; as such subspecies are mostly used in charismatic groups such as vertebrates and butterflies, and even in these there is a tendency given by the phylogenetic species concept to abandon subspecies and raise them often to full species, particularly in birds (Isaac et al. 2004).

If allopatric populations are morphologically (almost) inseparable, and also share most of the biology, we simply use the same species name over the entire area (e.g. *Ectoedemia occultella* in the entire Palearctic and Nearctic, making the same character-

istic mines on *Betula* and morphologically indistinguishable). In the case that there are more differences, we opt for separate species (e.g. *E. intimella* and *E. insularis*), particularly when consistent morphological differences are paired with a large diagnostic difference in DNA barcodes.

Very few subspecies have been described in the last 50 years in Nepticuloidea; for the few that were named (eg in two species of Neotropical *Pseudopostega*, (Davis and Stonis 2007)) we have raised the subspecies to full species when we see sufficient differences, or just left them as synonyms in other cases (e.g. *Stigmella anomalella pacifica* Puplesis, 1987). More than 50 years ago, the subspecies category had been used sometimes for biological forms (host plant races) and these are treated either as synonyms or raised to species level in a few cases.

Higher categories

Our classification of Nepticulidae follows our molecular phylogeny (Doorenweerd et al. 2016a), of which Fig. 24A shows the results in summary. Since the earlier recognised subfamily Nepticulinae and the tribe Nepticulini are not recovered, we abandon the use of subfamily and tribus here. After consultation with various lepidopterists we also choose to drop the subgenus category entirely, after we had to abandon them in *Ectoedemia* to maintain manageable monophyletic entities. In this way our classification resembles that of Diškus and Puplesis (2003).

The genus *Enteucha* is sister to all other Nepticulidae and therefore listed first. *Stigmella* is always split into two large clades that we name "Core *Stigmella*" for the clade with the type species *S. anomalella* and "Non-core *Stigmella*" for the other one. Splitting *Stigmella* was considered impractical, due to a lack of good morphological apomorphies for these clades and the fact that *Stigmella* remains a well supported and recognisable entity.

The new Neotropical genus *Ozadelpha* (van Nieukerken et al. 2016) forms a clade with the former Pectinivalvinae. Also in *Pectinivalva* we have raised the three recently recognised subgenera to full genus (Hoare and van Nieukerken 2013).

This clade is sistergroup to the remaining genera, previously collectively named Trifurculini. Within this clade one can recognise three groups: A poorly supported clade of *Bohemannia* together with the new Neotropic genera *Neotrifurcula* and *Hesperolyra*, a clade of *Glaucolepis* + *Trifurcula* and a clade of the former *Ectoedemia* plus *Acalyptris* and *Parafomoria*. Following the principle of abandoning subgenera, we split *Trifurcula* now into *Trifurcula* and *Glaucolepis*, whereas *Levarchama* is reduced to a species group in *Trifurcula*.

Whereas the finding of the large "*Ectoedemia* clade" occurs in all our analyses, the order of the various genera changes in different analyses. The classification we adopt here follows the best supported phylogeny. The raising of *Muhabbetana* and *Zimmermannia* to full genus is novel, even though Hering (1940) already proposed *Zimmermannia* as full genus, this was never adopted until now.

The classification of Opostegidae follows the treatments by Davis (1989) and Davis and Stonis (2007), but also here we abandon the use of subfamilies, and we do not follow the splitting of *Pseudopostega* in species groups (see below).

Species groups

Species groups are a practical category to combine groups of species within large genera that share morphological and biological characters, and have been extensively used in Nepticulidae, particularly *Stigmella* since Johansson (1971). Our molecular studies show that many of these species groups are indeed monophyletic entities (Doorenweerd et al. 2015a; 2016a), but on the other hand there has also been a proliferation of often monotypic species groups for species with different genitalia that cannot be placed easily. We have here recognised especially the well diagnosable species groups that often are also supported by molecular data. These include many of the Holarctic groups in *Stigmella* and *Ectoedemia*. Groups for which we have no or insufficient molecular data have been recognised when morphologically uniform and comprising more than one species, others are listed as unplaced within genera, or the larger clades that we recognise in *Stigmella*. When several groups together form a monophyletic clade, we sometimes use the term "cluster" for that clade.

In *Pseudopostega* we have abandoned all species groups for the time being, and list the species alphabetically by geographic region. We have been unable to find phylogenetic or molecular support for the groupings and found them hard to use. For details for these groups we refer to the revisions that introduced them (Puplesis and Robinson 1999; Davis and Stonis 2007).

Species group names are not governed by the ICZN rules, and for practical reasons we have therefore changed some names that were based on junior synonyms, or in one case replaced it by the name of the type species that is included in the group (*Acalyptris psammophricta* group rather than *repeteki* group). We give the authors who used the group name for the first time, even though they may have used a different composition. Synonyms are not complete, and many group names for single species are not given in synonymy, nor are the many names for *Pseudopostega* groups.

We group a few sibling species in species complexes, but only in cases where these species are really very hard to almost impossible to separate morphologically, or only by either biological or molecular data. This applies only to three complexes in European *Ectoedemia* (van Nieukerken et al. 2012b). Recently, Stonis and Remeikis (2015) recognised two species complexes in Neotropical *Acalyptris*, but since these are simply groupings of rather similar, but diagnosable species, we do not concur to use the term "complex" for such assemblages, because there are multiple similar examples in global Nepticuloidea. The *Stigmella nigriverticella* complex (Remeikis and Stonis 2015) is indeed a complicated group, but the *saginella* group to which the complex belongs, is overall still a taxonomic puzzle that requires more study both morphologically and genetically.

Order of the list

We order the genera and species groups according to our preferred phylogeny (Doorenweerd et al. 2016a), see also above. Species groups for which we do not have molecular data are listed at the end of each genus, monotypic species groups are largely abandoned, except in cases with an almost fully resolved phylogeny, where a single species does not group with any species group, as in the *Ectoedemia terebinthivora* group (Doorenweerd et al. 2015a), or when we know unnamed species that clearly group with the single species. The form genus *Stigmellites* for unplaced fossils is listed at the end of Nepticulidae and dubious taxa are listed at the end of each family. At the end of the checklist we list excluded taxa with their current taxonomic placement. These taxa were either originally described in *Nepticula* or *Opostega*, or once combined with these genera, but are now considered to belong to other families, in all 29 names, now belonging to 23 species in ca. eight to ten families. The list also includes four taxa of fossil leafmines, previously considered to possibly belong to Nepticulidae (Doorenweerd et al. 2015b).

Species are only listed in a phylogenetic order for those genera and groups where we have a detailed phylogeny, in practice only in *Ectoedemia* and a few species groups in *Stigmella*. Otherwise species are grouped by geographical region (order: WP, EP, OR, AFR, AUS, NEA, NEO) and listed alphabetically. Following the valid name we list the original combination plus those of the junior synonyms, followed by all now invalid subsequent combinations. Usually unavailable names are given at the end of the list of synonyms.

Distribution

For each species we give the biogeographical region(s) of occurrence with an abbreviation. When the abbreviation is placed in square brackets the species is assumed to be an introduction. For type localities and simple maps of country records we refer to the scratchpad http://nepticuloidea.info/, which is regularly updated, and the Catalogue of Life.

DNA barcodes and specimen data

Obtaining DNA barcodes has been standard in our taxonomic workflow since ca 2000, particularly for recent material or taxonomic relevant material such as types; in many cases additional genes have been sequenced. Our methodology has been explained in detail elsewhere (van Nieukerken et al. 2012a; 2012b; Doorenweerd et al. 2015a; 2016a). Where our classification in the first place is based on the phylogenetic analyses of several genes, we have used DNA barcodes often to place species that were not involved in those analyses. Further we used the barcodes as additional arguments in decid-

ing about species status, even though there is no absolute criterion. The judgement of the value of barcode distances is particularly problematic in vicariant populations, such as island populations, where exchange of genetic material has ceased often a long time ago (Mutanen et al. 2012; 2016). We only opted for separate species when a large distance was paired with morphological and/or biological differences, shown in sufficient material. In cases when single specimens on an island show large Barcode distances, but hardly any morphological ones, we usually decide against splitting until further material and data are available. This is for instance the case in *Pseudopostega chalcopepla* and some island populations of *Stigmella perpygmaeella*. We have discussed the use of barcodes extensively for *Ectoedemia* and some groups of *Stigmella* (van Nieukerken et al. 2012a; 2012b). Barcodes have also been of great use to couple the unknown sex, eg. in some *Zimmermannia* species, or to link species with unknown life histories to barcoded larvae. We would like to stress the added value of DNA barcodes to taxonomy, and plea for adding barcode data to all taxonomic treatments, whenever possible.

We release here our dataset of barcodes of well identified material as BOLD Dataset DS-NEPCAT (DOI: 10.5883/DS-NEPCAT). This means that identities of specimens included were either identified by us or by colleagues providing the data. Obviously various barcoded larvae that did not match any adult barcodes and could not otherwise be identified may still belong to named species with unknown hosts. This dataset includes DNA barcodes of 3205 specimens, belonging to 779 species (733 Nepticulidae and 46 Opostegidae) of which 2574 specimens belong to 444 formally named species (409 Nepticulidae and 35 species of Opostegidae). The remaining 642 specimens belong to ca. 335 unnamed (or as yet unidentified) species (324 Nepticulidae and 11 Opostegidae). In the dataset 3071 barcodes are assigned a Barcode Identification Number (BIN), representing 900 BIN's, which belong to 749 recognised species (717 Nepticulidae and 32 Opostegidae). A neighbour joining tree of the barcode data is supplied here as Supplementary material S2. Even though NJ trees contain phylogenetic signal, they cannot be considered as real phylogenetic trees, and species may be completely misplaced on the basis of barcodes alone.

Data for specimens to which we refer in our notes are listed in Supplementary material S1, and when barcoded these data are also available in dataset DS-NEPCAT.

Host plants

We refrain from compiling a list of host plants here, even though it would have made the catalogue much more complete. A critical catalogue of host plant records requires considerable work on checking literature records, verifying host plant identifications and nomenclature and interpretation. We are working on such a catalogue to publish in the future, including many records of yet unnamed and unidentified species, to provide an insight into the host plant choices of the family. A catalogue of hosts without reference to sources was given by Diškus and Puplesis (2003), and for the West Palearctic and northern European alone by respectively van Nieukerken (1986a) and

Johansson et al. (1990) and several more local revisions provide detailed host records. An analysis of all global host records on family level was used for the paper on the evolution of host associations in Lepidoptera (Menken et al. 2010).

Nomenclatorial practice

The most important source for nomenclatorial practice is the International Code of Zoological Nomenclature (International Commission on Zoological Nomenclature 1999), that helps solving most nomenclatorial issues. However, interpreting the code is often difficult and we therefore highlight how we dealt with some cases in general, whereas in notes we discuss our individual choices where needed.

We include only names that were published as real scientific names, all available names according to ICZN, but also all unavailable names that are formed as real binomial or trinomial names. Unavailable names are marked with a double dagger (‡). For completeness' sake we also include infrasubspecific names, that are not available according to the Code (ICZN 1.3.4). Many names based on leafmines only (the work of an animal) are unavailable when published after 1930 (ICZN 13.6.2), these are marked with the abbreviation "NNLM" (see below). After screening some of these descriptions, they appeared to be available after all, since part of the description dealt with the larva, eg. the colour, thus making the description available, even though it often is of little use in recognising the species.

We do not include any informal intermittent names (eg *Acalyptris* species 29135) or names given to unnamed barcoded species, as such names easily become obsolete and often are not published on paper or another fixed media anyway. Some of these intermittent names are listed on our scratchpad, though (van Nieukerken 2016) and many are used in the DNA barcodes dataset.

References

All original descriptions and references with new combinations and synonymies have been examined by EvN (mostly from his own collection of reprints, copies and pdf's), always trying to establish correct publication dates, which lead to changing a few publication dates of taxa. The Biodiversity Heritage Library (BHL) has been a particular great help, since this contains fully scanned volumes, allowing checking of dates on wrappers of issues and volumes, in addition to those available in our libraries. Nowadays there are many more repositories with scanned journals and books of particular countries, but since they do not always contain scans of wrappers and title pages, this checking is not always easy. It would be a great improvement if all these repositories were accessible through a single online source (BHL?); now it is not always easy to find these.

In the catalogue proper we present citations in the form as recommended by the Code (ICZN art. 50, 51, recommendations 51B–G), thus with comma, and use of

ampersand rather than "and". We always cite the page for the first description and for other nomenclatorial actions.

All references are listed here, and all references for original descriptions are also available in the scratchpad (van Nieukerken 2016). Where possible we added here the doi or url to the references.

Combinations

We here introduce several new combinations, that became necessary after raising subgenera to full genus. While doing this, we realised that it is often difficult to find where new combinations have been made in the past, since ICZN (in contrast to the Botanical Code) does not have special requirements for new combinations, other than that the publication fulfils availability. In practice many new combinations have been made unintentionally in checklists, faunistic papers, faunas, without marking these, whereas several of the new combinations marked as such in literature (including some of our own) appeared not to have been new at all. In Nepticulidae this is particularly the case with the combinations of many species with *Stigmella*: after the recognition by Walsingham (1908a) that in fact *Stigmella* is valid rather than *Nepticula*, some authors started to use this generic name, but many did not, and this uncertainty continued for 70 years (Wilkinson 1978)! There are in fact not many "official" new combinations in *Stigmella* and for European species many were given without further notice in two faunal works (Gerasimov 1952; Hering 1957). To document the combination history for all names in Nepticuloidea, we here give the first author using the combination – as far as we have been able to ascertain – as author name after the brackets, in a similar fashion as botanists do and as has been recommended by ICZN (Recommendation 51G). Combinations only published online on webpages or databases (including the scratchpad) and not in online journals, are not accepted as validly published, and thus are given here as new combinations when valid. When these combinations are no longer valid and have been used only online, we mention this by adding "online comb."

One deviation from the code is that we do not change the ending of species-group names to agree with the gender of the generic name (ICZN article 34.2). This follows the practice by most lepidopterists in leading catalogues and checklists, duly discussed by Sommerer (2002) and formally adopted by the Societas Europaea Lepidopterologica in a Resolution during their General Meeting at the 13th European Congress of Lepidopterology in Kørsør (Denmark) on June 4, 2002 (Sommerer *loc. cit.*). In Nepticuloidea this affects only few names anyway, since most generic names are feminine (except *Varius*, which is masculine).

Synonyms

Similar to the combinations, finding the source for new synonyms is often difficult, and we therefore give the source for all subjective synonymies that we have been able

to find. The few cases where we did not find it we leave this open, and we do not give a synonymy author for objective synonyms (names with the same type, such as new names, incorrect subsequent spellings) or for (infra)subspecific names within the same nominal species.

Types

We have tried to include information on the primary types for each name in the scratchpad (van Nieukerken 2016), including information on the depository, the genitalia slide number and the host plant of the type – when reared. Information on types is provided here only for cases discussed in the notes. We also started adding photos of types to the scratchpad, but this will be far from complete when this paper is published.

Type species of genera and type genera of family group names are always given in the list, using several similar abbreviations as suggested by Pullen et al. (2014).

Abbreviations and symbols used

AFR	Afrotropical region
AUS	Australian and Pacific regions
BIN	Barcode Identification number
BOLD	Barcoding of Life Database
EP	East Palearctic region
ICZN	International Code of Zoological Nomenclature (International Commission on Zoological Nomenclature 1999)
IOS	Incorrect original spelling
ISS	Incorrect subsequent spelling
JH	Junior Homonym [of Genus]
JPH	Junior Primary Homonym [of Species]
JSH	Junior Secondary Homonym [of Species]
ND	Nomen dubium, dubious name, of which identity is unknown and untraceable
NEA	Nearctic region
NEO	Neotropical region
NN	Nomen nudum, unavailable name failing to conform to ICZN art 12 or 13
NNLM	Nomen nudum, for a name after 1931 based on the description of a leafmine (ICZN 13.6.2)
NO	Nomen oblitum, forgotten name, not used as valid name after 1900
OR	Oriental region
RN	Replacement name
syn	Synonymised by
TG	Type Genus

TS/OD	Type Species by Original Designation
TS/OD,M	Type Species by Original Designation and Monotypy
TS/M	Type Species by Monotypy
TS/SD	Type Species by Subsequent Designation
UE	Unjustified emendation
URN	Unnecessary replacement name
WP	West Palearctic region
†	Fossil species
‡	The double dagger is given before an nomenclatorially unavailable name.
[1]	Superscript numbers refer to the taxonomic notes after the list. In the online html version of the list automatically hyperlinked.

Global Nepticuloidea: the state of the art

Table 4 provides an overview of the diversity of global Nepticuloidea for region and genus. Here we report 862 extant named species of Nepticulidae and 192 Opostegidae, a total of 1054. For the 18 fossil species we refer to our earlier catalogue (Doorenweerd et al. 2015b). We illustrate most genera with watercolours by Roland Johansson in Figures 25–40.

There is a strong bias for the West Palearctic with 321 Nepticulidae, but the highest number of Opostegidae is for the Neotropics (89), which probably better reflects the reality. Fig. 24B shows that there is a constant increase in numbers of described species over time, and we are already aware of large numbers of unnamed species from most areas, but particularly the tropics and East Asia. The following lines present a short summary of our current knowledge of Nepticuloidea per region.

Europe. As has also become clear from the taxonomic history, until recently most work concentrated on the European fauna, and probably the majority of species have been described by now. Europe, excluding Cyprus, but including Macaronesia, contains 280 named species. We know of about 20 unnamed species of *Trifurcula*, eight *Parafomoria* and a small number of *Stigmella*, particularly *Rhamnus* feeders and several species belonging to the *Stigmella salicis* complex (van Nieukerken et al. 2012a), but otherwise do not expect many more to be discovered. Several key works deal with parts of Europe (Johansson et al. 1990; Laštůvka and Laštůvka 1997; Bengtsson et al. 2008), but this still excludes southern Europe.

The largest genus is *Stigmella*, as in most regions, but also *Ectoedemia* is very diverse here with a particularly rich fauna associated with oaks (van Nieukerken et al. 2010; 2012b). Both genera feed particularly on Fagaceae, Rosaceae, Betulaceae and Salicaceae, and *Stigmella* also on Rhamnaceae and a few other families. Although Europe has no endemic genera, the three Mediterranean genera *Parafomoria*, *Glaucolepis* and *Trifurcula* have by far their largest diversity here, particularly in the Iberian Peninsula, and to a lesser extent in Italy and Greece. Most of these feed on shrubs and some herbs in the typical Mediterranean habitats as Garrigue or Maquis, with *Parafomoria* specialising

Table 4. Diversity of extant Nepticuloidea per geographic region and globally. Numbers are validly described species. When only unnamed species of a certain genus are known from a region this is indicated by a plus sign, brackets indicate occurrence just at the edge of the region. When generic assignment is uncertain, the number is given in *italics*.

	WP	EP	OR	AUS	AFR	NEA	NEO	Global
Enteucha	1	+	1			(2)	9	**11**
Varius					1			**1**
Simplimorpha	1				1			**2**
Stigmella	138	115	17	31	50	51	61	**428**
Ozadelpha							3	**3**
Roscidotoga				4				**4**
Casanovula				2				**2**
Menurella			1	11				**12**
Pectinivalva				7				**7**
Neotrifurcula							1	**1**
Hesperolyra							4	**4**
Bohemannia	3	5						**7**
Areticulata					1			**1**
Glaucolepis	34	2	1	+		1	*2*	**40**
Trifurcula	34		?		*2*			**36**
Fomoria	12	7	3	3	22	3	*1*	**48**
Muhabbetana	4				28			**32**
Parafomoria	8							**8**
Etainia	7	4	+		4	2		**16**
Acalyptris	23	1	7	+	22	9	32	**93**
Zimmermannia	9	4	+			5	+	**17**
Ectoedemia	47	29	+		5	14	*1*	**89**
Nepticulidae Total	**321**	**167**	**30**	**58**	**136**	**85**	**114**	**862**
# genera	**13**	**8**	**6**	**6**	**10**	**7**	**9**	**22**
Notiopostega							1	**1**
Eosopostega		1	1	+				**2**
Neopostega							6	**6**
Paralopostega				6				**6**
Opostegoides	1	6	15	1	4	1		**28**
Opostega	5	3	(1)					**7**
"Opostega"				17	3			**20**
Pseudopostega	3	2	22	+	8	9	82	**122**
Opostegidae total	**9**	**12**	**39**	**24**	**15**	**10**	**89**	**192**
# genera	**3**	**4**	**3**	**4**	**3**	**2**	**3**	**8**
Nepticuloidea Total	**330**	**179**	**69**	**82**	**151**	**95**	**203**	**1054**
# genera	**16**	**12**	**9**	**10**	**13**	**9**	**12**	**30**

on Cistaceae, *Trifurcula* on Fabaceae, mostly brooms (Genisteae) and Loteae, whereas *Glaucolepis* has groups of species feeding on Lamiaceae, Apiaceae: *Bupleurum* and Plantaginaceae: *Globularia*. *Fomoria* has a centre of diversity in Greece and Turkey with at



Figure 24A. Schematic phylogram of Nepticulidae genera based on the best resolved tree of our Molecular phylogeny (Doorenweerd et al. 2016a).

least six species feeding on *Hypericum*. *Acalyptris* also has most species in South East Europe, with the *staticis* group specialised on Plumbaginaceae and often occurring along the sea coast, and four species in the *platani* group on Platanaceae, Loranthaceae and Anacardiaceae. *Etainia* and *Zimmermannia* are widespread in Europe, the first associated with Sapindaceae (*Acer*) and Ericaceae (*Arctostaphylos*), whereas most *Zimmermannia* are barkminers in Fagaceae, and one in *Ulmus*. *Simplimorpha* has one species, *S. promissa*, oligophagous on Anacardiaceae in southern Europea, whereas the single *Enteucha, E. acetosae* occurs in Central and Eastern Europe on *Rumex* species (Polygonaceae).

Western Palearctic Region. Obviously Europe is part of this region, and its fauna continues along the Mediterranean coasts, but there is a great difference in the know-



Figure 24B. Cumulative number of valid species of Nepticulidae and Opostegidae described per year. Notice the long period that no Opostegidae were described (1935–1967) and the sudden increase in numbers of Nepticulidae since 1978.

ledge of the faunas inside and outside of Europe. The region has only 50 species that are not known from Europe (the total of 330 minus 280). The typical Mediterranean fauna continues in North Africa, Turkey and the Levant, but has been poorly sampled and will probably contain still many new taxa to discover. Turkey and Iran in particular are promising countries for high diversities, Turkey has a large diversity in such host plant genera as *Quercus* and *Hypericum*. In the desert areas of the Middle East, North Africa and the Arabian peninsula other groups become important, such as the *Acalyptris psammophricta* and *shafirkanus* groups (Puplesis et al. 1996; van Nieukerken 2010) and *Ectoedemia* and *Zimmermannia* are almost absent. Probably the best studied area in this region is Turkmenistan with nearly 50 recorded species (Puplesis and Diškus 2003a).

Eastern Palearctic Region. We separate West and East Palearctic more or less following the 64–65 East meridian, from North to South along the rivers Ob, Tobol, Turgay, Aral Sea, Karakum desert and the border between Iran and Afghanistan/Pakistan. In practice this means that we treat in Central Asia Turkmenistan and Iran as West Palearctic, and Afghanistan, Tadzhikistan and most of Uzbekistan and Kazakhstan as East Palearctic. This is a huge area, with major differences between the rather dry and mountainous Central Asia and the almost subtropical forested areas of the Sino-Japanese zone. The 167 named Nepticulidae and 12 Opostegidae certainly only represent the tip of the iceberg. Much work has been done on parts of the former Soviet Union: Central Asia and the Russian Far East (Puplesis 1994; Puplesis and Diškus 2003a; Rocienė and Stonis 2013; Stonis and Rocienė 2013), but for other regions descriptive work is only just beginning. In central Asia the best studied and most diverse country is Tadzhikistan with ca 40 species (Puplesis and Diškus 2003a). In this area



Figures 25–34. Diversity of Nepticulidae, all on same scale. **25** *Enteucha acetosae*, male, Austria **26** *Stigmella mespilicola*, male, Switzerland, holotype **27** *Roscidotoga callicomae*, female paratype, Australia, NSW **28** *Menurella libera*, male holotype, Australia, NSW **29** *Pectinivalva caenodora*, male holotype, Australia, NSW **30** *Glaucolepis lituanica* , male, Austria **31** *Bohemannia auriciliella*, male, The Netherlands **32** *Trifurcula iberica*, male paratype, Spain **33** *Fomoria weaveri*, female, Sweden **34** *Parafomoria helianthemella*, female, Czech Republic. Scale 1 mm. Watercolours by Roland Johansson, 25, 26, 31 and 33 published earlier by Johansson et al. (1990), 28 and 29 by Hoare et al. (1997). The left wings of 32 and 34 are digitally mirrored images of the right wings. These figures may be reproduced given that their author Roland Johansson and the present publication are credited.

Stigmella and *Ectoedemia* are particularly common in the mountainous areas, with species feeding particularly on Rosaceae, Salicaceae and Rhamnaceae, but in desert areas *Acalyptris* becomes an important element with members of the *psammophricta* and *shafirkanus* groups of which biologies are mostly completely unknown.

The northern part of the area, Siberia, has many trans-palearctic species, particularly *Stigmella* and *Ectoedemia* species feeding on *Betula* and Ericaceae feeders as *Stigmella lediella* on *Rhododendron* and *Fomoria weaveri* on *Vaccinium vitis-idaea*. A paper on Siberian nepticulids is in preparation (van Nieukerken, Kirichenko et al.).

Eastern Asia is a very rich faunistic area, with very diverse forests containing many potential host plants. The northernmost portion, Far east Russia, notably the Primorye region, has been best studied, with around 70 species known (Puplesis 1994; Rocienė and Stonis 2013; 2014, Stonis and Rocienė 2013). Initial fieldwork in China has resulted in the recognition of a very rich fauna of at least some 200 species, but to date only few have been described (van Driel and van Nieukerken 1985; van Nieukerken and Liu 2000). For Japan we have a working list of at least 120 species, the recent fauna work listed ca 75 (Hirano 2013; Hirowatari 2013). Obviously all these faunas have a large overlap, and are characterised by dominance of *Ectoedemia* and *Stigmella* (Kemperman et al. 1985; Hirano 2010; 2014). On many tree species there are multiple species in both genera, particularly in Fagaceae, Betulaceae, Juglandaceae, Rosaceae, Ulmaceae and Salicaceae, but also the fauna on Ericaceae and Sapindaceae is comparatively rich. Linking species names from species described from adults only to leafmines from which no adults have emerged is still a challenge, but due to an increasing number of available barcodes more and more of these are linked and publications are being prepared. In *Stigmella* there are also a few species feeding on monocots: the grass genus *Oplismenus* with *S. oplismeniella* (Kemperman et al. 1985) and an unnamed species on *Carex* in Japan. The smaller genera *Etainia*, *Fomoria* and *Bohemannia* are relatively rich in this area, and in the more southern parts *Acalyptris* and *Enteucha* become more important, the latter with a number of species feeding on Polygonaceae, to date all unnamed (van Nieukerken 1986b). In the Opostegidae the genus *Opostegoides* is particularly rich with six named species.

Oriental region. The number of named species for the oriental region is still very low with only 30 Nepticulidae, but the Opostegidae have been revised (Puplesis and Robinson 1999) and number 39 species. Most of the species were described by Meyrick from India, and some from Nepal by Puplesis and Diškus (2003a). From our fieldwork in Vietnam, Borneo and Taiwan, it is clear that there is a rich fauna of leafmining Nepticulidae in this region, but it is a challenge to find sufficient numbers of larvae that can be reared to adults. We have never been very successful with light collecting adults in this region, but some – often poor – material is available in various collections. The genera *Stigmella* and *Acalyptris* are most diverse in this area, *Ectoedemia* is becoming much rarer further away from the Palearctic, but still occurs on *Rubus* from the North as far as Borneo, forming a complex of closely related species. In Borneo we also discovered the first non-Australian species in the genus *Menurella* (formerly in *Pectinivalva*): *M. xenadelpha* on *Syzygium acuminatissimum* (Myrtaceae)



Figures 35–40. Diversity of Nepticulidae and Opostegidae, all on same scale. **35** *Etainia sericopeza*, male, Sweden **36** *Acalyptris platani*, male, Italy **37** *Zimmermannia atrifrontella*, male, Germany **38** *Ectoedemia klimeschi*, male, Austria **39** *Opostega spatulella*, female, Hungary **40** *Pseudopostega crepusculella*, male, Sweden. Scale 1 mm. Watercolours by Roland Johansson, published earlier by Johansson et al. (1990). These figures may be reproduced given that their author Roland Johansson and the present publication are credited.

(Hoare and van Nieukerken 2013). *Stigmella* species have a wide variety of hosts, including tropical groups such as Dipterocarpaceae, Meliaceae, Phyllanthaceae and Moraceae (*Ficus* species), the large majority belonging to non-core *Stigmella*. There

are also species in the *S. betulicola* group feeding on grasses such as *Oplismenus* and Cyperaceae (*Cyperus*): *S. xystodes* (van Nieukerken 2010). *Acalyptris* have an even wider range of hosts throughout the eudicots, most species belonging to the *A. platani* group.

The Opostegidae are characterised by many species in *Opostegoides* and *Pseudopostega*, and single species in *Opostega* and *Eosopostega* (Puplesis and Robinson 1999).

The fauna of northern parts of the Oriental region, Nepal, northern Vietnam, Taiwan, has a much more Palearctic character, often sharing the same host plant genera and comprises for instance also several *Ectoedemia*, the Polygonaceae feeding *Enteucha* species, including the only named one, *E. diplocosma* and *Fomoria* species feeding both on Lamiaceae: *Callicarpa* and *Vitex* and Hypericaceae (van Nieukerken 2008).

Afrotropical Region. The knowledge of the African fauna is very unbalanced, of the 136 named Nepticulidae species the great majority is known from southern Africa, thanks to the diligent rearing work by Lajos Vári, who also named several species (Vári 1955; 1963) and the revisions by Malcolm Scoble (Scoble 1978a; b; 1980a; b; 1983). Further just a few were described from Gambia (Gustafsson 1985) and outside that area just five species have been named; also collections are poor in unidentified material. But even in South Africa much of the diversity is still unknown, and every new collection contains unnamed species. Two monotypic endemic genera *Varius* and *Areticulata* have an uncertain placement and may be synonyms to existing genera. The genera *Simplimorpha* and *Muhabbetana* are near endemic, both also occur in the adjacent Mediterranean region. *Simplimorpha* is specialised on Anacardiaceae, and most African *Muhabbetana* feed on Ebenaceae and Celastraceae. *Fomoria* and *Acalyptris* are relatively diverse genera with broad host ranges. *Stigmella* is the largest genus, as everywhere, but with a dominance of species belonging to non-core *Stigmella*. The *Ectoedemia commiphorella* group has several species feeding on Burseraceae and is possibly sister to the northern hemisphere *Ectoedemia* (Doorenweerd et al. 2015a). Also the group of African *Etainia* species seems to be sister to the Holarctic *Etainia*, but hosts and biology are completely unknown. The two species assigned to *Trifurcula* (Scoble 1980a) may not belong there, a closer study of these species is needed.

The island fauna of the Indian Ocean is still poorly known: two species occur on Aldabra (Seychelles), see below under note 21, and one is named from Madagascar (*Fomoria scobleella*). However, we have seen examples or DNA barcodes of Nepticulidae from Madagascar that show the presence of the genera *Acalyptris*, *Muhabbetana*, *Ectoedemia* and *Stigmella*. No species are known from Réunion or Mauritius.

The African Opostegidae, with 15 named species, have not yet been revised, but we have been able to recombine several "*Opostega*" species here with *Opostegoides* or *Pseudopostega*, the only genera known with certainty from Africa.

Australian Region. Australia proper has a very rich and special fauna with an estimate of about 250 species of Nepticulidae as currently available in collections (Hoare 1998), of which approximately 30 have been named. Australia has four (almost) endemic genera: the small genus *Roscidotoga* (4 species) in the eastern rainforest, specialised on plants in the Oxalidales (Hoare 2000a; van Nieukerken et al. 2011), and *Casanovula*, *Menurella* and *Pectinivalva* with around 160 species, all but one (that feeds on

Quintinia, Paracryphiaceae) feeding on Myrtaceae, with a large number feeding on *Eucalyptus* (Hoare and van Nieukerken 2013). The other large genus is *Stigmella* with an estimated number of at least 80 species, with the most important host families being Rutaceae, Fabaceae (including *Acacia*), Sapindaceae, Euphorbiaceae, Phyllanthaceae and Rhamnaceae (Hoare 1998); a large number of these species belong to non-core *Stigmella*. *Fomoria* has just a few species in the *vannifera* group on Brassicaceae and one in the *weaveri* group on Salicaceae (Hoare 2000b; Doorenweerd et al. 2016a), *Acalyptris* includes ca. seven unnamed species in the *A. platani* group (hosts Phyllanthaceae, Melastomataceae, Loranthaceae) and there is also one *Glaucolepis* in the *raikhonae* group. There is a rich opostegid fauna with 24 named species, but they have not yet been revised. The genera *Opostegoides* and *Pseudopostega* at least occur here, and there is possibly also a species of *Eosopostega* (Hoare 1998).

The fauna of New Guinea is virtually unknown, we have only seen a few species of *Stigmella*.

New Zealand has a fauna quite unrelated to Australia, only the genus *Stigmella*, with the *S. ogygia* group occurs here, with 27 named and ca 12 unnamed species. Interestingly, its sistergroup is the *epicosma* group in South America. Many New Zealand species feed on shrubby and herbaceous Asteraceae, such as *Olearia* or *Senecio*, but there are also species feeding on Malvaceae, Ericaceae and Nothofagaceae (Donner and Wilkinson 1989). There are no opostegids known from New Zealand.

On the many Pacific islands only few Nepticulidae are known, *S. ebbenielseni*, feeding on *Pipturus* (Urticaceae) was described from Guam (van Nieukerken and van den Berg 2003), and a few other Urticaceae feeding *Stigmella* are reported in the same paper from Polynesia and Fiji. No nepticulids are known from Hawaii, but there occurs the interesting endemic genus *Paralopostega*, with a small radiation of species, making leafmines on *Melicope* (formerly *Pelea*, Rutaceae) (Swezey 1921; Zimmerman 1978, Davis, 1989).

On New Caledonia many mines of Nepticulidae have been seen (RJBH), several on Cunoniaceae, but no serious collecting has taken place. The adults of two species are known, but these have not been placed to genus, and require further study. The fauna could well be diverse and important, in common with the very rich and unusual flora of this island.

Nearctic region. The fauna of the Nearctic is relatively poor with 85 named Nepticulidae and ten Opostegidae. The Nepticulidae were revised in the early 1980's (Wilkinson 1979; 1981; Wilkinson and Scoble 1979; Wilkinson and Newton 1981; Newton and Wilkinson 1982) and the Opostegidae recently (Davis and Stonis 2007), but much material remained unstudied and a lot has been collected since the 1980's. Two of us (EvN and CD) have been collecting leafmines throughout North America, whereas Davis has draft descriptions of 20 new species and several other people are contributing to a much better knowledge. Some of our results are shown in this list with notes, and several manuscripts are underway. Even though there are still quite a few unnamed species to describe, overall the fauna is not as rich as in the Palearctic, or even Europe alone. We do not know the cause, but it is interesting to note that other

groups of leafminers are much more diverse in the Nearctic than in the Palearctic (Tischeriidae, Bucculatricidae, several groups of Gracillariidae).

By far the largest genus is *Stigmella*, with groups specialising on amongst others Fagaceae, Betulaceae, Juglandaceae, Rhamnaceae, Rosaceae and Anacardiaceae; typical species groups for this region are the *saginella* and *quercipulchella* groups with oak feeding species and the *prunifoliella* group feeding on Anacardiaceae, Rhamnaceae (*Ceanothus*) and Rosaceae (*Prunus*); most other species groups are shared with the Palearctic. Particularly in California and Arizona there are largely unstudied radiations of species on *Quercus* and *Ceanothus*. *Ectoedemia* is not nearly as diverse as in the Palearctic, and for instance has not more than three species feeding on oaks, but it has some other hosts including Platanaceae and Cornaceae (*Nyssa*). The genus *Acalyptris* still has a large undiscovered diversity, with several species specialising on Cyperaceae in wetlands, like the type species of *Microcalyptris*, *A. scirpi*. *Fomoria*, *Etainia* and *Glaucolepis* are small genera with just a few species, but the barkmining *Zimmermannia* has a few more species (even though we synonymise here eight names) and is a widespread element, with associations with Fagaceae, Salicaceae and possibly Betulaceae. The fauna of southern Florida is more Neotropical with its two species of *Enteucha* on seagrape *Coccoloba uvifera* (Polygonaceae) and various other species in *Acalyptris*, *Stigmella* and *Pseudopostega*. In northern North America there are several Holarctic species, particularly feeding on *Betula* (eg *Ectoedemia occultella*, *minimella*) or Ericaceae (*Fomoria weaveri*) and some European species have been introduced (paper in preparation).

Neotropical Region. Currently 123 named species of Nepticulidae (plus 13 informally named species) and 89 Opostegidae are known, but due to active research new species are added regularly, particularly by Stonis and co-authors (reviewed by van Nieukerken et al. (2016) and see above). The fauna differs remarkably from most other regions, with three endemic genera in Nepticulidae: *Ozadelpha*, *Neotrifurcula* and *Hesperolyra*, and two endemic genera in Opostegidae: *Notiopostega* and *Neopostega*. Most of the named Nepticulidae belong to *Stigmella*, but also here with endemic species groups: the *epicosma* group, the *eurydesma* group, the *barbata* group and the *purpurimaculae* group. Host plant relationships are also special: as in the New Zealand *ogygia* group, many species in the *epicosma* group, particularly common in the Andes, feed on Asteraceae (Stonis et al. 2015a; 2016b). The *Nothofagus* forest of austral South America is the locality for several endemics, and *Notiopostega* is known to make extremely long mines in the cambium of *Nothofagus* trees. It is possible that species of *Neotrifurcula* are also barkminers of *Nothofagus*, and for the *Stigmella purpurimaculae* group there is a strong suspicion that they make leafmines in *Nothofagus* (Stonis et al. 2014). *Ozadelpha* species are associated with Myrtales: Melastomataceae and Myrtaceae, and the only species of *Hesperolyra* where the host is known also feeds on Myrtaceae. There are nine species of *Enteucha*, and where known they feed on Polygonaceae: *Coccoloba*. For the large number of *Acalyptris* species there are only few host records, including Fabaceae and Verbenaceae. In the northern part of the Neotropics there is greater similarity with the Nearctic fauna, and this is particularly the case for the recently discovered diversity of *Quercus* miners in Guatemala and Colombia (Stonis et al. 2013e; Remeikis and Stonis 2015).

The generic placement of the few Neotropical species now placed in *Ectoedemia*, *Fomoria* and *Glaucolepis* requires further study, *Zimmermannia* occurs in Mesoamerica with an unnamed species with genitalia very similar to *Z. bosquella* (Puplesis and Robinson 2000).

In the Opostegidae the genus *Pseudopostega* is remarkably diverse with 82 species, unfortunately as yet without any knowledge of host associations (Davis and Stonis 2007).

Atlantic Islands. Not a single nepticuloid species is known from the Oceanic Atlantic islands south of Macaronesia (such as St. Helena, Ascension, Tristan da Cunha, the Falklands), but their occurrence still could be possible. The fauna of Macaronesia is mostly endemic, particularly on the Canarian Islands, with mostly Mediterranean or African elements of *Glaucolepis*, *Muhabbetana*, *Fomoria*, *Acalyptris* and *Stigmella*. The fauna of the Azores and Madeira is very poor with respectively one and four species, some of which have been introduced (Karsholt and Vieira 2005; Aguiar and Karsholt 2006). No nepticulids are yet known from Cabo Verde, but are expected to occur there, and in the North Atlantic there are no Nepticulidae known from Greenland or Iceland, nor any of the smaller islands.

Catalogue

SUPERFAMILY **NEPTICULOIDEA** Stainton, 1854: 295
FAMILY **NEPTICULIDAE** Stainton, 1854: 295 (TG: *Nepticula* Heyden, 1843)
Family Stigmellidae Hampson, 1918: 387 (TG: *Stigmella* Schrank, 1802)
Subfamily Pectinivalvinae Scoble, 1983: 12 (TG: *Pectinivalva* Scoble, 1983) (syn.: Puplesis, 1994: 36)
Subfamily Nepticulinae Stainton, 1854
Subfamily Stigmellinae Hampson, 1918
Subfamily Trifurculinae Scoble, 1983: 16 (TG: *Trifurcula* Zeller, 1848), **syn. n.**
Tribe Nepticulini Stainton, 1854: 295
Tribe Stigmellini Hampson, 1918
Tribe Trifurculini Scoble, 1983: 16 **syn. n.**

Enteucha Meyrick, 1915a: 241 (TS/M: *Enteucha cyanochlora* Meyrick, 1915) [1]
Johanssonia Borkowski, 1972a: 702; JH of *Johanssonia* Selensky, 1914 (Annelida, Hirudinea) (TS/OD,M: *Nepticula acetosae* Stainton, 1854) (syn: van Nieukerken, 1986a: 7)
Artaversala Davis, 1978: 219 (TS/OD,M: *Artaversala gilvafascia* Davis, 1978) (syn: van Nieukerken, 1986a: 7)
Oligoneura Davis, 1978: 217; JH of *Oligoneura* Bigot, 1878 (Brachiopoda) (TS/OD,M: *Oligoneura basidactyla* Davis, 1978) (syn: van Nieukerken, 1986a: 7)
Manoneura Davis, 1979: 276; RN for *Oligoneura* Davis, 1978 (syn: van Nieukerken, 1986a: 7) [1]

Johanssoniella Koçak, 1981: 99; RN for *Johanssonia* Borkowski (syn: van Nieukerken, 1986a: 7)

Enteucha acetosae (Stainton, 1854) van Nieukerken, 1986a: 7 (Fig. 25) WP
‡ *Nepticula acetosae* Shield, 1853: 4153 NN
Nepticula acetosae Stainton, 1854: 303
Nepticula acetosella Doubleday, 1859: 36 UE
Nepticula arifoliella Klimesch, 1940b: 92
Stigmella acetosae (Stainton, 1854) Beirne, 1945: 200
Johanssonia acetosae (Stainton, 1854) Borkowski, 1972a: 702
Johanssoniella acetosae (Stainton, 1854) Koçak, 1981: 99
Stigmella arifoliella (Klimesch, 1940) Hering, 1957: 912
‡ *Nepticula arifoliella* var. *altvateri* Skala, 1941b: 79

Enteucha diplocosma (Meyrick, 1921) Diškus & Puplesis, 2003: 321 OR
Nepticula diplocosma Meyrick, 1921a: 411

Enteucha acuta Puplesis & Diškus in Puplesis et al., 2002: 21 NEO

Enteucha basidactyla (Davis, 1978) van Nieukerken, 1986a: 54 [1] NEA,NEO
Oligoneura basidactyla Davis, 1978: 218
Manoneura basidactyla (Davis, 1978) Davis, 1979: 276

Enteucha contracolorea Puplesis & Robinson, 2000: 20 NEO

Enteucha cyanochlora Meyrick, 1915a: 241 NEO

Enteucha gilvafascia (Davis, 1978) van Nieukerken, 1986a: 54 NEA,NEO
Artaversala gilvafascia Davis, 1978: 221

Enteucha hilli Puplesis & Robinson, 2000: 19 NEO

Enteucha snaddoni Puplesis & Robinson, 2000: 21 NEO

Enteucha trinaria (Puplesis & Robinson, 2000), **comb. n.** NEO
Manoneura trinaria Puplesis & Robinson, 2000: 23

Enteucha terricula Puplesis & Robinson, 2000: 20 NEO

Varius Scoble, 1983: 14 (TS/OD,M: *Stigmella ochnicola* Vári, 1955)

Varius ochnicola (Vári, 1955) Scoble, 1983: 14 AFR
Stigmella ochnicola Vári, 1955: 336
Varius ochnicolus (Vári, 1955) Scoble, 1983: 14 [variant]

Simplimorpha Scoble, 1983: 15 (TS/OD,M: *Stigmella lanceifoliella* Vári, 1955)

Simplimorpha promissa (Staudinger, 1871) van Nieukerken, 1986a: 6 WP
Nepticula promissa Staudinger, 1871: 325
Nepticula robiniella Gustafsson, 1973: 197 (syn: van Nieukerken, 1986a: 6)
Stigmella promissa (Staudinger, 1871) Klimesch, 1951b: 64

Simplimorpha lanceifoliella (Vári, 1955) Scoble, 1983: 15 AFR
Stigmella lanceifoliella Vári, 1955: 331

Stigmella Schrank, 1802: 169 (TS/SD (Walsingham, 1908a: 1007)[2]: *Phalaena* (*Tinea*) *anomalella* Goeze, 1783)
Nepticula Heyden, 1843: 208 (TS/SD (Walsingham, 1908a: 1007): *Tinea aurella* Fabricius, 1755) (syn: Walsingham, 1908a: 1008) [2]
Dysnepticula Börner, 1925: 370 (TS/OD: *Phalaena* (*Tinea*) *anomalella* Goeze, 1783)
Astigmella Puplesis, 1984a: 111 (TS/OD: *Astigmella dissona* Puplesis, 1984) (syn: van Nieukerken, 1986a: 7)
[*Microsetia* Stephens, 1834 *sensu* Kirby, 1897: 313 (TS/incorrect SD by Kirby, 1897: *Nepticula microtheriella* Stainton, 1854) (see Wilkinson 1978: 17)]

Non-core *Stigmella* [3]

ungrouped species in Non-core *Stigmella*

Stigmella freyella (Heyden, 1858) Vári, 1950: 182 WP
Nepticula freyella Heyden, 1858: 175
Stigmella kurilensis Puplesis, 1987: 8 EP
Stigmella ebbenielseni van Nieukerken & Van den Berg, 2003: 28 AUS
Stigmella resplendensella (Chambers, 1875) Newton & Wilkinson, 1982: 456 [4] NEA
Nepticula resplendensella Chambers, 1875b: 118
Stigmella unifasciella (Chambers, 1875) Newton & Wilkinson, 1982: 438 NEA
Nepticula unifasciella Chambers, 1875b: 119
Stigmella gallicola van Nieukerken & Nishida in van Nieukerken et al., 2016: 7 NEO

***Stigmella prunifoliella* group** (Newton & Wilkinson, 1982: 385) [5]
Stigmella prunetorum group (Johansson, 1971: 245)
Stigmella bifasciella group (Wilkinson & Scoble, 1979: 59)
Stigmella prunetorum (Stainton, 1855) Beirne, 1945: 198 WP,EP
Nepticula prunetorum Stainton, 1855: 72
Nepticula dimidiatella Herrich-Schäffer, 1855a: 352 (syn: Herrich-Schäffer, 1860: 59)
Nepticula perpusillella Herrich-Schäffer, 1855a: 353 (syn: Frey, 1856: 390)
Nepticula prunetella Doubleday, 1859: 36 UE
Nepticula ligustrella Rössler, 1867: 395 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula punctella Threlfall, 1884: 113 ISS
‡ *Nepticula prunetorum* var. *aviella* Skala, 1934b: 6 NNLM
‡ *Stigmella prunetorum* var. *aviella* (Skala, 1934b) NNLM
Stigmella diniensis (Klimesch, 1975) Leraut, 1980: 49 WP
Nepticula diniensis Klimesch, 1975c: 5
Stigmella ceanothi (Braun, 1910) Newton & Wilkinson, 1982: 387 NEA
Nepticula ceanothi Braun, 1910: 172
Stigmella cerea (Braun, 1917) Newton & Wilkinson, 1982: 407 NEA
Nepticula cerea Braun, 1917: 172
Stigmella intermedia (Braun, 1917) Wilkinson & Scoble, 1979: 62 NEA
Nepticula intermedia Braun, 1917: 171

Stigmella prunifoliella (Clemens, 1861) Newton & Wilkinson, 1982: 385 NEA
Nepticula prunifoliella Clemens, 1861: 84
Nepticula bifasciella Clemens, 1862: 133 (syn: Newton & Wilkinson, 1982: 385)
Nepticula serotinaeella Chambers, 1873: 126 (syn: Braun, 1917: 170)
Stigmella bifasciella (Clemens, 1862) Wilkinson & Scoble, 1979: 59
Stigmella rhoifoliella (Braun, 1912) Newton & Wilkinson, 1982: 391 NEA
Nepticula rhoifoliella Braun, 1912: 93
Stigmella gossypii (Forbes & Leonard, 1930) Newton & Wilkinson, 1982: 404 NEO
Nepticula gossypii Forbes & Leonard, 1930: 149
Stigmella schinivora van Nieukerken, 2016 in van Nieukerken et al., 2016: 15 NEO

Stigmella ultima **group** (Puplesis, 1984a: 116)
Stigmella aceris (Frey, 1857) Gerasimov, 1952: 222 WP
Nepticula aceris Frey, 1857: 386
Nepticula penicillata Heinemann & Wocke, [1876]: 744 [6] (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula szöcsi Klimesch, 1956: 423 (syn: Klimesch, 1978a: 246)
Nepticula szoecsi Klimesch, 1956: 423 ISS (syn: Klimesch, 1978a: 246)
Stigmella acerna Puplesis, 1988: 277 [7] WP
‡ *Stigmella acerifoliella* Dovnar-Zapolski, 1969: 20 NNLM; **syn. n.** [7]
Stigmella bicolor Puplesis, 1988: 276 EP
Stigmella bumbegerensis Puplesis, 1984d: 509 EP
Stigmella kozlovi Puplesis, 1984a: 118 EP
Stigmella monella Puplesis, 1984a: 117 EP
Stigmella orientalis Kemperman & Wilkinson, 1985: 21 EP
Stigmella semiaurea Puplesis, 1988: 275 WP,EP
Stigmella tegmentosella Puplesis, 1984a: 117 EP
Stigmella ultima Puplesis, 1984a: 117 EP
Stigmella japonica Kemperman & Wilkinson, 1985: 20 (syn: Puplesis, 1994: 81)

Stigmella ulmivora **group** (Johansson, 1971: 244)
Stigmella kazakhstanica Puplesis in Puplesis et al. 1991: 70 WP,EP
Stigmella pimschoorli Puplesis, 1994: 64 (syn: Puplesis et al., 1996: 192)
Stigmella ulmiphaga (Preissecker, 1942) Klimesch, 1948b: 62 WP
Nepticula ulmiphaga Preissecker, 1942: 208
Nepticula gracilivora Skala, 1942: 6 (syn: Klimesch, 1975c: 4)
Stigmella ulmivora (Fologne, 1860) Beirne, 1945: 199 [8] WP
Nepticula ulmivora Fologne, 1860: 92
Nepticula ulmifoliae Hering, 1931: 531 (syn: Emmet, 1974d: 151)
Nepticula ulmicola Hering, 1932: 569 (syn: Emmet, 1974d: 151)
Stigmella ulmifoliae (Hering, 1931) Vári, 1944b: xxv
Stigmella ulmicola (Hering, 1932) Vári, 1944b: xxv
‡ *Nepticula ulmella* Hofmann, 1858: 191 NN (syn: Segerer, 1997: 188) [8]

Stigmella viscerella (Stainton, 1853) Beirne, 1945: 199 WP
Nepticula viscerella Stainton, 1853: 3958
Nepticula subvirescens Meyrick, 1934b: 523 (syn: Diškus & Puplesis, 2003: 343)
Nepticula tauromeniella Groschke, 1944: 117 (syn: Klimesch, 1975c: 2)
Stigmella tauromeniella (Groschke, 1944) Hering, 1957: 1090
Stigmella alisa Puplesis, 1985c: 63 EP
Stigmella amuriella Puplesis, 1985c: 62 EP
Stigmella auricularia Puplesis, Diškus & Juchnevič in Puplesis & Diškus, 2003a: 245 EP
Stigmella multispicata Rocienė & Stonis in Stonis & Rocienė, 2014: 205 [9] EP
Stigmella nireae Kemperman & Wilkinson, 1985: 18 EP
Stigmella palionisi Puplesis, 1984b: 596[7] EP
Stigmella nakamurai Kemperman & Wilkinson, 1985: 16 **syn. n.** [10]

***Stigmella eurydesma* group** (Puplesis & Robinson, 2000: 32)
Stigmella albilamina Puplesis & Robinson, 2000: 33 NEO
Stigmella eurydesma (Meyrick, 1915a) Davis, 1984: 18 NEO
Nepticula eurydesma Meyrick, 1915a: 255
Stigmella fuscilamina Puplesis & Robinson, 2000: 34 NEO

***Stigmella saginella* group** (Wilkinson & Scoble, 1979: 39)
Stigmella castaneaefoliella (Chambers, 1875b) Wilkinson & Scoble, 1979: 44 NEA
Nepticula castaneaefoliella Chambers, 1875b: 117
Stigmella flavipedella (Braun, 1914) Wilkinson & Newton, 1981: 31 NEA
Nepticula flavipedella Braun, 1914: 19
Stigmella macrocarpae (Freeman, 1967), **comb. n.** [11] NEA
Nepticula latifasciella Chambers, 1878: 106 JPH of *Nepticula latifasciella* Herrich-Schäffer, 1855 [11] (syn: Wilkinson & Scoble, 1979: 47)
Nepticula macrocarpae Freeman, 1967: 19
Stigmella latifasciella (Chambers, 1878) Wilkinson & Scoble, 1979: 47
Stigmella nigriverticella (Chambers, 1875b) Newton & Wilkinson, 1982: 423 NEA
Nepticula nigriverticella Chambers, 1875b: 118
Nepticula maculosella Chambers, 1880a: 193 (syn: Braun, 1917: 194)
Stigmella saginella (Clemens, 1861) Wilkinson & Scoble, 1979: 39 NEA
Nepticula saginella Clemens, 1861: 85
Nepticula fuscocapitella Chambers, 1873: 128 (syn: Braun, 1917: 195)
Nepticula quercicastanella Chambers, 1873: 127 (syn: Braun, 1917: 195)
Stigmella sclerostylota Newton & Wilkinson, 1982: 429 NEA
Stigmella aurifasciata Diškus & Stonis in Stonis et al., 2013c: 8 NEO
Stigmella crassifoliae Remeikis & Stonis, 2015: 410 NEO
Stigmella jaguari Remeikis & Stonis in Stonis et al., 2013c: 6 NEO
Stigmella lauta Diškus & Stonis in Stonis et al., 2013c: 6 NEO
Stigmella robleae Remeikis & Stonis, 2015: 411 NEO
Stigmella sublauta Remeikis & Stonis in Stonis et al., 2013c: 8 NEO

***Stigmella paliurella* group** (van Nieukerken, 1986a: 8)
Stigmella birgittae Gustafsson, 1985: 171 [12] WP,AFR
Stigmella omani Puplesis & Diškus, 2003a: 207 (syn: van Nieukerken, 2010: 493)
‡ *Nepticula amseli* Skala, 1941b: 78 NNLM; **syn. n.** [12]
Stigmella abaiella Klimesch, 1979: 21 [13] WP
Stigmella ficulnea Puplesis & Krasnilnikova, in Puplesis, 1994: 65 [13] WP
Stigmella longicornuta Puplesis & Diškus, 2003a: 217 WP
Stigmella paliurella Gerasimov, 1937: 285 [14] WP
Nepticula paliurella (Gerasimov, 1937) Klimesch, 1940a: 177 [14]
Nepticula paliurella Klimesch, 1940c
Stigmella turbatrix Puplesis, 1994: 66 [15] WP,EP
‡ *Stigmella celtivora* Dovnar-Zapolski, 1969: 39 NNLM
Stigmella zizyphi Walsingham, 1911: 190 WP,AFR
Stigmella ziziphivora Gustafsson, 1985: 171 (syn: van Nieukerken, 2010: 495)
Nepticula zizyphi (Walsingham, 1911) Skala, 1938a: 45
Stigmella morivora Hirano, 2010: 128 EP
Stigmella sruogai Puplesis & Diškus, 2003a: 204 EP
Stigmella isochalca (Meyrick, 1916b) Diškus & Puplesis, 2003: 325 OR
Nepticula isochalca Meyrick, 1916b: 6
Stigmella nepali Puplesis & Diškus, 2003a: 206 OR
Stigmella phyllanthina (Meyrick, 1906b) Common, 1990: 156 AUS
Nepticula phyllanthina Meyrick, 1906b: 60

***Stigmella naturnella* group** (new)
Stigmella dissona group (Puplesis, 1994: 58)
Stigmella naturnella (Klimesch, 1936) Klimesch, 1948b: 65 WP,EP
Nepticula naturnella Klimesch, 1936: 205
Astigmella dissona Puplesis, 1984a: 112 (syn: van Nieukerken et al., 2004a: 133)
Stigmella dissona (Puplesis, 1984a) Puplesis, 1994: 58
Stigmella mirabella (Puplesis, 1984a) Puplesis, 1994: 58 EP
Astigmella mirabella Puplesis, 1984a: 112

***Stigmella tiliae* group** (Johansson, 1971: 245)
Stigmella tiliae (Frey, 1856) Beirne, 1945: 198 WP
Nepticula tiliae Frey, 1856: 375
Stigmella sashai Puplesis, 1984b: 594 EP
Stigmella regina Puplesis, 1984b: 596 (syn: Rocienė & Stonis, 2013: 95)

***Stigmella betulicola* group** (Johansson, 1971: 245)
Stigmella corylifoliella group (Wilkinson & Scoble, 1979: 50)
Stigmella alnetella (Stainton, 1856) Beirne, 1945: 198 WP
Nepticula alnetella Stainton, 1856: 43

Stigmella betulicola (Stainton, 1856) Beirne, 1945: 198 WP,EP,NEA
Nepticula betulicola Stainton, 1856: 42
Nepticula betulicolella Doubleday, 1859: 36 UE
Nepticula betulicola var. *nanivora* Petersen, 1930: 61 (syn: Johansson, 1971: 245)
Stigmella nanivora (Petersen, 1930) Hering, 1957: 183
Stigmella glutinosae (Stainton, 1858) Beirne, 1945: 198 WP
Nepticula glutinosae Stainton, 1858: 96
Nepticula distinguenda Heinemann, 1862b: 305 (syn: Schoorl & Wilkinson, 1986: 234)
Nepticula rubescens Heinemann, 1871: 214 (syn: Klimesch, 1950a: 27)
Nepticula glutinosella Porritt, 1883: 173 UE
Stigmella rubescens (Heinemann, 1871) Gerasimov, 1952: 256
‡ *Nepticula glutinosae* var. *alni-viridis* Skala, 1939e: 111 NNLM (syn: Klimesch, 1950a: 27)
‡ *Nepticula rubescens* var. *incanae* Skala, 1941a: 57 NNLM (syn: Klimesch, 1950a: 27)
Stigmella gutlebiella A. Laštuvka & Huemer, 2002: 604 WP
Stigmella luteella (Stainton, 1857a) Beirne, 1945: 198 WP,EP
Nepticula luteella Stainton, 1857a: 110
Nepticula luteellina Skala, 1941b: 79 (syn: Skala, 1948: 121)
Stigmella microtheriella (Stainton, 1854) Fletcher & Clutterbuck, 1945: 59 [16] WP,EP,[NEA,AUS]
Nepticula microtheriella Stainton, 1854: 302
Stigmella cathepostis Kemperman & Wilkinson, 1985: 10 **syn. n.** [16]
Microsetia microtheriella (Stainton, 1854) Kirby, 1897: 313
Stigmella nivenburgensis (Preissecker, 1942) Klimesch, 1951b: 59 [17] WP,EP
Nepticula nivenburgensis Preissecker, 1942: 209
Stigmella populnea Kemperman & Wilkinson, 1985: 13 **syn. n.** [17]
Stigmella sakhalinella Puplesis, 1984a: 115 WP,EP
Stigmella discidia Schoorl & Wilkinson, 1986: 237 (syn: van Nieukerken & Johansson, 1987: 470)
Nepticula distinguenda auct. [misapplied] (syn: van Nieukerken & Johansson, 1987: 470)
Stigmella distinguenda auct. [misapplied]
Stigmella attenuata Puplesis, 1985c: 62 EP
Stigmella betulifoliae Puplesis & Diškus, 2003a: 179 EP
Stigmella conchyliata Kemperman & Wilkinson, 1985: 11 EP
Stigmella cornuta Rocienė & Stonis in Stonis et al., 2013e: 206 [18] EP
Stigmella excelsa Puplesis & Diškus, 2003a: 182 EP
Stigmella kumashidei Hirano, 2014: 20 EP
Stigmella oplismeniella Kemperman & Wilkinson, 1985: 12 EP
Stigmella pamirbetulae Puplesis & Diškus, 2003a: 180 EP

Stigmella titivillitia Kemperman & Wilkinson, 1985: 14 EP
Stigmella caryaefoliella (Clemens, 1861), **stat. rev., comb. n.** [19] NEA
Nepticula caryaefoliella Clemens, 1861: 84
Stigmella corylifoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 50 NEA
Nepticula corylifoliella Clemens, 1861: 83
Nepticula virginiella Clemens, 1861: 83 (syn: Braun, 1917: 179)
Nepticula minimella Chambers, 1873: 179 (syn: Braun, 1917: 179)
Nepticula opulifoliella Braun, 1914: 22 (syn: Wilkinson & Scoble, 1979: 50)
Nepticula paludicola Braun, 1917: 177 (syn: Wilkinson & Scoble, 1979: 51)
Nepticula exasperata Braun, 1930: 17 (syn: Wilkinson & Scoble, 1979: 51)
Stigmella juglandifoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 57 NEA
Nepticula juglandifoliella Clemens, 1861: 84
Stigmella longisacca Newton & Wilkinson, 1982: 436 NEA
Stigmella myricafoliella (Busck, 1900) Grossbeck, 1917: 145 [20] NEA
Nepticula myricafoliella Busck, 1900: 238
Nepticula obscurella Braun, 1912: 95 **syn. n.** [20]
Stigmella ostryaefoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 54 [21] NEA
Nepticula ostryaefoliella Clemens, 1861: 83
Stigmella himalayai Puplesis & Diškus, 2003a: 208 OR
Stigmella xystodes (Meyrick, 1916b) Diškus & Puplesis, 2003: 328 WP,OR
Nepticula xystodes Meyrick, 1916b: 6
Nepticula liochalca Meyrick, 1916b: 6 (syn: van Nieukerken, 2010: 496)
Nepticula homophaea Meyrick, 1918b: 181 (syn: van Nieukerken, 2010: 496)
Stigmella liochalca (Meyrick, 1916b) Diškus & Puplesis, 2003: 328
Stigmella homophaea (Meyrick, 1918b) Diškus & Puplesis, 2003: 363
Stigmella allophylica Scoble, 1978b: 97 AFR
Stigmella allophylivora Gustafsson, 1985: 167 AFR
Stigmella androflavus Scoble, 1978b: 104 AFR
Stigmella generalis Scoble, 1978b: 102 AFR
Stigmella geranica Scoble, 1978b: 96 AFR
Stigmella hortorum Scoble, 1978b: 99 AFR
Stigmella pelanodes (Meyrick, 1920b), **comb. n.** [22] AFR
Nepticula pelanodes Meyrick, 1920b: 116
Stigmella potgieteri Scoble, 1978b: 99 AFR
Stigmella satarensis Scoble, 1978b: 97 AFR
Stigmella tragilis Scoble, 1978b: 98 AFR
Stigmella tropicatella Legrand, 1965: 27 [23] AFR
Stigmella triumfettica Scoble, 1978b: 107 AFR

***Stigmella divina* group** (Puplesis & Diškus, 2003a: 212)
Stigmella divina Puplesis, Diškus & van Nieukerken, 1997: 55 WP
Stigmella maculifera Puplesis & Diškus, 2003a: 212 WP
Stigmella skulei Puplesis & Diškus, 2003a: 213 WP

Unplaced tropical species - most probably in non-core *Stigmella*

African species

Stigmella fluida group (Scoble, 1978b: 92)

Stigmella ingens group (Scoble, 1978b: 111)

Stigmella abachausi (Janse, 1948) Scoble, 1978b: 104 AFR

Nepticula abachausi Janse, 1948: 162

Stigmella abutilonica Scoble, 1978b: 93 AFR

Stigmella ampullata Scoble, 1978b: 108 AFR

Stigmella angustivalva Scoble, 1978b: 113 AFR

Stigmella caliginosa (Meyrick, 1921b) Scoble, 1983: 43 AFR

Nepticula caliginosa Meyrick, 1921b: 140

Stigmella celtifoliella Vári, 1955: 338 AFR

Stigmella charistis Vári, 1963: 71 AFR

Stigmella confinalis Scoble, 1978b: 111 AFR

Stigmella crotonica Scoble, 1978b: 100 AFR

Stigmella dombeyivora Scoble, 1978b: 107 AFR

Stigmella ficivora Gustafsson, 1985: 170 AFR

Stigmella fluida (Meyrick, 1911a) Scoble, 1978b: 94 AFR

Nepticula fluida Meyrick, 1911a: 236

Stigmella galactacma (Meyrick, 1924b) Diškus & Puplesis, 2003: 364 AFR

Nepticula galactacma Meyrick, 1924b: 89

Stigmella grewiae Scoble, 1978b: 112 AFR

Stigmella gustafssoni (Căpuşe, 1975) Diškus & Puplesis, 2003: 366 AFR

Nepticula gustafssoni Căpuşe, 1975: 211

Stigmella ingens (Meyrick, 1913) Scoble, 1978b: 112 AFR

Nepticula ingens Meyrick, 1913: 327

Stigmella irrorata (Janse, 1948) Scoble, 1978b: 95 AFR

Nepticula irrorata Janse, 1948: 168

Stigmella letabensis Scoble, 1978b: 113 AFR

Stigmella liota Vári, 1963: 73 AFR

Stigmella maytenivora Gustafsson, 1985: 174 AFR

Stigmella naibabi Mey, 2004: 29 AFR

Stigmella nigrata (Meyrick, 1913) Scoble, 1978b: 106 AFR

Nepticula nigrata Meyrick, 1913: 326

Stigmella panconista (Meyrick, 1920a) Diškus & Puplesis, 2003: 363 AFR

Nepticula panconista Meyrick, 1920a: 312

Stigmella parinarella Vári, 1955: 337 AFR

Stigmella perplexa (Janse, 1948) Scoble, 1978b: 103 AFR

Nepticula perplexa Janse, 1948: 172

Stigmella platyzona Vári, 1963: 67 AFR

Stigmella porphyreuta (Meyrick, 1917a) Scoble, 1978b: 110 AFR

Nepticula porphyreuta Meyrick, 1917a: 13

Stigmella pretoriata Scoble, 1978b: 109 AFR
Stigmella protosema (Meyrick, 1921b) Scoble, 1978b: 109 AFR
Nepticula protosema Meyrick, 1921b: 140
Stigmella rhomboivora Gustafsson, 1985: 167 AFR
Stigmella rhynchosiella Vári, 1955: 338 AFR
Stigmella urbica (Meyrick, 1913) Scoble, 1978b: 103 AFR
Nepticula urbica Meyrick, 1913: 326
Stigmella uwusebi Mey, 2004: 30 AFR
Stigmella varii Scoble, 1978b: 95 AFR
Stigmella wollofella (Gustafsson, 1972) Gustafsson, 1985: 174 [24] AFR
Nepticula wollofella Gustafsson, 1972: 158
Nepticula mandingella Gustafsson, 1972: 157 **syn. n.** [24]
Stigmella mandingella (Gustafsson, 1972) Diškus & Puplesis, 2003: 366
Stigmella worcesteri Scoble, 1983: 16 RN for *S. pallida* Scoble, 1978 AFR
Stigmella pallida Scoble, 1978b: 105 JSH of *Stigmella pallida* (Braun, 1917)

Australian species
Stigmella leucargyra (Meyrick, 1906b) Nielsen, 1996: 16 AUS
Nepticula leucargyra Meyrick, 1906b: 57
Stigmella symmora (Meyrick, 1906b) Nielsen, 1996: 16 AUS
Nepticula symmora Meyrick, 1906b: 59

Oriental species
Stigmella aeriventris (Meyrick, 1932) Diškus & Puplesis, 2003: 364 OR
Nepticula aeriventris Meyrick, 1932: 312
Stigmella alicia (Meyrick, 1928) Diškus & Puplesis, 2003: 364 OR
Nepticula alicia Meyrick, 1928b: 461
Stigmella argyrodoxa (Meyrick, 1918) Fletcher, 1933: 83 OR
Nepticula argyrodoxa Meyrick, 1918b: 181
Stigmella auxozona (Meyrick, 1934) Diškus & Puplesis, 2003: 365 OR
Nepticula auxozona Meyrick, 1934a: 468
Stigmella elachistarcha (Meyrick, 1934) Diškus & Puplesis, 2003: 365 OR
Nepticula elachistarcha Meyrick, 1934a: 467
Stigmella hoplometalla (Meyrick, 1934) Puplesis & Diškus, 2003a: 215 OR
Nepticula hoplometalla Meyrick, 1934a: 467
Stigmella ipomoeella (Gustafsson, 1976) Diškus & Puplesis, 2003: 328 OR
Nepticula ipomoeella Gustafsson, 1976: 45
Stigmella neodora (Meyrick, 1918) Diškus & Puplesis, 2003: 363 OR
Nepticula neodora Meyrick, 1918b: 182
Stigmella oligosperma (Meyrick, 1934) Diškus & Puplesis, 2003: 365 OR
Nepticula oligosperma Meyrick, 1934a: 468
Stigmella polydoxa (Meyrick, 1911) Diškus & Puplesis, 2003: 363 OR
Nepticula polydoxa Meyrick, 1911c: 107

Core *Stigmella* [3]

Stigmella rhamnella/lapponica/sanguisorbae cluster

***Stigmella tiliella* group** (Puplesis et al., 2002: 63)

Stigmella tiliella (Braun, 1912) Newton & Wilkinson, 1982: 442 NEA
Nepticula tiliella Braun, 1912: 90

Stigmella kimae Puplesis & Robinson, 2000: 35 NEO

***Stigmella rhamnella* group** (new)

Stigmella alaternella (Le Marchand, 1937) Klimesch, 1948b: 63 WP
Nepticula alaternella Le Marchand, 1937: 234

Stigmella armeniana Puplesis, 1994: 90 WP

Stigmella catharticella (Stainton, 1853) Beirne, 1945: 199 WP
Nepticula catharticella Stainton, 1853: 3955

Stigmella crenulatae (Klimesch, 1975) van Nieukerken, 1986a: 8 WP
Nepticula crenulatae Klimesch, 1975c: 2

Stigmella kopetdagica Puplesis, 1994: 92 WP

Stigmella pyrellicola (Klimesch, 1978) van Nieukerken, 1986a: 9 WP
Nepticula pyrellicola Klimesch, 1978b: 264

Stigmella rhamnella (Herrich-Schäffer, 1860) Klimesch, 1951b: 56 WP
Nepticula rhamnella Herrich-Schäffer, 1860: 60
Nepticula rhamnella var. *rhamnipumilae* Klimesch, 1950a: 49

Stigmella rhamnophila (Amsel, 1934) van Nieukerken, 1986a: 8 WP
Nepticula rhamnella rhamnophila Amsel, 1934: 317
‡ *Nepticula rhamnophila* Amsel & Hering, 1931: 142 NNLM

Stigmella klimeschi Puplesis, 1988: 274 EP

Stigmella kurotsubarai Kemperman & Wilkinson, 1985: 15 EP

Stigmella taigae Puplesis, 1984a: 112 EP

Stigmella condaliafoliella (Busck, 1900) Grossbeck, 1917: 145 NEA
Nepticula condaliafoliella Busck, 1900: 238

Stigmella diffasciae (Braun, 1910) Newton & Wilkinson, 1982: 398 NEA
Nepticula diffasciae Braun, 1910: 172

Stigmella inconspicuella Newton & Wilkinson, 1982: 400 NEA

Stigmella rhamnicola (Braun, 1916) Newton & Wilkinson, 1982: 393 NEA
Nepticula rhamnella Braun, 1912: 96 JPH of *Nepticula rhamnella* Herrich-Schäffer, 1860
Nepticula rhamnicola Braun, 1916: 55 RN for *N. rhamnella* Braun, 1912

Stigmella maya Remeikis & Stonis in Stonis et al., 2013b: 224 NEO

***Stigmella sanguisorbae* group** (van Nieukerken, 1986a: 10)
Stigmella rosaefoliella group (Wilkinson & Scoble, 1979: 14)

Stigmella muricatella (Klimesch, 1978) van Nieukerken, 1986a: 9 WP,EP
Nepticula muricatella Klimesch, 1978b: 266

Stigmella polymorpha Puplesis & Diškus, 2003a: 183 WP
Stigmella rolandi van Nieukerken, 1990c: 239 WP,EP
Stigmella sanguisorbae (Wocke, 1865) Gerasimov, 1952: 258 WP
Nepticula sanguisorbae Wocke, 1865: 269
Stigmella thuringiaca (Petry, 1904) Gerasimov, 1952: 263 WP,EP
Nepticula thuringiaca Petry, 1904: 267
Nepticula nickerli Rebel in Nickerl, 1908: 116 (syn: Rebel, 1909: (269))
Stigmella fasciola Puplesis & Diškus, 2003a: 185 EP
Stigmella trisyllaba Puplesis in Puplesis et al., 1992: 51 EP
Stigmella rosaefoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 14 [25] NEA
Nepticula rosaefoliella Clemens, 1861: 85
Stigmella rosaefoliella rosaefoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 14[25]

Stigmella lapponica **group** (Johansson, 1971: 245)
Stigmella malella group (Johansson, 1971: 245)
Stigmella confusella (Wood & Walsingham, 1894) Vári, 1944a: 215 WP,EP,NEA
Nepticula confusella Wood & Walsingham, 1894: 272
Stigmella lapponica (Wocke, 1862) Fletcher & Clutterbuck, 1945: 61 WP,EP,NEA
Nepticula lapponica Wocke, 1862: 251
Nepticula lapponicella Herrich-Schäffer, 1863c: 23 UE
Nepticula lusatica Schütze, 1905: 204 (syn: Johansson & Nielsen, 1990: 141)
Nepticula vossensis Grønlien, 1928: 217 (syn: Krogerus, 1971: 31)
Stigmella lusatica (Schütze, 1905) Beirne, 1945: 200
Stigmella vossensis (Grønlien, 1928) Gerasimov, 1952: 269
Stigmella malella (Stainton, 1854) Beirne, 1945: 199 WP
Nepticula malella Stainton, 1854: 304
Nepticula angustella Heinemann & Wocke, [1876]: 756 [6] (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula nigrobrunnella Groschke, 1939: 716 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula nigrobrunella auct. ISS
Stigmella nigrobrunnella (Groschke, 1939) Hering, 1957: 837
‡ *Nepticula malella* var. *prunicola* Skala, 1939f: 126 NN
Stigmella maloidica Puplesis in Puplesis & Arutyunova, 1991: 573 EP
Stigmella braunella (Jones, 1933) Wilkinson & Scoble, 1979: 13 NEA
Nepticula braunella Jones, 1933: 49
Stigmella slingerlandella (Kearfott, 1908) Wilkinson & Scoble, 1979: 19 NEA
Nepticula slingerlandella Kearfott, 1908: 187

unplaced in *rhamnella/lapponica/sanguisorbae* cluster
Stigmella boehmeriae Kemperman & Wilkinson, 1985: 54 EP
Stigmella costaricensis van Nieukerken & Nishida in van Nieukerken et al., 2016: 19 NEO
Stigmella intronia van Nieukerken & Nishida in van Nieukerken et al., 2016: 21 NEO

***Stigmella salicis* cluster**

***Stigmella ogygia* group** (new) [26]

Stigmella aigialeia Donner & Wilkinson, 1989: 17 AUS
Stigmella aliena Donner & Wilkinson, 1989: 17 AUS
Stigmella atrata Donner & Wilkinson, 1989: 18 AUS
Stigmella cassiniae Donner & Wilkinson, 1989: 18 AUS
Stigmella childi Donner & Wilkinson, 1989: 19 AUS
Stigmella cypracma (Meyrick, 1916) Dugdale, 1988: 53 AUS
Nepticula cypracma Meyrick, 1916c: 419
Nepticula perissopa Meyrick, 1919: 354 (syn: Donner & Wilkinson, 1989: 20)
Stigmella perissopa (Meyrick, 1919) Dugdale, 1988: 54
Stigmella erysibodea Donner & Wilkinson, 1989: 21 AUS
Stigmella fulva (Watt, 1921) Dugdale, 1988: 53 AUS
Nepticula fulva Watt, 1921: 215
Stigmella hakekeae Donner & Wilkinson, 1989: 22 AUS
Stigmella hamishella Donner & Wilkinson, 1989: 23 AUS
Stigmella hoheriae Donner & Wilkinson, 1989: 24 AUS
Stigmella ilsea Donner & Wilkinson, 1989: 25 AUS
Stigmella insignis (Philpott, 1927) Dugdale, 1988: 53 AUS
Nepticula insignis Philpott, 1927: 89
Stigmella kaimanua Donner & Wilkinson, 1989: 26 AUS
Stigmella laquaeorum (Dugdale, 1971) Dugdale, 1988: 53 AUS
Nepticula laquaeorum Dugdale, 1971: 117
Stigmella lucida (Philpott, 1919) Dugdale, 1988: 54 AUS
Nepticula lucida Philpott, 1919: 225
Stigmella maoriella (Walker, 1864) Dugdale, 1988: 54 AUS
Tinea maoriella Walker, 1864: 1008
Stigmella ogygia (Meyrick, 1889) Dugdale, 1988: 54 AUS
Nepticula ogygia Meyrick, 1889: 187
Nepticula erechtitus Watt, 1924: 686 (syn: Donner & Wilkinson, 1989: 20)
Stigmella erechtitus (Watt, 1924) Dugdale, 1988: 53
Stigmella oriastra (Meyrick, 1917) Dugdale, 1988: 54 AUS
Nepticula oriastra Meyrick, 1917b: 247
Stigmella palaga Donner & Wilkinson, 1989: 31 AUS
Stigmella platina Donner & Wilkinson, 1989: 32 AUS
Stigmella progama (Meyrick, 1924) Dugdale, 1988: 54 AUS
Nepticula progama Meyrick, 1924a: 662
Stigmella progonopis (Meyrick, 1921) Dugdale, 1988: 54 AUS
Nepticula progonopis Meyrick, 1921c: 336
Stigmella propalaea (Meyrick, 1889) Dugdale, 1988: 54 AUS
Nepticula propalaea Meyrick, 1889: 187
Stigmella sophorae (Hudson, 1939) Dugdale, 1988: 54 AUS
Nepticula sophorae Hudson, 1939: 469

Stigmella tricentra (Meyrick, 1889) Dugdale, 1988: 54 AUS
Nepticula tricentra Meyrick, 1889: 187
Stigmella watti Donner & Wilkinson, 1989: 35 AUS

***Stigmella epicosma* group** (new) [27]
Stigmella andina (Meyrick, 1915) Davis, 1984: 18 NEO
Nepticula andina Meyrick, 1915a: 255
Stigmella baccharicola Diškus & Stonis in Stonis et al., 2016b: 119 NEO
Stigmella bipartita Diškus & Stonis in Stonis et al., 2016b: 107 NEO
Stigmella confertae Diškus & Stonis in Stonis et al., 2016b: 124 NEO
Stigmella costalimai (Bourquin, 1961) Davis, 1984: 18 [28] NEO
Nepticula costalimai Bourquin, 1961: 31
Stigmella cuprata (Meyrick, 1915) Davis, 1984: 18 NEO
Nepticula cuprata Meyrick, 1915a: 255
Stigmella emarginatae Diškus & Stonis in Stonis et al., 2016b: 104 NEO
Stigmella epicosma (Meyrick, 1915) Davis, 1984: 18 NEO
Nepticula epicosma Meyrick, 1915a: 255
Stigmella guittonae (Bourquin, 1961) Davis, 1984: 18 [28] NEO
Nepticula guittonae Bourquin, 1961: 32
Stigmella hamata Puplesis & Robinson, 2000: 30 NEO
Stigmella imperatoria Puplesis & Robinson, 2000: 30 NEO
Stigmella johannis (Zeller, 1877) Davis, 1984: 18 NEO
Nepticula johannis Zeller, 1877: 454
Stigmella latifoliae Remeikis, Diškus & Stonis in Stonis et al., 2016b: 115 NEO
Stigmella marmorea Puplesis & Robinson, 2000: 26 NEO
Stigmella mevia Remeikis & Stonis in Stonis & Remeikis, 2016: 311 NEO
Stigmella montanotropica Puplesis & Diškus in Puplesis et al., 2002: 23 NEO
Stigmella nubimontana Puplesis & Diškus in Puplesis et al., 2002: 24 NEO
Stigmella olyritis (Meyrick, 1915) Davis, 1984: 18 NEO
Nepticula olyritis Meyrick, 1915a: 256
Stigmella pangorica Diškus & Stonis in Stonis et al., 2015a: 580 NEO
Stigmella peruanica Puplesis & Robinson, 2000: 27 NEO
Stigmella podanthae Diškus & Stonis in Stonis et al., 2016a: 120 NEO
Stigmella racemifera Šimkevičiūtė & Stonis in Šimkevičiūtė et al., 2009: 270 NEO
Stigmella rubeta Puplesis & Diškus in Puplesis et al., 2002: 24 NEO
Stigmella rudis Puplesis & Robinson, 2000: 26 NEO
Stigmella schoorli Puplesis & Robinson, 2000: 29 NEO
Stigmella serpentina Diškus & Stonis in Stonis et al., 2015a: 576 NEO
Stigmella sinuosa Remeikis & Stonis in Stonis & Remeikis, 2016: 310 NEO
Stigmella tripartita Diškus & Stonis in Stonis et al., 2016b: 110 NEO

***Stigmella salicis* group** (Johansson, 1971: 244)
Stigmella fuscotibiella group (Newton & Wilkinson, 1982: 385)

Stigmella aiderensis Puplesis, 1988: 277 WP
Stigmella arbusculae (Klimesch, 1951) Hering, 1957: 930 WP
Nepticula arbusculae Klimesch, 1951c: 149
Stigmella assimilella (Zeller, 1848) Fletcher & Clutterbuck, 1945: 61 WP,EP
Nepticula assimilella Zeller, 1848: 327
Nepticula tremulaefoliella Sorhagen, 1922: 48 (syn: Johansson & Nielsen, 1990: 200)
Stigmella tremulaefoliella (Sorhagen, 1922) Gerasimov, 1952: 265
‡ *Lyonetia nigricornella* Mann in Zeller, 1848: 327 NN
Stigmella benanderella (Wolff, 1955) Hering, 1957: 930 WP
Nepticula benanderella Wolff, 1955b: 49
‡ *Nepticula scandicella* Jonasson in Krogerus et al., 1971: 30 NN
Stigmella flavescens Puplesis, 1994: 131 WP, EP
Stigmella myrtillella (Stainton, 1857b) Vári, 1944a: 215 WP
Nepticula myrtillella Stainton, 1857b: 44
‡ *Nepticula myrtillella* var. *uliginosi* Skala, 1941b: 80 NNLM
Stigmella obliquella (Heinemann, 1862) Vári, 1944a: 215 WP,EP
Nepticula obliquella Heinemann, 1862b: 316
Nepticula wockeella Heinemann, 1871: 223 (syn: Johansson & Nielsen, 1990: 198)
Nepticula diversa Glitz, 1872: 24 (syn: Glitz, 1887: 277)
Stigmella babylonicae Hartig, 1949: 94 (syn: van Nieukerken, 1986a: 11)
Stigmella wockeella (Heinemann, 1871) Gerasimov, 1952: 269
Stigmella pallidiciliella Klimesch, 1948a: 165 [29] WP
Nepticula purpureae Skala, 1948: 121 (syn: van Nieukerken, 1986a: 11)
Nepticula pallidiciliella (Klimesch, 1948a) Wolff, 1955a: 86
Stigmella salicis (Stainton, 1854) Fletcher & Clutterbuck, 1945: 60 [30] WP,EP,NEA
Nepticula salicis Stainton, 1854: 302
Nepticula salicella Herrich-Schäffer, 1855a: 354 UE
Nepticula salicivorella Doubleday, 1859: 36 UE
Nepticula uniformis Heinemann, 1871: 210 (syn: van Nieukerken, 1986a: 11)
Nepticula dewitziella Sorhagen, 1885: 285 (syn: Johansson & Nielsen, 1990: 192)
Nepticula auritella Skala, 1939f: 128 (syn: van Nieukerken, 1986a: 11)
Stigmella libiezi Dufrane, 1949: 8 (syn: van Nieukerken, 1986a: 11)
Stigmella uniformis (Heinemann, 1871) Gerasimov, 1952: 267
Stigmella auritella (Skala, 1939) Hering, 1957: 929
‡ *Nepticula salicis* ab. *crombruggheella* Dufrane, 1930: 30
‡ *Nepticula salicis* ab. *februella* Crombrugghe, 1907: 14
‡ *Nepticula salicis* ab. *interrupta* Skala, 1933a: 32
Stigmella trimaculella (Haworth, 1828) Fletcher & Clutterbuck, 1945: 61 WP,EP
Tinea trimaculella Haworth, 1828: 583
Lyonetia rufella Zeller, 1839: 215 (syn: Herrich-Schäffer, 1855a: 358)

Nepticula populella Herrich-Schäffer, 1855a: 357 (syn: Frey, 1856: 381)
Nepticula albicornella Kollar in Nowicki, 1860: 231 (syn: Rebel, 1901: 228)
Nepticula gilvella Rössler, 1867: 395 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula populicola Sorhagen, 1922: 88 (syn: Skala, 1948: 121)
Stigmella subtrimaculella Dufrane, 1949: 10 (syn: Borkowski, 1969: 107)
Microsetia trimaculella (Haworth, 1828) Stephens, 1834: 269
Nepticula trimaculella (Haworth, 1828) Stainton, 1849: 29
Nepticula rufella (Zeller, 1839) Zeller, 1848: 328
Stigmella populicola (Sorhagen, 1922) Gerasimov, 1952: 252
‡ *Nepticula trimaculella* ab. *semipictella* Steudel in Steudel & Hoffmann, 1882: 244

Stigmella vimineticola (Frey, 1856) Fletcher & Clutterbuck, 1945: 60 WP
Nepticula vimineticola Frey, 1856: 382

Stigmella zelleriella (Snellen, 1875) van Nieukerken, 1983a: 60 WP
Nepticula zelleriella Snellen, 1875: 116
Nepticula repentiella Wolff, 1955a: 82 (syn: van Nieukerken, 1983a: 60)
Nepticula lappovimella Svensson, 1976: 204 (syn: van Nieukerken, 1983a: 60)
Stigmella repentiella (Wolff, 1955) Hering, 1957: 929
Stigmella lappovimella (Svensson, 1976) Svensson, 1985: 78

Stigmella azusa Hirano, 2010: 129 EP

Stigmella johanssoni Puplesis & Diškus, 1996c: 181 EP

Stigmella juratae Puplesis, 1988: 279 EP

Stigmella kondarai Puplesis, 1988: 277 EP

Stigmella sexcornuta Rocienė & Stonis in Stonis & Rocienė, 2014: 205 EP

Stigmella tenryuensis Hirano, 2014: 26 EP

Stigmella tranocrossa Kemperman & Wilkinson, 1985: 27 EP
Stigmella ussurica Puplesis, 1987: 8 (syn: Puplesis, 1994: 128)

Stigmella vittata Kemperman & Wilkinson, 1985: 28 EP

Stigmella fibigeri Puplesis & Diškus, 2003a: 209 OR

Stigmella aromella Wilkinson & Scoble, 1979: 27 NEA

Stigmella fuscotibiella (Clemens, 1862) Wilkinson & Scoble, 1979: 23 NEA
Nepticula fuscotibiella Clemens, 1862: 133
Nepticula ciliaefuscella Chambers, 1873: 128 (syn: Chambers, 1875a: 117)
Nepticula discolorella Braun, 1912: 86 (syn: Braun, 1917: 185)

Stigmella pallida (Braun, 1912) Newton & Wilkinson, 1982: 418 NEA
Nepticula pallida Braun, 1912: 85

Stigmella populetorum (Frey & Boll, 1878) Wilkinson & Scoble, 1979: 26 NEA
Nepticula populetorum Frey & Boll, 1878: 276

Stigmella molinensis van Nieukerken & Snyers in van Nieukerken et al., 2016: 22 NEO

***Stigmella quercipulchella/anomalella/oxyacanthella* cluster**

***Stigmella quercipulchella* group** (Wilkinson & Scoble, 1979: 65)

Stigmella altella (Braun, 1914) Wilkinson & Newton, 1981: 58 NEA
Nepticula altella Braun, 1914: 21

Stigmella quercipulchella (Chambers, 1882) Wilkinson & Scoble, 1979: 65 NEA
Nepticula quercipulchella Chambers, 1882: 105
Nepticula terminella Braun, 1914: 23 (syn: Wilkinson & Scoble, 1979: 65)

Stigmella variella (Braun, 1910) Wilkinson & Scoble, 1979: 67 NEA
Nepticula variella Braun, 1910: 173

Stigmella guatemalensis Diškus & Stonis in Stonis et al., 2013c: 10 NEO

***Stigmella anomalella* group** (Johansson, 1971: 245)

Stigmella anomalella (Goeze, 1783) Walsingham, 1908a: 1008 [2] WP,EP,[NEA]
Phalaena anomalella Goeze, 1783: 168
Phalaena grisearosae Retzius, 1783: 55
Tinea penicilla Thunberg, 1794: 88 (syn: Karsholt & Nielsen, 1986: 452)
Tinea rosella Schrank, 1802: 139 (syn: Stainton, 1854: 297)
Nepticula aeneella Heinemann, 1862a: 254 (syn: Schoorl et al., 1985: 98)
Nepticula fletcheri Tutt, 1899: 211 (syn: Krogerus, 1971: 30)
Nepticula laticuniculella Sauber, 1904: 55 (syn: Hering, 1957: 902)
Stigmella rubicurrens Walsingham, 1908a: 1009 [2] (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula rosarum Sorhagen, 1922: 30 (syn: van Nieukerken & Johansson, 1987: 470)
Nepticula zermattensis Weber, 1936: 668 (syn: van Nieukerken, 1986a: 9)
Nepticula helbigi Hartig, 1941: 160 (syn: van Nieukerken & Johansson, 1987: 468)
Stigmella caulescentella Klimesch, 1948a: 162 [29] (syn: van Nieukerken, 1986a: 9)
Stigmella anomalella pacifica Puplesis, 1987: 10
Nepticula anomalella (Goeze, 1783) Stainton, 1854: 297
Dysnepticula anomalella (Goeze, 1783) Börner, 1925: 370
Nepticula rosella (Schrank, 1802) Sand, 1879: 200
Stigmella rosella (Schrank, 1802) Walsingham, 1908a: 1008[1]
Stigmella fletcheri (Tutt, 1899) Fletcher & Clutterbuck, 1945: 59
Nepticula rubicurrens (Walsingham, 1908a) Rebel, 1910: 373
Stigmella rosarum (Sorhagen, 1922) Gerasimov, 1952: 256
Stigmella zermattensis (Weber, 1936) Gerasimov, 1952: 270
Stigmella anomalella fletcheri (Tutt, 1899) Hering, 1957: 902
Stigmella anomalella helbigi (Hartig, 1941) Hering, 1957: 902

Stigmella centifoliella (Zeller, 1848) Beirne, 1945: 199 [31] WP,[NEA]
‡ *Nepticula centifoliella* Heyden, 1843: 208 NN
Nepticula centifoliella Zeller, 1848: 315
Nepticula hodgkinsoni Stainton, 1884: 103 (syn: Bradley et al., 1972: 2)

Stigmella rosaefoliella pectocatena Wilkinson & Scoble, 1979: 18 **syn. n.**[31]
Stigmella hodgkinsoni (Stainton, 1884) Gerasimov, 1952: 243
Stigmella spinosissimae (Waters, 1928) Beirne, 1945: 198 WP,EP
Nepticula spinosissimae Waters, 1928: 105

***Stigmella hybnerella* group** (Johansson, 1971: 245)
Stigmella nitidella group (Johansson, 1971: 245)
Stigmella paradoxa group (Emmet, 1976: 238)
Stigmella irregularis group (Puplesis, 1994: 61)
Stigmella hybnerella (Hübner, 1796) Fletcher & Clutterbuck, 1945: 59 WP
Tinea hybnerella Hübner, 1796: pl. 34: 236
Caloptilia ampelipennella Hübner, [1825]: 427 URN
Tinea posticella Haworth, 1828: 584 (syn: Stainton, 1849: 29)
Oecophora gratiosella Duponchel, 1843: 323 (syn: Fletcher & Clutterbuck, 1945: 59)
Nepticula ignobilella Stainton, 1849: 29 (syn: Emmet, 1974c: 77)
Nepticula latifasciella Herrich-Schäffer, 1855a: 352 (syn: Herrich-Schäffer, 1855a: 352)
Microsetia posticella (Haworth, 1828) Stephens, 1834: 269
Lyonetia hübnerella (Hübner, 1796) Zeller, 1839: 215
Nepticula gratiosella (Duponchel, 1843) Stainton, 1849: 29
Lithocolletis gratiosella (Duponchel, 1843) Bruand, [1851]: 86
Stigmella gratiosella (Duponchel, 1843) Beirne, 1945: 200
Stigmella ignobilella (Stainton, 1849) Fletcher & Clutterbuck, 1945: 60
Nepticula ignobiliella auct. ISS
Stigmella mespilicola (Frey, 1856) Klimesch, 1948b: 57 (Fig. 26) WP
Nepticula mespilicola Frey, 1856: 392
Nepticula ariella Herrich-Schäffer, 1860: 60 (syn: Frey, 1880: 422)
Stigmella ariella (Herrich-Schäffer, 1860) Klimesch, 1948b: 57
Stigmella irregularis Puplesis, 1994: 61 WP
Stigmella inopinata A. Laštuvka & Z. Laštuvka, 1990: 197 WP
Stigmella paradoxa (Frey, 1858) Emmet, 1970: 3 WP
Nepticula paradoxa Frey, 1858a: 14
Nepticula nitidella Heinemann, 1862a: 257 (syn: Heinemann & Wocke, [1876]: 734)[6]
Stigmella juryi Puplesis, 1991: 125 (syn: van Nieukerken et al., 2004a: 140)
Stigmella nitidella (Heinemann, 1862) Gerasimov, 1952: 249
Stigmella pyrivora Gustafsson, 1981: 457 WP
Stigmella malifoliella Puplesis in Puplesis & Arutyunova, 1991: 571 EP
Stigmella montana Puplesis, 1991: 126 EP
Stigmella taeniola (Braun, 1925) Newton & Wilkinson, 1982: 382 NEA
Nepticula taeniola Braun, 1925b: 226
Stigmella stigmaciella Wilkinson & Scoble, 1979: 38 NEA

***Stigmella incognitella* group** (new)

Stigmella pomella group (Johansson, 1971: 244)

Stigmella azaroli (Klimesch, 1978) van Nieukerken, 1986a: 13 [32] WP

Nepticula azaroli Klimesch, 1978b: 261

Stigmella fuscacalyptriella Puplesis, 1994: 149 WP

Stigmella incognitella (Herrich-Schäffer, 1855) van Nieukerken, 1986a: 13 WP

Nepticula incognitella Herrich-Schäffer, 1855a: 349

Nepticula pomella Vaughan, 1858: 43 (syn: van Nieukerken, 1986a: 13)

Nepticula mali Hering, 1932: 568 (syn: van Nieukerken, 1986a: 13)

Stigmella mali (Hering, 1932) Gerasimov, 1952: 247

Stigmella pomella (Vaughan, 1858) Fletcher & Clutterbuck, 1945: 58

Stigmella perpygmaeella (Doubleday, 1859) Karsholt & Nielsen, 1976: 18 [32] WP

Tinea pygmaeella Haworth, 1828: 586; JPH of *Tinea pygmaeella* [Denis & Schiffermüller], 1775), now *Argyresthia pygmaeella* (Argyresthiidae)

Nepticula perpygmaeella Doubleday, 1859: 36; RN for *Tinea pygmaeella* Haworth, 1828

Microsetia pygmaeella (Haworth, 1828) Stephens, 1834: 269

Nepticula pygmaeella (Haworth, 1828) Stainton, 1853: 3958

Stigmella pygmaeella (Haworth, 1828) Klimesch, 1951b: 55

Stigmella elegantiae Puplesis & Diškus, 2003a: 210 OR

***Stigmella oxyacanthella* group** (Johansson, 1971: 245)

Stigmella crataegifoliella group (Wilkinson & Scoble, 1979: 29)

Stigmella caspica Puplesis, 1994: 109 WP

Stigmella crataegella (Klimesch, 1936) Klimesch, 1948b: 62 WP

Nepticula crataegella Klimesch, 1936: 200

Stigmella indigena Puplesis, 1994: 111 (syn: Puplesis et al., 1996: 192)

Nepticula gratiosella sensu Tutt, 1899: 253

Stigmella desperatella (Frey, 1856) Beirne, 1945: 200 WP

Nepticula desperatella Frey, 1856: 374

Nepticula pyricola Wocke, 1877: 49 (syn: Schoorl et al., 1985: 94)

Stigmella pyricola (Wocke, 1877) Klimesch, 1951b: 57

Stigmella hahniella (Wörz, 1937) Gerasimov, 1952: 241 WP

Nepticula hahniella Wörz, 1937: 290

Stigmella lanceolata Puplesis, 1994: 111 WP

Stigmella magdalenae (Klimesch, 1950) Emmet, 1979: 25 [33] WP

Nepticula nylandriella var. *magdalenae* Klimesch, 1950b: 72

Nepticula nylandriella auct. [misapplied, before 1972] (syn: Borkowski, 1975: 523)

Stigmella nylandriella magdalenae (Klimesch, 1950b) Klimesch, 1961: 752

Nepticula magdalenae Klimesch, 1950b (Borkowski, 1975: 523)

Stigmella minusculella (Herrich-Schäffer, 1855) Beirne, 1945: 198 WP,[NEA]

Nepticula minusculella Herrich-Schäffer, 1855a: 348

Nepticula chalybeia Braun, 1914: 20 (syn: Schoorl et al., 1985: 92)

Nepticula embonella Klimesch, 1978b: 259 (syn: Schoorl et al., 1985: 92)
Stigmella chalybeia (Braun, 1914) Wilkinson & Scoble, 1979: 35
Stigmella nylandriella (Tengström, 1848) Beirne, 1945: 200 [34] WP
Lyonetia nylandriella Tengström, 1848: 152
Nepticula aucupariae Frey, 1857: 376 (syn: Borkowski, 1975: 522)
Nepticula aucupariella Porritt, 1883: 172 UE
Nepticula nylandriella (Tengström, 1848) Herrich-Schäffer, 1855: 359 [34]
Stigmella aucupariae (Frey, 1857) Beirne, 1945: 199
Stigmella oxyacanthella (Stainton, 1854) Beirne, 1945: 200 [35] WP,[NEA]
Nepticula oxyacanthella Stainton, 1854: 298
Nepticula oxyacanthaecolella Doubleday, 1859: 298 URN
Micropteryx pomivorella Packard, 1870: 237 **syn. n.** [35]
Nepticula cotoneastri Sorhagen, 1922: 42 (syn: Schoorl et al., 1985: 87)
Nepticula aeneella auct. [misapplied, before 1985]
Stigmella aeneella auct. [misapplied, before 1985]
Nepticula pomivorella (Packard, 1870) Busck, 1901: 52
Stigmella pomivorella (Packard, 1870) Wilkinson & Scoble, 1979: 33
Stigmella cotoneastri (Sorhagen, 1922) Klimesch, 1948b: 60
‡ *Nepticula chaenomelis* Skala, 1936: 79 NNLM (syn: van Nieukerken, 1986a: 10)
‡ *Nepticula oxyacanthella* var. *mespili* Skala, 1940: 144 NNLM
‡ *Nepticula oxyacanthella* var. *oxymalella* Skala, 1933b: 130 NNLM
‡ *Nepticula oxyacanthella* var. *oxysorbi* Skala, 1933b: 130 NNLM
‡ *Stigmella chaenomelis* (Skala, 1936) Hering, 1957: 275
Stigmella pyri (Glitz, 1865) Vári, 1944a: 214 WP
Nepticula pyri Glitz, 1865: 42
Stigmella regiella (Herrich-Schäffer, 1855) Vári, 1944a: 214 WP
Nepticula regiella Herrich-Schäffer, 1855a: 351
Nepticula corvimontana Hering, 1935: 6 (syn.: Borkowski, 1969: 103)
Stigmella corvimontana (Hering, 1935) Gerasimov, 1952: 234
Stigmella stettinensis (Heinemann, 1871) Gerasimov, 1952: 262 WP
Nepticula stettinensis Heinemann, 1871: 210
Stigmella torminalis (Wood, 1890) Beirne, 1945: 200 WP
Nepticula torminalis Wood, 1890: 209
Stigmella alaurulenta Kemperman & Wilkinson, 1985: 23 EP
Stigmella aurora Puplesis, 1984a: 119 WP,EP
Stigmella chaenomelae Kemperman & Wilkinson, 1985: 23 EP
Stigmella crataegi Gerasimov, 1937: 283 EP
Stigmella hissariella Puplesis, 1994: 112 EP
Stigmella honshui Kemperman & Wilkinson, 1985: 24 EP
Stigmella micromelis Puplesis, 1985c: 59 [36] EP
Stigmella crataegivora Puplesis, 1985c: 60 **syn. n.** [36]
Stigmella nostrata Puplesis, 1984a: 113 EP
Stigmella sorbivora Kemperman & Wilkinson, 1985: 25 EP

Stigmella zumii Kemperman & Wilkinson, 1985: 26 EP
Stigmella amelanchierella (Clemens, 1861) Davis & Wilkinson, 1983: 3 [37] NEA
Nepticula amelanchierella Clemens, 1861: 84
Stigmella crataegifoliella (Clemens, 1861) Wilkinson & Scoble, 1979: 30 NEA
Nepticula crataegifoliella Clemens, 1861: 83
Stigmella heteromelis Newton & Wilkinson, 1982: 405 NEA
Stigmella purpuratella (Braun, 1917) Newton & Wilkinson, 1982: 381 [38] NEA
Nepticula purpuratella Braun, 1917: 176
Stigmella scinanella Wilkinson & Scoble, 1979: 36 **syn. n.** [38]
Stigmella scintillans (Braun, 1917) Wilkinson & Scoble, 1979: 33 NEA
Nepticula scintillans Braun, 1917: 167

***Stigmella argentifasciella* group** (new)
Stigmella argentifasciella (Braun, 1912) Newton & Wilkinson, 1982: 453 NEA
Nepticula argentifasciella Braun, 1912: 100

***Stigmella aurella/ruficapitella* cluster** [39]

***Stigmella styracicolella* group** (new)
Stigmella styracicolella (Klimesch, 1978) van Nieukerken, 1986a: 14 WP
Nepticula styracicolella Klimesch, 1978b: 267
Stigmella egonokii Kemperman & Wilkinson, 1985: 53 EP

***Stigmella speciosa* group** (new) [39]
Stigmella kuznetzovi Puplesis, 1994: 152 WP
Stigmella speciosa (Frey, 1858) Walsingham, 1916: 159 WP
Nepticula speciosa Frey, 1858b: 27
Nepticula pseudoplatanella Weber, 1936: 671 (syn: Borkowski, 1969: 98)
Stigmella pseudoplatanella (Weber, 1936) Gerasimov, 1952: 254
‡ *Nepticula aceris* var. *pseudoplatanella* Skala, 1933b: 132 NNLM
‡ *Nepticula speciosa* var. *monspessulani* Skala, 1939d: 144 NNLM
Stigmella lonicerarum (Frey, 1857) Gerasimov, 1952: 246 WP
Nepticula lonicerarum Frey, 1857: 383
Nepticula lonicerarum var. *lentinensis* Skala, 1935: 79 (syn: van Nieukerken, 1986a: 13)
Nepticula lonicerarum var. *livonica* Skala, 1935: 79 (syn: van Nieukerken, 1986a: 13)
Nepticula lonicerarum var. *teutonica* Skala, 1935: 79 (syn: van Nieukerken, 1986a: 13)
Stigmella monticulella Puplesis, 1984a: 114 [40] EP
Stigmella gracilipae Hirano, 2014: 22 **syn. n.** [40]

***Stigmella aurella* group** (Johansson, 1971: 243) [39]
Stigmella lediella group (Puplesis, 1984b:583)

Stigmella aeneofasciella (Herrich-Schäffer, 1855) Gerasimov, 1952: 222 WP
Nepticula aeneofasciella Herrich-Schäffer, 1855a: 353
Nepticula aeneofasciata Frey, 1856: 376 UE
Stigmella aurella (Fabricius, 1775) Walsingham, 1908a: 1009 [2] WP
Tinea aurella Fabricius, 1775: 666
Nepticula fragariella Heinemann, 1862a: 263 (syn: Borkowksi, 1975: 503)
Nepticula nitens Fologne, 1862: 164 (syn: Klimesch, 1981: 114)
Nepticula gei Wocke, 1871: 336 (syn: Borkowksi, 1975: 503)
Nepticula albicomella Heinemann & Wocke, [1876]: 748 [6] (syn: Johansson & Nielsen, 1990: 207)
Nepticula fruticosella Müller-Rutz in Vorbrodt & Müller-Rutz, 1914: 591 (syn: van Nieukerken, 1986a: 12)
Microsetia aurella (Fabricius, 1775) Stephens, 1834: 268
Nepticula aurella (Fabricius, 1775) Heyden, 1843: 209
Stigmella fragariella (Heinemann, 1862a) Vári, 1944a: 214
Stigmella nitens (Fologne, 1862) Vári, 1944a: 214
Stigmella gei (Wocke, 1871) Gerasimov, 1952: 240
Stigmella fruticosella (Müller-Rutz in Vorbrodt & Müller-Rutz, 1914) Gerasimov, 1952: 240
‡ *Nepticula gei* ab. *semicolorella* Eppelsheim, 1891: 351 NNLM
‡ *Nepticula gei* var. *geirubi* Skala, 1940: 143 NNLM
Stigmella auromarginella (Richardson, 1890) Gerasimov, 1952: 229 WP
Nepticula auromarginella Richardson, 1890: 30
Stigmella dryadella (Hofmann, 1868) Klimesch, 1951b: 58 WP
Nepticula dryadella Hofmann, 1868: 29
Stigmella filipendulae (Wocke, 1871) Gerasimov, 1952: 238 [41] WP,EP
Nepticula filipendulae Wocke, 1871: 338
Nepticula ulmariae Wocke, 1879: 79 (syn: van Nieukerken et al., 2012a: 250)
Stigmella palmatae Puplesis, 1984a: 115 **syn. n.** [41]
Stigmella ulmariae (Wocke, 1879) Gerasimov, 1952: 266
Stigmella geimontani (Klimesch, 1940) Klimesch, 1961: 754 WP
Nepticula geimontani Klimesch, 1940b: 89
Stigmella lediella (Schleich, 1867) Gerasimov, 1952: 245 [42] WP,EP
Nepticula lediella Schleich, 1867: 449
Stigmella magica Puplesis, 1985c: 63 (syn: Puplesis, 1994: 146)
Stigmella rhododendri Puplesis, 1985c: 65 (syn: Puplesis, 1994: 146)
Stigmella sesplicata Kemperman & Wilkinson, 1985: 36 **syn. n.** [42]
‡ *Nepticula lediella* ab. *auromarginata* Petersen, 1930
‡ *Stigmella rhododendrifolia* Dovnar-Zapolski & Tomilova, 1978: 29 NNLM; **syn. n.** [42]
Stigmella poterii (Stainton, 1857) Fletcher & Clutterbuck, 1945: 59 WP
Nepticula poterii Stainton, 1857c: 116
Nepticula poteriella Doubleday, 1859: 36 UE

Nepticula comari Wocke, 1862: 253 (syn: Borkowksi, 1975: 506)
Nepticula geminella Frey, 1870: 288 (syn: Karsholt & Nielsen, 1976: 17)
Nepticula palustrella Frey, 1870: 287 (syn: van Nieukerken, 1986a: 12)
Nepticula occultella Heinemann, 1871: 215 (syn: Borkowksi, 1975: 506)
Nepticula tengströmi Nolcken, 1871: 776 (syn: Borkowksi, 1975: 506)
Nepticula potentillae Glitz, 1872: 24 (syn with *occultella*: Heinemann &Wocke, [1876]: 749)
Nepticula diffinis Wocke, 1874: 100 (syn: Borkowksi, 1975: 506)
Nepticula serella Stainton, 1888a: 260 (syn: Borkowksi, 1975: 506)
Nepticula elisabethella Szőcs, 1957: 321 (syn: van Nieukerken, 1986a: 12)
Stigmella comari (Wocke, 1862) Gerasimov, 1952: 234
Stigmella geminella (Frey, 1870) Gerasimov, 1952: 240
Stigmella occultella (Heinemann, 1871) Gerasimov, 1952: 250
Stigmella tengstroemi (Nolcken, 1871) Gerasimov, 1952: 263
Stigmella diffinis (Wocke, 1874) Gerasimov, 1952: 236
Stigmella serella (Stainton, 1888a) Gerasimov, 1952: 259

Stigmella pretiosa (Heinemann, 1862) Gerasimov, 1952: 253 WP
Nepticula pretiosa Heinemann, 1862a: 261
Nepticula bollii Frey, 1873: 144 (syn: van Nieukerken, 1986a: 12)
Stigmella geimontani tatrensis Borkowski, 1970b: 546 (syn: Borkowksi, 1975: 507)
Stigmella bollii (Frey, 1873) Gerasimov, 1952: 231

Stigmella splendidissimella (Herrich-Schäffer, 1855) Klimesch, 1951b: 58 WP
Nepticula splendidissimella Herrich-Schäffer, 1855a: 353
Nepticula splendidissima Frey, 1856: 393 UE
Nepticula dulcella Heinemann, 1862a: 267 (syn: Karsholt & Nielsen, 1976: 17)
Nepticula inaequalis Heinemann, 1862b: 302 (syn: Johansson & Nielsen, 1990: 209)
Nepticula fragarivora Carolsfeld-Krause, 1944: 158 (syn: Karsholt & Nielsen, 1976: 17)
Stigmella dulcella (Heinemann, 1862a) Gerasimov, 1952: 237
Stigmella inaequalis (Heinemann, 1862b) Gerasimov, 1952: 244
Stigmella fragarivora (Carolsfeld-Krause, 1944) Hering, 1957: 455
‡ *Nepticula peterseniella* Skala, 1941b: 78 NN (syn: van Nieukerken, 1986a: 12)
‡ *Stigmella fragarivora peterseniella* (Skala, 1941b) Hering, 1957: 455

Stigmella stelviana (Weber, 1938) Klimesch, 1948b: 67 WP
‡ *Nepticula stelviana* Wocke, 1881: 205 NN
Nepticula stelviana Weber, 1938: 5
Nepticula crantziella Weber, 1945: 401 (syn: Klimesch, 1950a: 27)
Stigmella crantziella (Weber, 1945) Klimesch, 1948b: 69

Stigmella tormentillella (Herrich-Schäffer, 1860) Gerasimov, 1952: 264 WP
Nepticula tormentillella Herrich-Schäffer, 1860: 60

Stigmella acrochaetia Kemperman & Wilkinson, 1985: 31 EP

Stigmella alikurokoi Kemperman & Wilkinson, 1985: 31 EP

Stigmella ichigoiella Kemperman & Wilkinson, 1985: 35 EP
Stigmella longispina Puplesis, 1994: 166 [43] WP,EP
Stigmella spiculifera Kemperman & Wilkinson, 1985: 37 [44] EP
Stigmella oa Kemperman & Wilkinson, 1985: 52 **syn. n.** [44]
Stigmella villosella (Clemens, 1861) Newton & Wilkinson, 1982: 410 NEA
Nepticula villosella Clemens, 1861: 84
Nepticula dallasiana Frey & Boll, 1876: 288 (syn: Braun, 1917: 174)

***Stigmella sorbi* group** (Johansson, 1971: 244) [36]
Stigmella amygdali group (van Nieukerken, 1986a: 13)
Stigmella amygdali (Klimesch, 1978) van Nieukerken, 1986a: 13 WP
Nepticula amygdali Klimesch, 1978b: 264
Stigmella cerasi Puplesis & Diškus, 1996b: 178 WP
Stigmella plagicolella (Stainton, 1854) Fletcher & Clutterbuck, 1945: 60 WP
Nepticula plagicolella Stainton, 1854: 303
‡ *Nepticula plagicolella* var. *avianella* Skala, 1934c: 30 NNLM
‡ *Stigmella plagicolella avianella* (Skala, 1934c) Hering, 1957: 839 NNLM
Stigmella sorbi (Stainton, 1861) Fletcher & Clutterbuck, 1945: 60 WP,EP
Nepticula sorbi Stainton, 1861: 91
Nepticula sorbiella Porritt, 1883: 171 UE
Nepticula sorbi var. *cotoneastrella* Weber, 1936: 670
Stigmella cotoneastrella (Weber, 1936) Hering, 1957: 338
‡ *Nepticula plagicolella* var. *malicola* Skala, 1939d: 95 NNLM
‡ *Stigmella plagicolella malicola* (Skala, 1939) Hering, 1957: 664 NNLM
Stigmella aflatuniae Puplesis & Diškus, 1996b: 180 EP
Stigmella azukinashii Hirano, 2014: 25 EP
Stigmella hamamelella Hirano, 2014: 20 EP
Stigmella lurida Puplesis, 1994: 132 [45] EP
Stigmella motiekaitisi Puplesis, 1994: 135 WP,EP
Stigmella pourthiaeella Hirano, 2014: 24 EP
Stigmella subsorbi Puplesis, 1994: 134 EP
Stigmella tenebrica Puplesis & Diškus, 2003a: 214 OR

***Stigmella lemniscella* group** (new) [39]
Stigmella marginicolella group (Johansson, 1971: 243)
Stigmella continuella (Stainton, 1856) Fletcher & Clutterbuck, 1945: 59 WP,EP
Nepticula continuella Stainton, 1856: 42
Stigmella uigurica Puplesis, 1985c: 62 (syn: Puplesis, 1994: 137)
Stigmella lemniscella (Zeller, 1839) van Nieukerken, 1986a: 11 WP
Lyonetia lemniscella Zeller, 1839: 215
Nepticula marginicolella Stainton, 1853: 3958 (syn: van Nieukerken, 1986a: 12)
Nepticula suberosella Toll, 1934b: 76 (syn: Hering, 1957: 1089)
Nepticula fulvomacula Skala, 1936: 79 (syn: Borkowski, 1969: 114)

Nepticula lemniscella (Zeller, 1839) Zeller, 1848: 313
Stigmella marginicolella (Stainton, 1853) Fletcher & Clutterbuck, 1945: 59
Stigmella fulvomacula (Skala, 1936) Gerasimov, 1952: 239

Stigmella zagulaevi Puplesis, 1994: 139 WP

Stigmella gimmonella (Matsumura, 1931) Kuroko, 1982a: 448 EP
Nepticula gimmonella Matsumura, 1931: 1114

Stigmella talassica Puplesis in Puplesis et al., 1992: 54 EP

Stigmella apicialbella (Chambers, 1873) Newton & Wilkinson, 1982: 413 NEA
Nepticula apicialbella Chambers, 1873: 127
Nepticula leucostigma Braun, 1912: 88 (syn: Braun, 1914: 21)

***Stigmella floslactella* group** (Johansson, 1971: 244) [39]

Stigmella carpinella (Heinemann, 1862) Gerasimov, 1952: 232 WP
Nepticula carpinella Heinemann, 1862a: 251

Stigmella floslactella (Haworth, 1828) Fletcher & Clutterbuck, 1945: 61 WP
Tinea floslactella Haworth, 1828: 585
Nepticula saxatilella Grönlien, 1932: 114 (syn: van Nieukerken & Johansson, 1987: 461)
Microsetia floslactella (Haworth, 1828) Stephens, 1834: 268
Nepticula floslactella (Haworth, 1828) Stainton, 1849: 29
Stigmella saxatilella (Grönlien, 1932) Gerasimov, 1952: 258
‡ *Stigmella floslactella* f. *interrupta* Dufrane, 1949: 10

Stigmella johanssonella A. Laštuvka & Z. Laštuvka, 1997: 70 WP

Stigmella tityrella (Stainton, 1854) Hering, 1957: 439 WP
Nepticula tityrella Stainton, 1854: 304
Nepticula turicella Herrich-Schäffer, 1855a: 355 (syn: Carolsfeld-Krausé, 1949: 310)
Nepticula turicensis Frey, 1856: 391 UE
Nepticula castanella Stainton, 1859a: 123 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula hemargyrella auct. [misapplied] (syn: Carolsfeld-Krausé, 1949: 304)
Stigmella turicella (Herrich-Schäffer, 1855a) Fletcher & Clutterbuck, 1945: 60
Stigmella castanella (Stainton, 1859a) Gerasimov, 1952: 232

***Stigmella ruficapitella* group - s.l.** (Johansson, 1971: 245) [39]
Stigmella hemargyrella group (Johansson, 1971: 244)
Stigmella procrastinella group (Wilkinson & Scoble, 1979: 69)
Stigmella castanopsiella group (Puplesis, 1984b: 583)

Stigmella hemargyrella (Kollar, 1832) Gerasimov, 1952: 242 WP
Oecophora hemargyrella Kollar, 1832: 98
Nepticula basalella Herrich-Schäffer, 1855a: 354 (syn: Carolsfeld-Krausé, 1949: 310)
Nepticula fagella Herrich-Schäffer, 1855a: 354 (syn: van Nieukerken & Johansson, 1987: 461)

Nepticula fagi Frey, 1856: 384 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula nobilella Heinemann & Wocke, [1876]: 755 [6] (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula fulgens Stainton, 1888b: 12 (syn: Carolsfeld-Krausé, 1949: 310)
Nepticula tityrella auct. [misapplied] (see: Carolsfeld-Krausé, 1949: 307)
Lyonetia hemargyrella (Kollar, 1832) Zeller, 1839: 215
Nepticula hemargyrella (Kollar, 1832) Zeller, 1848: 323
Stigmella basalella (Herrich-Schäffer, 1855) Fletcher & Clutterbuck, 1945: 60

Stigmella castanopsiella (Kuroko, 1978) Kuroko, 1982a: 448 EP
Nepticula castanopsiella Kuroko, 1978: 1

Stigmella kurokoi Puplesis, 1984b: 594 EP
Stigmella valvaurigemmata Kemperman & Wilkinson, 1985: 45 (syn: Puplesis, 1994: 161)

Stigmella lithocarpella van Nieukerken & Liu, 2000: 169 EP

Stigmella vandrieli van Nieukerken & Liu, 2000: 171 EP

Stigmella circumargentea van Nieukerken & Liu, 2000: 165 EP

Stigmella kao van Nieukerken & Liu, 2000: 166 EP,OR

Stigmella alba Wilkinson & Scoble, 1979: 73 NEA

Stigmella procrastinella (Braun, 1927) Wilkinson & Scoble, 1979: 70 NEA
Nepticula procrastinella Braun, 1927: 59

Stigmella humboldti Remeikis & Stonis, 2015: 412 [46] NEO

***Stigmella ruficapitella* group - s.s.** (Johansson, 1971: 245) [39]
Stigmella atricapitella group (Emmet, 1976: 239)
Stigmella caesurifasciella group (Kemperman & Wilkinson, 1985: 38)
Stigmella suberivora group (Kemperman & Wilkinson, 1985: 38)

Stigmella atricapitella (Haworth, 1828) Beirne, 1945: 198 WP
Tinea atricapitella Haworth, 1828: 585
Nepticula discrepans Sorhagen, 1922: 41 (syn: van Nieukerken & Johansson, 1987: 461)
Microsetia atricapitella (Haworth, 1828) Stephens, 1834: 269
Nepticula atricapitella (Haworth, 1828) Stainton, 1849: 28
Stigmella discrepans (Sorhagen, 1922) Gerasimov, 1952: 237

Stigmella basiguttella (Heinemann, 1862) Vári, 1944b: xxv WP
Nepticula basiguttella Heinemann, 1862a: 258
Stigmella cerricolella Klimesch, 1948a: 160 [29] (syn: Johansson, 1971: 253)
Nepticula cerricolella (Klimesch, 1948) Johansson, 1971: 253
Nepticula basiguttella cerricolella (Klimesch, 1948) Johansson, 1971: 253

Stigmella bicuspidata van Nieukerken & Johansson, 2003: 341 WP

Stigmella cocciferae van Nieukerken & Johansson, 2003: 329 WP

Stigmella dorsiguttella (Johansson, 1971) Pröse, 1984: 107 WP
Nepticula dorsiguttella Johansson, 1971: 251

Stigmella eberhardi (Johansson, 1971) Kasy, 1979: 4 WP
Nepticula eberhardi Johansson, 1971: 258

Stigmella fasciata van Nieukerken & Johansson, 2003: 321 WP
Stigmella ilicifoliella (Mendes, 1918) Gómez Bustillo, 1981: 18 WP
Nepticula ilicifoliella Mendes, 1918: 127
Stigmella ilicivora nigra Dufrane, 1955: 192 (syn: van Nieukerken & Johansson, 2003: 326)
Stigmella karsholti van Nieukerken & Johansson, 2003: 343 WP
Stigmella macrolepidella (Klimesch, 1978) van Nieukerken, 1986b: 13 WP
Nepticula macrolepidella Klimesch, 1978b: 257
Stigmella roborella (Johansson, 1971) Emmet, 1976: 241 WP
Nepticula roborella Johansson, 1971: 258
Nepticula ruficapitella auct. partim before 1971
Stigmella ruficapitella (Haworth, 1828) Beirne, 1945: 198 WP
Tinea ruficapitella Haworth, 1828: 586
Tinea violacella Haworth, 1828: 585 (syn: Stainton, 1849: 28)
Microsetia ruficapitella (Haworth, 1828) Stephens, 1834: 269
Nepticula ruficapitella (Haworth, 1828) Stainton, 1849: 28
Microsetia violaceella (Haworth, 1828) Stephens, 1834: 269
‡ *Nepticula lamprotornella* Heyden in Herrich-Schäffer, 1855b: 69 NN (syn: Herrich-Schäffer, 1855b: 69)
‡ *Nepticula ruficapitella* var. *ruficastaneae* Skala, 1949: 129 NNLM
Stigmella samiatella (Zeller, 1839) Vári, 1950: 180 [47] WP
Lyonetia samiatella Zeller, 1839: 215
Nepticula chaoniella Herrich-Schäffer, 1863b: 170 NO **syn. n.** [47]
Nepticula quercella Herrich-Schäffer, 1863a: 23 NN (syn: Segerer, 1997: 190) [47]
Nepticula samiatella (Zeller, 1839) Zeller, 1848: 303
Stigmella suberivora (Stainton, 1869) Beirne, 1945: 197 WP
Nepticula suberivora Stainton, 1869b: 228
Nepticula aureocapitella Millière, 1876 (syn: van Nieukerken & Johansson, 1987: 461)
Nepticula aureocaputella Millière, 1876: 374 IOS
Nepticula ilicivora Peyerimhoff, 1871: 413 (syn: Johansson, 1971: 246)
Nepticula ilicella Walsingham, 1891: 152 (syn: van Nieukerken, 2004: 111)
Stigmella ilicivora (Peyerimhoff, 1871) Gerasimov, 1952: 244
Stigmella ilicella (Walsingham, 1891) Le Marchand, 1946b: 284
Stigmella svenssoni (Johansson, 1971) Emmet, 1976: 243 WP
Nepticula svenssoni Johansson, 1971: 249
Stigmella szoecsiella (Borkowski, 1972) van Nieukerken, 1986a: 13 WP
Nepticula szoecsiella Borkowski, 1972b: 776
Stigmella tristis (Wocke, 1862) Gerasimov, 1952: 265 WP
Nepticula tristis Wocke, 1862: 251
Stigmella trojana Z. Laštuvka & A. Laštuvka, 1998: 313 WP
Stigmella zangherii (Klimesch, 1951) Zangheri, 1969: 1014 WP
Nepticula zangherii Klimesch, 1951d: 61

Stigmella acuta Diškus, Navickaitė & Remeikis in Stonis et al., 2013e: 202 EP
Stigmella aladina Puplesis, 1984a: 115 EP
Stigmella quercifaga Kemperman & Wilkinson, 1985: 44 (syn: van Nieukerken & Liu, 2000: 161)
Stigmella azuminoensis Hirano, 2010: 125 EP
Stigmella caesurifasciella Kemperman & Wilkinson, 1985: 38 EP
Stigmella egregilustrata Kemperman & Wilkinson, 1985: 39 (syn: van Nieukerken & Liu, 2000: 174)
Stigmella clisiotophora Kemperman & Wilkinson, 1985: 48 EP
Stigmella crenatiella Hirano, 2010: 125 EP
Stigmella dentatae Puplesis, 1984a: 114 EP
Stigmella pulla Kemperman & Wilkinson, 1985: 43 (syn: Puplesis, 1994: 162)
Stigmella fervida Puplesis, 1984b: 593 EP
Stigmella fumida Kemperman & Wilkinson, 1985: 42 EP
Stigmella chrysopterella Kemperman & Wilkinson, 1985: 48 (syn: van Nieukerken & Liu, 2000: 157)
Stigmella kurii Kemperman & Wilkinson, 1985: 51 (syn: van Nieukerken & Liu, 2000: 157)
Stigmella hisaii Kuroko, 2004: 238 EP
Stigmella hisakoae Hirano, 2010: 126 EP
Stigmella kasyi van Nieukerken & Johansson, 2003: 331 EP
Stigmella omelkoi Puplesis, 1984b: 593 EP
Stigmella kumatai Kemperman & Wilkinson, 1985: 50 (syn: Puplesis, 1994: 163)

Unplaced Species groups

***Stigmella barbata* group** (Puplesis et al., 2002: 63)
Stigmella plumosetaeella Newton & Wilkinson, 1982: 455 NEA
Stigmella austroamericana Puplesis & Diškus in Puplesis et al., 2002: 25 NEO
Stigmella barbata Puplesis & Robinson, 2000: 37 NEO

***Stigmella purpurimaculae* group** (Remeikis & Stonis in Stonis et al., 2014: 351)
Stigmella cana Remeikis & Stonis in Stonis et al., 2014: 324 NEO
Stigmella concreta Remeikis & Stonis in Stonis et al., 2014: 328 NEO
Stigmella pseudoconcreta Remeikis & Stonis in Stonis et al., 2014: 329 NEO
Stigmella purpurimaculae Remeikis & Stonis in Stonis et al., 2014: 323 NEO
Stigmella quadrata Remeikis & Stonis in Stonis et al., 2014: 329 NEO
Stigmella sceptra Remeikis & Stonis in Stonis et al., 2014: 327 NEO
Stigmella truncata Remeikis & Stonis in Stonis et al., 2014: 326 NEO

***Stigmella* unplaced and ungrouped species**
Stigmella arbatella (Chrétien, 1922) Rungs, 1979: 25 WP
Nepticula arbatella Chrétien, 1922: 373
Stigmella georgiana Puplesis, 1994: 165 WP

Stigmella grandistyla Puplesis, 1994: 113 WP
Stigmella brutea Remeikis & Stonis in Stonis et al., 2014: 331 NEO
Stigmella hylomaga (Meyrick, 1931a) Davis, 1984: 18 NEO
Nepticula hylomaga Meyrick, 1931a: 415
Stigmella pruinosa Puplesis & Robinson, 2000: 38 NEO
Stigmella pseudodigitata Remeikis & Stonis in Stonis et al., 2014: 332 NEO
Stigmella semilactea Remeikis & Stonis in Stonis et al., 2014: 330 NEO

Ozadelpha van Nieukerken in van Nieukerken et al., 2016: 26 (TS/OD: *Ozadelpha conostegiae* van Nieukerken & Nishida, 2016)
Ozadelpha conostegiae van Nieukerken & Nishida in van Nieukerken et al., 2016: 28 NEO
Ozadelpha guajavae (Puplesis & Diškus, 2002) van Nieukerken et al., 2016: 27 NEO
Enteucha guajavae Puplesis & Diškus in Puplesis et al., 2002: 22
Ozadelpha ovata (Puplesis & Robinson, 2000) van Nieukerken et al., 2016: 27 NEO
Stigmella ovata Puplesis & Robinson, 2000: 39

Roscidotoga Hoare, 2000a: 293 (TS/OD: *Roscidotoga callicomae* Hoare, 2000)
Roscidotoga callicomae Hoare, 2000a: 295 (Fig. 27) AUS
Roscidotoga eucryphiae Hoare, 2000a: 296 AUS
Roscidotoga lamingtonia van Nieukerken, van den Berg & Hoare, 2011: 194 AUS
Roscidotoga sapphiripes Hoare, 2000a: 297 AUS

Casanovula Hoare in Hoare & van Nieukerken, 2013: 24 (TS/OD: *Pectinivalva brevipalpa* Hoare, 2013), **stat. n.**
Casanovula brevipalpa (Hoare, 2013), **comb. n.** AUS
Pectinivalva brevipalpa Hoare in Hoare & van Nieukerken, 2013: 27
Casanovula minotaurus (Hoare, 2013), **comb. n.** AUS
Pectinivalva minotaurus Hoare in Hoare & van Nieukerken, 2013: 29

Menurella Hoare in Hoare & van Nieukerken, 2013: 35 (TS/OD: *Pectinivalva scotodes* Hoare, 2013), **stat. n.**
Menurella acmenae (Hoare, 2013), **comb. n.** AUS
Pectinivalva acmenae Hoare in Hoare & van Nieukerken, 2013: 41
Menurella anazona (Meyrick, 1906), **comb. n.** AUS
Nepticula anazona Meyrick, 1906b: 58
Pectinivalva anazona (Meyrick, 1906) Nielsen, 1996: 16
Menurella funeralis (Meyrick, 1906), **comb. n.** AUS
Nepticula funeralis Meyrick, 1906b: 59
Pectinivalva funeralis (Meyrick, 1906) Nielsen, 1996: 16
Menurella libera (Meyrick, 1906), **comb. n.** (Fig. 28) AUS
Nepticula libera Meyrick, 1906b: 61

Pectinivalva libera (Meyrick, 1906) Nielsen, 1996: 16
Menurella planetis (Meyrick, 1906), **comb. n.** AUS
Nepticula planetis Meyrick, 1906b: 58
Pectinivalva planetis (Meyrick, 1906) Nielsen, 1996: 16
Menurella primigena (Meyrick, 1906), **comb. n.** AUS
Nepticula primigena Meyrick, 1906b: 58
Pectinivalva primigena (Meyrick, 1906) Nielsen, 1996: 16
Menurella quintiniae (Hoare & van Nieukerken, 2013), **comb. n.** AUS
Pectinivalva quintiniae Hoare & van Nieukerken, 2013: 47
Menurella scotodes (Hoare, 2013), **comb. n.** AUS
Pectinivalva scotodes Hoare in Hoare & van Nieukerken, 2013: 37
Menurella trepida (Meyrick, 1906), **comb. n.** AUS
Nepticula trepida Meyrick, 1906b: 61
Pectinivalva trepida (Meyrick, 1906) Nielsen, 1996: 16
Menurella tribulatrix (van Nieukerken & Hoare, 2013), **comb. n.** AUS
Pectinivalva tribulatrix van Nieukerken & Hoare in Hoare & van Nieukerken, 2013: 48
Menurella warburtonensis (Wilson, 1939), **comb. n.** AUS
Nepticula warburtonensis Wilson, 1939: 239
Pectinivalva warburtonensis (Wilson, 1939) Nielsen, 1996: 16
Menurella xenadelpha (van Nieukerken & Hoare, 2013), **comb. n.** OR
Pectinivalva xenadelpha van Nieukerken & Hoare in Hoare & van Nieukerken, 2013: 44

Pectinivalva Scoble, 1983: 12
Pectinivalva caenodora (Meyrick, 1906) Nielsen, 1996: 16 (Fig. 29) AUS
Nepticula caenodora Meyrick, 1906b: 58
Pectinivalva chalcitis (Meyrick, 1906) Nielsen, 1996: 16 AUS
Nepticula chalcitis Meyrick, 1906b: 60
Pectinivalva commoni Scoble, 1983: 13 AUS
Pectinivalva endocapna (Meyrick, 1906) Nielsen, 1996: 16 AUS
Nepticula endocapna Meyrick, 1906b: 60
Pectinivalva gilva (Meyrick, 1906b) Nielsen, 1996: 16 AUS
Nepticula gilva Meyrick, 1906b: 59
Pectinivalva melanotis (Meyrick, 1906) Nielsen, 1996: 16 AUS
Nepticula melanotis Meyrick, 1906b: 59
Pectinivalva mystaconota Hoare in Hoare & van Nieukerken, 2013: 20 AUS

Neotrifurcula van Nieukerken, 2016 in van Nieukerken et al., 2016: 36 (TS/OD: *Neotrifurcula gielisorum* van Nieukerken, 2016)
Neotrifurcula gielisorum van Nieukerken, 2016 in van Nieukerken et al., 2016: 38 NEO

Hesperolyra van Nieukerken, 2016 in van Nieukerken et al., 2016: 44 (TS/OD: *Fomoria diskusi* Puplesis & Robinson, 2000)
Fomoria molybditis group Puplesis et al., 2002
Hesperolyra diskusi (Puplesis & Robinson, 2000) van Nieukerken et al., 2016: 44 NEO
Fomoria diskusi Puplesis & Robinson, 2000: 43
Hesperolyra molybditis (Zeller, 1877) van Nieukerken et al., 2016: 46 NEO
Nepticula molybditis Zeller, 1877: 453
Stigmella molybditis (Zeller, 1877) Davis, 1984: 18
Fomoria molybditis (Zeller, 1877) Puplesis & Robinson, 2000: 43
Hesperolyra repanda (Puplesis & Diškus, 2002) van Nieukerken et al., 2016: 46 NEO
Fomoria repanda Puplesis & Diškus in Puplesis et al., 2002: 26
Hesperolyra saopaulensis van Nieukerken, 2016 in van Nieukerken et al., 2016: 52 NEO

Bohemannia Stainton, 1859a: 439 (TS/M: *Nepticula quadrimaculella* Boheman, 1853)
Scoliaula Meyrick, 1895: 727; URN for *Bohemannia* Stainton, 1859 (TS/OD,M: *Nepticula quadrimaculella* Boheman, 1853)
Bohemannia aschaueri Fischer, 2013: 88 WP†
Bohemannia butzmanni Fischer, 2013: 86 WP†
Bohemannia auriciliella (Joannis, 1909) van Nieukerken, 1986a: 16 (Fig. 31) WP
Nepticula auriciliella Joannis, 1909: 822
Ectoedemia bradfordi Emmet, 1974: 269 (syn: van Nieukerken, 1986a: 16)
Stigmella auriciliella (Joannis, 1909) Lhomme, [1963]: 1192
Bohemannia pulverosella (Stainton, 1849) van Nieukerken, 1982: 105 [48] WP,EP,[NEA]
Trifurcula pulverosella Stainton, 1849: 30
Bohemannia piotra Puplesis, 1984b: 586 **syn. n.**[48]
Nepticula pulverosella (Stainton, 1849) Meyrick, 1895: 726
Stigmella pulverosella (Stainton, 1849) Fletcher & Clutterbuck, 1945: 61
Dechtiria pulverosella (Stainton, 1849) Beirne, 1945: 206
Ectoedemia pulverosella (Stainton, 1849) Bradley et al., 1972: 3
Scoliaula pulverosella (Stainton, 1849) Borkowski, 1975: 489
‡ *Nepticula cineretella* Frey in Herrich-Schäffer, 1855b: 70 NN (syn: Herrich-Schäffer, 1855b: 70)
Bohemannia quadrimaculella (Boheman, 1853) Stainton, 1859a: 439 [49] WP
Nepticula quadrimaculella Boheman, 1853: 167
Bucculatrix antispilella Meess, 1907: 129 (syn: Disqué, 1912: 75) [49]
Scoliaula quadrimaculella (Boheman, 1853) Meyrick, 1895: 727
Trifurcula quadrimaculella (Boheman, 1853) Johansson, 1971: 246
Bohemannia ussuriella Puplesis, 1984b: 588 EP
Bohemannia manschurella Puplesis, 1984b: 587 [50] EP
Bohemannia nipponicella Hirano, 2010: 129 **syn. n.** [50]

Bohemannia nubila Puplesis, 1985c: 69 EP
Bohemannia suiphunella Puplesis, 1984b: 588 EP

Areticulata Scoble, 1983: 11 (key), 40 (TS/OD,M: *Areticulata leucosideae* Scoble, 1983)
Areticulata leucosideae Scoble, 1983: 40 AFR

Glaucolepis Braun, 1917: 161 (key), 201 (TS/OD,M: *Nepticula saccharella* Braun, 1912)
Fedalmia Beirne, 1945: 207 (TS/OD,M: *Nepticula headleyella* Stainton, 1854) (syn: Puplesis, 1985a: 11)
Sinopticula Yang, 1989: 79 [81] (TS/OD,M: *Sinopticula sinica* Yang, 1989: 80) (syn: van Nieukerken & Puplesis, 1991: 202)

***Glaucolepis raikhonae* group** (van Nieukerken & Puplesis, 1991: 202)
Glaucolepis melanoptera (van Nieukerken & Puplesis, 1991) Puplesis, 1994: 219 WP
Trifurcula melanoptera van Nieukerken & Puplesis, 1991
Glaucolepis oishiella (Matsumura, 1931), **comb. n.** [51] EP
Trifurcula oishiella Matsumura, 1931: 1114
Sinopticula sinica Yang, 1989: 80 **syn. n.** [51]
Trifurcula sinica (Yang, 1989) van Nieukerken & Puplesis, 1991: 205
Glaucolepis sinica (Yang, 1989) Diškus & Puplesis, 2003: 404
Glaucolepis raikhonae Puplesis, 1985c: 71 EP
Trifurcula raikhonae (Puplesis, 1985) van Nieukerken, 1986a: 68

***Glaucolepis saccharella* group** (new)
Glaucolepis saccharella (Braun, 1912) Braun, 1917: 201 NEA
Nepticula saccharella Braun, 1912: 97
Trifurcula saccharella (Braun, 1912) van Nieukerken, 1986a: 68

***Glaucolepis headleyella* group** (Puplesis, 1994: 219)
Glaucolepis albiflorella (Klimesch, 1978) Diškus & Puplesis, 2003: 406 WP
Trifurcula albiflorella Klimesch, 1978b: 274
Glaucolepis alypella (Klimesch, 1975) Diškus & Puplesis, 2003: 406 WP
Trifurcula alypella Klimesch, 1975c: 12
Glaucolepis andalusica (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula andalusica Z. Laštuvka & A. Laštuvka, 2007: 102
Glaucolepis bleonella (Chrétien, 1904) Puplesis, 1994: 219 WP
Nepticula bleonella Chrétien, 1904: 164
Stigmella bleonella (Chrétien, 1904) Gerasimov, 1952: 231
Ectoedemia bleonella (Chrétien, 1904) Klimesch, 1975a: 861
Trifurcula bleonella (Chrétien, 1904) Leraut, 1980: 49
Glaucolepis bupleurella (Chrétien, 1907) Diškus & Puplesis, 2003: 405 WP
Nepticula bupleurella Chrétien, 1907: 91
Stigmella bupleurella (Chrétien, 1907) Gerasimov, 1952: 232

Ectoedemia bupleurella (Chrétien, 1907) Klimesch, 1975a: 863
Trifurcula bupleurella (Chrétien, 1907) Leraut, 1980: 49

Glaucolepis chretieni (Z. Laštůvka, A. Laštůvka & van Nieukerken, 2013), **comb. n.** WP
Trifurcula chretieni Z. Laštůvka, A. Laštůvka & van Nieukerken, 2013: 198

Glaucolepis corleyi (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula corleyi Z. Laštuvka & A. Laštuvka, 2007: 102

Glaucolepis globulariae (Klimesch, 1975) Diškus & Puplesis, 2003: 406 WP
Trifurcula globulariae Klimesch, 1975c: 7

Glaucolepis hamirella (Chrétien, 1915) Diškus & Puplesis, 2003: 406 [52] WP
Nepticula hamirella Chrétien, 1915: 364
Ectoedemia hamirella (Chrétien, 1915) Klimesch, 1975a: 863
Trifurcula hamirella (Chrétien, 1915) van Nieukerken, 1986a: 15

Glaucolepis saturejae (Parenti, 1963) Diškus & Puplesis, 2003: 406 [52] WP
Stigmella saturejae Parenti, 1963: 101
Fedalmia saturejae (Parenti, 1963) Klimesch, 1976: 45
Trifurcula saturejae (Parenti, 1963) van Nieukerken, 1986a: 15

Glaucolepis headleyella (Stainton, 1854) Puplesis, 1994: 219 WP
Nepticula headleyella Stainton, 1854: 298
Nepticula argyrostigma Frey, 1856: 379 (syn: Frey, 1880: 425)
Nepticula dubiella Hauder, 1912: 273 (syn: Klimesch, 1948b: 76)
Trifurcula rodella Svensson, 1982: 299 (syn: van Nieukerken, 1986a: 15)
Fedalmia headleyella (Stainton, 1854) Beirne, 1945: 207
Stigmella headleyella (Stainton, 1854) Klimesch, 1948b: 76
Trifurcula headleyella (Stainton, 1854) Johansson, 1971: 245
Stigmella dubiella (Hauder, 1912) Klimesch, 1948b: 76

Glaucolepis helladica (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula helladica Z. Laštuvka & A. Laštuvka, 2007: 101

Glaucolepis istriae (A. Laštuvka & Z. Laštuvka, 2000) Diškus & Puplesis, 2003: 406 WP
Trifurcula istriae A. Laštuvka & Z. Laštuvka, 2000a: 290

Glaucolepis kalavritana (Z. Laštuvka & A. Laštuvka, 1998) Diškus & Puplesis, 2003: 407 WP
Trifurcula kalavritana (Z. Laštuvka & A. Laštuvka, 1998): 314

Glaucolepis lavandulae (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula lavandulae Z. Laštuvka & A. Laštuvka, 2007: 104

Glaucolepis liskai (A. Laštuvka & Z. Laštuvka, 2000) Diškus & Puplesis, 2003: 407 WP
Trifurcula liskai A. Laštuvka & Z. Laštuvka, 2000a: 291

Glaucolepis lituanica (Ivinskis & van Nieukerken, 2012), **comb. n.** (Fig. 30) WP
Trifurcula lituanica Ivinskis & van Nieukerken, 2012: 43

Glaucolepis magna (A. Laštuvka & Z. Laštuvka, 1997) Diškus & Puplesis, 2003: 407 [53] WP

Trifurcula magna (A. Laštuvka & Z. Laštuvka, 1997): 132
Trifurcula collinella Nel, 2012: 24 **syn. n.**[53]

Glaucolepis megaphallus (van Nieukerken, Z. Laštůvka & A. Laštůvka, 2013), **comb. n.** WP
Trifurcula megaphallus van Nieukerken, Z. Laštůvka & A. Laštůvka in Z. Laštůvka et al., 2013: 195

Glaucolepis micromeriae (Walsingham, 1908) Diškus & Puplesis, 2003: 405 WP
Stigmella micromeriae Walsingham, 1908a: 1010[1]
Nepticula micromeriae (Walsingham, 1908) Rebel, 1910: 374
Trifurcula micromeriae (Walsingham, 1908) Klimesch, 1977: 196

Glaucolepis montana (Z. Laštuvka, A. Laštuvka & van Nieukerken), **comb. n.** WP
Trifurcula montana Z. Laštuvka, A. Laštuvka & van Nieukerken in Z. & A. Laštuvka, 2007: 103

Glaucolepis pederi (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula pederi Z. Laštuvka & A. Laštuvka, 2007: 102

Glaucolepis rosmarinella (Chrétien, 1914) Diškus & Puplesis, 2003: 405 WP
Nepticula rosmarinella Chrétien, 1914: 270
Stigmella rosmarinella (Chrétien, 1914) Gerasimov, 1952: 256
Trifurcula rosmarinella (Chrétien, 1914) Klimesch, 1975b: 23

Glaucolepis salicinae (Klimesch, 1975) Diškus & Puplesis, 2003: 406 WP
Trifurcula salicinae Klimesch, 1975c: 10

Glaucolepis salvifoliae (Z. Laštuvka & A. Laštuvka, 2007), **comb. n.** WP
Trifurcula salvifoliae Z. Laštuvka & A. Laštuvka, 2007: 103

Glaucolepis sanctaecrucis (Walsingham, 1908) Diškus & Puplesis, 2003: 405 WP
Stigmella sanctaecrucis Walsingham, 1908a: 1010[1]
Nepticula sanctaecrucis (Walsingham, 1908) Rebel, 1910: 374
Fedalmia sanctaecrucis (Walsingham, 1908) Klimesch, 1976: 44
Trifurcula sanctaecrucis (Walsingham, 1908) Klimesch, 1977: 196

Glaucolepis sanctibenedicti (Klimesch, 1979) Diškus & Puplesis, 2003: 406 WP
Trifurcula sanctibenedicti Klimesch, 1979: 24

Glaucolepis siciliae (Z. Laštůvka, A. Laštůvka & van Nieukerken, 2013), **comb. n.** WP
Trifurcula siciliae Z. Laštůvka, A. Laštůvka & van Nieukerken, 2013: 201

Glaucolepis stoechadella (Klimesch, 1975) Diškus & Puplesis, 2003: 406 WP
Trifurcula stoechadella Klimesch, 1975c: 23

Glaucolepis teucriella (Chrétien, 1914) Diškus & Puplesis, 2003: 405 WP
Nepticula teucriella Chrétien, 1914: 270
Stigmella teucriella (Chrétien, 1914) Gerasimov, 1952: 263
Trifurcula teucriella (Chrétien, 1914) Leraut, 1980: 49

Glaucolepis thymi (Szőcs, 1965) Diškus & Puplesis, 2003: 406 WP
Nepticula thymi Szőcs, 1965: 89
Fedalmia thymi Borkowski, 1970a: 74; JSH of *Nepticula thymi* Szőcs, 1965
Trifurcula thymi (Szőcs, 1965) van Nieukerken, 1986a: 15

Glaucolepis trilobella (Klimesch, 1978) Diškus & Puplesis, 2003: 406 WP
Trifurcula trilobella Klimesch, 1978b: 271
Glaucolepis zollikofferiela (Chrétien, 1914) Diškus & Puplesis, 2003: 405 WP
Nepticula zollikofferiela Chrétien, 1914: 271
Stigmella zollikofferiela (Chrétien, 1914) Gerasimov, 1952: 270
Ectoedemia zollikofferiela (Chrétien, 1914) Klimesch, 1975a: 862
Trifurcula zollikofferiela (Chrétien, 1914) van Nieukerken, 1986a: 15
Glaucolepis rusticula (Meyrick, 1916) Diškus & Puplesis, 2003: 406 OR
Nepticula rusticula Meyrick, 1916b: 7
Trifurcula rusticula (Meyrick, 1916) online comb.

Unassigned to group [54]
Glaucolepis aerifica (Meyrick, 1915) Puplesis & Robinson, 2000: 56 NEO
Nepticula aerifica Meyrick, 1915a: 255
Stigmella aerifica (Meyrick, 1915) Davis, 1984: 18
Trifurcula aerifica (Meyrick, 1915) online comb.
Glaucolepis argentosa Puplesis & Robinson, 2000: 57 [54] NEO
Trifurcula argentosa (Puplesis & Robinson, 2000) online comb.

Trifurcula Zeller, 1848: 249 (key), 330 (TS/SD (Tutt, 1899: 355): *Trifurcula pallidella* Zeller, 1848)
Trifurcella Chambers, 1878: 165 ISS
Levarchama Beirne, 1945: 206 (TS/OD: *Nepticula cryptella* Stainton, 1856) (syn: Johansson, 1971: 246)

***Trifurcula cryptella* group** (new)
Trifurcula anthyllidella Klimesch, 1975c: 14 WP
Trifurcula cryptella (Stainton, 1856) Johansson, 1971: 246 WP
Nepticula cryptella Stainton, 1856: 41
Nepticula trifolii Sorhagen, 1885: 280 (syn: Hering, 1957: 1067)
Stigmella cryptella (Stainton, 1856) Fletcher & Clutterbuck, 1945: 61
Levarchama cryptella (Stainton, 1856) Beirne, 1945: 207
Trifurcula eurema (Tutt, 1899) Johansson, 1971: 246 WP
Nepticula eurema Tutt, 1899: 332
Nepticula heurema Meess, 1910: 481 UE
Nepticula dorycniella Suire, 1928: 128 (syn: van Nieukerken, 1986a: 15)
Nepticula gozmanyi Szőcs, 1959: 417 (syn: van Nieukerken, 1986a: 15)
Levarchama eurema (Tutt, 1899) Beirne, 1945: 207
Stigmella eurema (Tutt, 1899) Klimesch, 1951b: 66
Stigmella heurema (Meess, 1910) Gerasimov, 1952: 243
Stigmella dorycniella (Suire, 1928) Klimesch, 1951b: 66
Trifurcula manygoza van Nieukerken, A. Laštuvka & Z. Laštuvka in van Nieukerken, 2007b: 125 WP

Trifurcula ortneri (Klimesch, 1951) van Nieukerken, 1986a: 15 WP
Nepticula ortneri Klimesch, 1951b: 66
Stigmella ortneri (Klimesch, 1951) Klimesch, 1961: 763
Trifurcula peloponnesica van Nieukerken, 2007b: 118 WP
Trifurcula ridiculosa (Walsingham, 1908) Klimesch, 1975b: 15 WP
Stigmella ridiculosa Walsingham, 1908a: 1011[1]
Nepticula ridiculosa (Walsingham, 1908) Rebel, 1910: 364

***Trifurcula subnitidella* group** (van Nieukerken, 1990b: 208)
Trifurcula coronillae van Nieukerken, 1990b: 217 WP
Trifurcula iberica van Nieukerken, 1990b: 228 (Fig. 32) WP
Trifurcula josefklimeschi van Nieukerken, 1990b: 225 WP
Trifurcula luteola van Nieukerken, 1990b: 215 WP
Trifurcula puplesisi van Nieukerken, 1990b: 215 WP,EP
Trifurcula silviae van Nieukerken, 1990b: 230 WP
Trifurcula subnitidella (Duponchel, 1843) van Nieukerken & Johansson, 1987: 462 WP
Elachista subnitidella Duponchel, 1843: 326
Trifurcula griseella Wolff, 1957: 21 (syn: van Nieukerken & Johansson, 1987: 462)
Lyonetia subnitidella (Duponchel, 1843) Duponchel, 1844: 378
Nepticula subnitidella (Duponchel, 1843) Zeller, 1848: 305
Trifurcula victoris van Nieukerken, 1990b: 219 WP

***Trifurcula pallidella* group** (van Nieukerken, 1990b: 208)
Trifurcula aurella Rebel, 1933: 82 WP
Trifurcula austriaca van Nieukerken, 1990b: 213 WP
Trifurcula baldensis A. Laštuvka & Z. Laštuvka, 2005a: 8 WP
Trifurcula beirnei Puplesis, 1984a: 124 WP
Trifurcula bicolorella (Chrétien, 1915), **comb. n.** [55] WP
Bucculatrix bicolorella Chrétien, 1915: 364
Trifurcula calycotomella A. Laštuvka & Z. Laštuvka, 1997: 148 WP
Trifurcula chamaecytisi A. Laštuvka & Z. Laštuvka, 1994: 207 WP
Trifurcula corothamni A. Laštuvka & Z. Laštuvka, 1994: 202 WP
Trifurcula cytisanthi A. Laštuvka & Z. Laštuvka, 2005a: 8 WP
Trifurcula etnensis A. Laštuvka & Z. Laštuvka, 2005a: 7 WP
Trifurcula graeca Z. Laštuvka & A. Laštuvka, 1998: 315 WP
Trifurcula immundella (Zeller, 1839) Zeller, 1848: 332 WP
Lyonetia immundella Zeller, 1839: 215
Trifurcula macedonica Z. Laštuvka & A. Laštuvka, 1998: 315 WP
Trifurcula moravica A. Laštuvka & Z. Laštuvka, 1994: 205 WP
Trifurcula orientella Klimesch, 1953a: 168 WP
Trifurcula pallidella (Duponchel, 1843) Joannis, 1915: 129 WP
Oecophora pallidella Duponchel, 1843: 339

Trifurcula pallidella Zeller, 1848: 332; JSH of *Trifurcula pallidella* (Duponchel, 1843)
Lithocolletis pallidella (Duponchel, 1843) Bruand, [1851]: 86
Trifurcula incognitella Toll, 1936: 409 (syn: van Nieukerken, 1986a: 15)
‡ *[no genus] pallidulella* Herrich-Schäffer, 1853: pl 108 NN
Trifurcula serotinella Herrich-Schäffer, 1855a: 359 WP
Trifurcula confertella Fuchs, 1895: 47 (syn: van Nieukerken, 1986a: 15)
Trifurcula squamatella Stainton, 1849: 30 WP
Trifurcula maxima Klimesch, 1953a: 167 (syn: van Nieukerken, 1987b: 180)
Trifurcula trasaghica A. Laštuvka & Z. Laštuvka, 2005a: 9 WP

***Trifurcula barbertonensis* group** (new)
Trifurcula barbertonensis Scoble, 1980a: 142 AFR
Trifurcula pullus Scoble, 1980a: 140 AFR

Fomoria Beirne, 1945: 208 (TS/OD: *Nepticula weaveri* Stainton, 1855)

***Fomoria vannifera* group** (Hoare, 2000b: 300)
Fomoria asiatica group (Puplesis, 1994: 208)
Fomoria asiatica Puplesis, 1988: 27 EP
Ectoedemia asiatica (Puplesis, 1988) Hoare, 2000a: 301
Fomoria glycystrota (Meyrick, 1928) Diškus & Puplesis, 2003: 385 OR
Nepticula glycystrota Meyrick, 1928b: 462
Ectoedemia glycystrota (Meyrick, 1928b) Hoare, 2000a: 302
Fomoria fuscata (Janse, 1948) Diškus & Puplesis, 2003: 385 AFR
Nepticula fuscata Janse, 1948: 165
Ectoedemia fuscata (Janse, 1948) Scoble, 1983: 36
Fomoria hobohmi (Janse, 1948) Diškus & Puplesis, 2003: 385 AFR
Nepticula hobohmi Janse, 1948: 167
Ectoedemia hobohmi (Janse, 1948) Scoble, 1983: 38
Fomoria kharuxabi (Mey, 2004), **comb. n.** AFR
Ectoedemia kharuxabi Mey, 2004: 32
Fomoria uisebi (Mey, 2004), **comb. n.** AFR
Ectoedemia uisebi Mey, 2004: 32
Fomoria vannifera (Meyrick, 1914) Diškus & Puplesis, 2003: 385 AFR
Nepticula vannifera Meyrick, 1914: 203
Ectoedemia vannifera (Meyrick, 1914) Scoble, 1983: 37
Fomoria hadronycha (Hoare, 2000) Diškus & Puplesis, 2003: 385 AUS
Ectoedemia hadronycha Hoare, 2000b: 307
Fomoria pelops (Hoare, 2000) Diškus & Puplesis, 2003: 385 AUS
Ectoedemia pelops Hoare, 2000b: 304
Fomoria squamibunda (Hoare, 2000) Diškus & Puplesis, 2003: 385 AUS
Ectoedemia squamibunda Hoare, 2000b: 304

***Fomoria groschkei* group** (Hoare, 2000b: 313)

Fomoria aegaeica (Z. Laštuvka, A. Laštuvka & Johansson, 1998) Diškus & Puplesis, 2003: 391 WP

Ectoedemia aegaeica Z. Laštuvka, A. Laštuvka & Johansson in Z. & A. Laštuvka, 1998: 316

Fomoria groschkei (Skala, 1943) Diškus & Puplesis, 2003: 388 WP

Nepticula groschkei Skala, 1943: 86

Stigmella groschkei (Skala, 1943) Klimesch, 1948b: 77

Ectoedemia groschkei (Skala, 1943) van Nieukerken, 1986a: 17

Fomoria thermae (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR

Ectoedemia thermae Scoble, 1983: 36

***Fomoria weaveri* group** (Puplesis, 1994: 205)

Fomoria degeeri (van Nieukerken, 2008), **comb. n.** WP

Ectoedemia degeeri van Nieukerken, 2008: 124

Fomoria deschkai (Klimesch, 1978) Diškus & Puplesis, 2003: 384 WP

Trifurcula deschkai Klimesch, 1978b: 274

Ectoedemia deschkai (Klimesch, 1978b) van Nieukerken, 1986a: 17

Fomoria empetrifolii (A. Laštuvka & Z. Laštuvka, 2000) Diškus & Puplesis, 2003: 385 WP

Ectoedemia empetrifolii A. Laštuvka & Z. Laštuvka, 2000b: 22

Fomoria eriki (A. Laštuvka & Z. Laštuvka, 2000) Diškus & Puplesis, 2003: 385 WP

Ectoedemia eriki A. Laštuvka & Z. Laštuvka, 2000b: 21

Fomoria luisae Klimesch, 1978a: 89 WP

Ectoedemia luisae (Klimesch, 1978) van Nieukerken, 1986a: 17

Fomoria septembrella (Stainton, 1849) Beirne, 1945: 209 WP

Nepticula septembrella Stainton, 1849: 29

Stigmella septembrella (Stainton, 1849) Fletcher & Clutterbuck, 1945: 61

Trifurcula septembrella (Stainton, 1849) Johansson, 1971: 246

Ectoedemia septembrella (Stainton, 1849) Scoble, 1983: 32

Fomoria variicapitella (Chrétien, 1908) Diškus & Puplesis, 2003: 384 WP

Nepticula variicapitella Chrétien, 1908: 363

Stigmella variicapitella (Chrétien, 1908) Gerasimov, 1952: 263

Trifurcula variicapitella (Chrétien, 1908) Klimesch, 1977: 197

Ectoedemia variicapitella (Chrétien, 1908) van Nieukerken, 1986a: 17

Fomoria weaveri (Stainton, 1855) Beirne, 1945: 209 (Fig. 33) WP,EP,NEA

Nepticula weaveri Stainton, 1855: 49

Nepticula weaweri Herrich-Schäffer, 1855a: 346 ISS

Nepticula weaverella Doubleday, 1859: 36 UE

Stigmella weaveri (Stainton, 1855) Gerasimov, 1952: 269

Trifurcula weaveri (Stainton, 1855) Johansson, 1971: 246

Ectoedemia weaveri (Stainton, 1855) Scoble, 1983: 32

‡ *Fomoria weaveri* f. *fuliginella* Vári, 1947: 523

Fomoria festivitatis (van Nieukerken, 2008), **comb. n.** OR, EP
Ectoedemia festivitatis van Nieukerken, 2008: 117
Fomoria hypericifolia Kuroko, 1982: 49 EP
Ectoedemia hypericifolia (Kuroko, 1982) van Nieukerken, 1986a: 84
Fomoria permira Puplesis, 1984b: 592 EP
Ectoedemia permira (Puplesis, 1984) van Nieukerken, 1986a: 84
Fomoria hypericella (Braun, 1925) Wilkinson, 1979: 84 NEA
Nepticula hypericella Braun, 1925a: 17
Ectoedemia hypericella (Braun, 1925a) van Nieukerken, 2008: 116
Fomoria pteliaeella (Chambers, 1880) Wilkinson, 1979: 84 NEA
Nepticula pteliaeella Chambers, 1880c: 137
Ectoedemia pteliaeella (Chambers, 1880) van Nieukerken, 2008: 117
Fomoria ruwenzoriensis (Bradley, 1965), **comb. n.** AFR
Stigmella ruwenzoriensis Bradley, 1965: 120
Acalyptris ruwenzoriensis (Bradley, 1965) Diškus & Puplesis, 2003: 394
Ectoedemia ruwenzoriensis (Bradley, 1965) van Nieukerken, 2008: 117

Fomoria lacrimulae **group** (Diškus & Puplesis, 2003: 386)
Fomoria lacrimulae Puplesis & Diškus, 1996c: 185 WP
Ectoedemia lacrimulae (Puplesis & Diškus, 1996) online comb.
Fomoria knysnaensis (Scoble, 1983) Diškus & Puplesis, 2003: 386 AFR
Ectoedemia knysnaensis Scoble, 1983: 33

African unplaced *Fomoria*
Fomoria alexandria (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia alexandria Scoble, 1983: 35
Fomoria gambiana (Gustafsson, 1972) Diškus & Puplesis, 2003: 389 AFR
Nepticula gambiana Gustafsson, 1972: 156
Ectoedemia gambiana (Gustafsson, 1972) online comb.
Fomoria incisaevora (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia incisaevora Scoble, 1983: 35
Fomoria indicaevora (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia indicaevora Scoble, 1983: 33
Fomoria leptodictyae (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia leptodictyae Scoble, 1983: 35
Fomoria lucidae (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia lucidae Scoble, 1983: 34
Fomoria malelanensis (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia malelanensis Scoble, 1983: 36
Fomoria myrtinaecola (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia myrtinaecola Scoble, 1983: 34
Fomoria oleivora (Vári, 1955) Diškus & Puplesis, 2003: 388 AFR
Stigmella oleivora Vári, 1955: 336
Ectoedemia oleivora (Vári, 1955) Scoble, 1983: 32

Fomoria pappeivora (Vári, 1963) Diškus & Puplesis, 2003: 388 AFR
Stigmella pappeivora Vári, 1963: 68
Ectoedemia pappeivora (Vári, 1963) Scoble, 1983: 32
Fomoria portensis (Scoble, 1983) Diškus & Puplesis, 2003: 390 AFR
Ectoedemia portensis Scoble, 1983: 36
Fomoria primaria (Meyrick, 1913) Diškus & Puplesis, 2003: 388 AFR
Nepticula primaria Meyrick, 1913: 326
Ectoedemia primaria (Meyrick, 1913) Scoble, 1983: 38
Fomoria scobleella (Minet, 2004), **comb. n.** [56] AFR
Ectoedemia scoblei Minet, 1990: 220; JPH of *Ectoedemia scoblei* Puplesis, 1984a
Ectoedemia scobleella Minet, 2004: 366; RN for *E. scoblei* Minet, 1990
Fomoria scoblei (Minet, 1990) Diškus & Puplesis, 2003: 391
Fomoria tecomariae (Vári, 1955) Diškus & Puplesis, 2003: 388 AFR
Stigmella tecomariae Vári, 1955: 333
Ectoedemia tecomariae (Vári, 1955) Scoble, 1983: 34

other unplaced *Fomoria*

Fomoria viridissimella (Caradja, 1920) Diškus & Puplesis, 2003: 385 WP
Nepticula viridissimella Caradja, 1920: 162
Nepticula nowakowskii Toll, 1957: 199 (syn: van Nieukerken, 1987a: 142)
Ectoedemia viridissimella (Caradja, 1920) van Nieukerken, 1987a: 142
Ectoedemia nowakowskii (Toll, 1957) van Nieukerken, 1986a: 17
Fomoria argyraspis (Puplesis & Diškus, 1995), **comb. n.** [57] EP
Acalyptris argyraspis Puplesis & Diškus, 1995: 51
Fomoria flavimacula Puplesis & Diškus, 1996c: 183 EP
Ectoedemia flavimacula (Puplesis & Diškus, 1996c) online comb.
Fomoria sporadopa (Meyrick, 1911), **comb. n.** [58] OR
Nepticula sporadopa Meyrick, 1911c: 108
Acalyptris sporadopa (Meyrick, 1911) Diškus & Puplesis, 2003: 393
Fomoria tabulosa Puplesis & Diškus in Puplesis et al., 2002: 27 NEO

Muhabbetana Koçak & Kemal, 2007: 5 **stat. n.**; RN for *Laqueus* Scoble, 1983
Laqueus Scoble, 1983: 12 (key), 20; JH of *Laqueus* Dall, 1870 (Brachiopoda) (TS/OD: *Nepticula grandinosa* Meyrick, 1911) [as subgenus of *Ectoedemia*]

***Muhabbetana grandinosa* group** (new)

Muhabbetana furcella (Scoble, 1983), **comb. n.** AFR
Ectoedemia furcella Scoble, 1983: 24
Fomoria furcella (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana grandinosa (Meyrick, 1911), **comb. n.** AFR
Nepticula grandinosa Meyrick, 1911a: 236
Ectoedemia grandinosa (Meyrick, 1911) Scoble, 1983: 21
Fomoria grandinosa (Meyrick, 1911) Diškus & Puplesis, 2003: 386

Muhabbetana guerkiae (Scoble, 1983), **comb. n.** AFR
Ectoedemia guerkiae Scoble, 1983: 22
Fomoria guerkiae (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana jupiteri (Scoble, 1983), **comb. n.** AFR
Ectoedemia jupiteri Scoble, 1983: 25
Fomoria jupiteri (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana macrochaeta (Meyrick, 1921), **comb. n.** AFR
Nepticula macrochaeta Meyrick, 1921b: 140
Ectoedemia macrochaeta (Meyrick, 1921) Scoble, 1983: 23
Fomoria macrochaeta (Meyrick, 1921) Diškus & Puplesis, 2003: 387
Muhabbetana maritima (Scoble, 1983), **comb. n.** AFR
Ectoedemia maritima Scoble, 1983: 24
Fomoria maritima (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana scabridae (Scoble, 1983), **comb. n.** AFR
Ectoedemia scabridae Scoble, 1983: 24
Fomoria scabridae (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana simiicola (Scoble, 1983), **comb. n.** AFR
Ectoedemia simiicola Scoble, 1983: 22
Fomoria simiicola (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana stimulata (Meyrick, 1913), **comb. n.** AFR
Nepticula stimulata Meyrick, 1913: 326
Ectoedemia stimulata (Meyrick, 1913) Scoble, 1983: 21
Fomoria stimulata (Meyrick, 1913) Diškus & Puplesis, 2003: 386
Muhabbetana umdoniella (Scoble, 1983), **comb. n.** AFR
Ectoedemia umdoniella Scoble, 1983: 24
Fomoria umdoniella (Scoble, 1983) Diškus & Puplesis, 2003: 387
Muhabbetana wilkinsoni (Scoble, 1983), **comb. n.** AFR
Ectoedemia wilkinsoni Scoble, 1983: 21
Fomoria wilkinsoni (Scoble, 1983) Diškus & Puplesis, 2003: 387

***Muhabbetana euphorbiella* group** (new)
Muhabbetana euphorbiella (Stainton, 1869), **comb. n.** WP
Nepticula euphorbiella Stainton, 1869b: 229
Nepticula tergestina Klimesch, 1940a: 79 (syn: Laštůvka & Laštůvka, 1997: 167)
Stigmella euphorbiella (Stainton, 1869) Gerasimov, 1952: 238
Ectoedemia euphorbiella (Stainton, 1869) van Nieukerken, 1986a: 17
Fomoria euphorbiella (Stainton, 1869) Diškus & Puplesis, 2003: 384
Stigmella tergestina (Klimesch, 1940) Hering, 1957: 434
Ectoedemia tergestina (Klimesch, 1940) van Nieukerken, 1986a: 17
Muhabbetana jubae (Walsingham, 1908), **comb. n.** WP
Stigmella jubae Walsingham, 1908a: 1011 [2]
Nepticula jubae (Walsingham, 1908) Rebel, 1910: 364

Trifurcula jubae (Walsingham, 1908) Klimesch, 1977: 197
Ectoedemia jubae (Walsingham, 1908) van Nieukerken, 1986b: 17
Fomoria jubae (Walsingham, 1908) Diškus & Puplesis, 2003: 384

Muhabbetana nigrifasciata (Walsingham, 1908), **comb. n.** WP
Stigmella nigrifasciata Walsingham, 1908a: 1011 [2]
Nepticula nigrifasciata (Walsingham, 1908) Rebel, 1910: 364
Dechtiria nigrifasciata (Walsingham, 1908) Klimesch, 1972: 1
Trifurcula nigrifasciata (Walsingham, 1908) Klimesch, 1977: 200
Fomoria nigrifasciata (Walsingham, 1908) Diškus & Puplesis, 2003: 387
Ectoedemia nigrifasciata (Walsingham, 1908) van Nieukerken, 1986b: 17

Muhabbetana vincamajorella (Hartig, 1964), **comb. n.** WP
Nepticula vincamajorella Hartig, 1964: 8
Fomoria vincamajorella (Hartig, 1964) Diškus & Puplesis, 2003: 389
Ectoedemia vincamajorella (Hartig, 1964) van Nieukerken, 1986a: 17

***Muhabbetana* – unplaced species**

Muhabbetana bicarina (Scoble, 1983), **comb. n.** AFR
Ectoedemia bicarina Scoble, 1983: 27
Fomoria bicarina (Scoble, 1983) Diškus & Puplesis, 2003: 389

Muhabbetana capensis (Scoble, 1983), **comb. n.** AFR
Ectoedemia capensis Scoble, 1983: 28
Fomoria capensis (Scoble, 1983) Diškus & Puplesis, 2003: 389

Muhabbetana craspedota (Vári, 1963), **comb. n.** AFR
Stigmella craspedota Vári, 1963: 73
Ectoedemia craspedota (Vári, 1963) Scoble, 1983: 27
Fomoria craspedota (Vári, 1963) Diškus & Puplesis, 2003: 388

Muhabbetana crispae (Scoble, 1983), **comb. n.** AFR
Ectoedemia crispae Scoble, 1983: 31
Fomoria crispae (Scoble, 1983) Diškus & Puplesis, 2003: 390

Muhabbetana denticulata (Scoble, 1983), **comb. n.** AFR
Ectoedemia denticulata Scoble, 1983: 26
Fomoria denticulata (Scoble, 1983) Diškus & Puplesis, 2003: 389

Muhabbetana digitata (Scoble, 1983), **comb. n.** AFR
Ectoedemia digitata Scoble, 1983: 27
Fomoria digitata (Scoble, 1983) Diškus & Puplesis, 2003: 389

Muhabbetana gymnosporiae (Vári, 1955), **comb. n.** AFR
Stigmella gymnosporiae Vári, 1955: 334
Ectoedemia gymnosporiae (Vári, 1955) Scoble, 1983: 29
Fomoria gymnosporiae (Vári, 1955) Diškus & Puplesis, 2003: 388

Muhabbetana insulata (Meyrick, 1911), **comb. n.** AFR
Nepticula insulata Meyrick, 1911b: 79
Ectoedemia insulata (Meyrick, 1911) Scoble, 1983: 29
Fomoria insulata (Meyrick, 1911) Diškus & Puplesis, 2003: 388

Muhabbetana kowynensis (Scoble, 1983), **comb. n.** AFR
Ectoedemia kowynensis Scoble, 1983: 30
Fomoria kowynensis (Scoble, 1983) Diškus & Puplesis, 2003: 389
Muhabbetana limburgensis (Scoble, 1983), **comb. n.** AFR
Ectoedemia limburgensis Scoble, 1983: 28
Fomoria limburgensis (Scoble, 1983) Diškus & Puplesis, 2003: 389
Muhabbetana nigrisquama (Scoble, 1983), **comb. n.** AFR
Ectoedemia nigrisquama Scoble, 1983: 26
Fomoria nigrisquama (Scoble, 1983) Diškus & Puplesis, 2003: 389
Muhabbetana nylstroomensis (Scoble, 1983), **comb. n.** AFR
Ectoedemia nylstroomensis Scoble, 1983: 30
Fomoria nylstroomensis (Scoble, 1983) Diškus & Puplesis, 2003: 389
Muhabbetana psarodes (Vári, 1963), **comb. n.** AFR
Stigmella psarodes Vári, 1963: 70
Ectoedemia psarodes (Vári, 1963) Scoble, 1983: 29
Fomoria psarodes (Vári, 1963) Diškus & Puplesis, 2003: 388
Muhabbetana rhabdophora (Scoble, 1983), **comb. n.** AFR
Ectoedemia rhabdophora Scoble, 1983: 31
Fomoria rhabdophora (Scoble, 1983) Diškus & Puplesis, 2003: 390
Muhabbetana royenicola (Vári, 1955), **comb. n.** AFR
Stigmella royenicola Vári, 1955: 335
Ectoedemia royenicola (Vári, 1955) Scoble, 1983: 25
Fomoria royenicola (Vári, 1955) Diškus & Puplesis, 2003: 388
Muhabbetana subnitescens (Meyrick, 1937), **comb. n.** AFR
Trifurcula subnitescens Meyrick, 1937: 90
Ectoedemia subnitescens (Meyrick, 1937) Scoble, 1983: 28
Fomoria subnitescens (Meyrick, 1937) Diškus & Puplesis, 2003: 387
Muhabbetana undatae (Scoble, 1983), **comb. n.** AFR
Ectoedemia undatae Scoble, 1983: 27
Fomoria undatae (Scoble, 1983) Diškus & Puplesis, 2003: 389

Parafomoria Borkowski, 1975: 498 (TS/OD: *Nepticula helianthemella* Herrich-Schäffer, 1860: 60)
Parafomoria van Nieukerken, 1983b: 454 JH of *Parafomoria* Borkowski, 1975

***Parafomoria liguricella* group** (new)
Parafomoria ladaniphila (Mendes, 1910) van Nieukerken, 1983b: 468 WP
Nepticula ladaniphila Mendes, 1910a: 102
Stigmella ladaniphila (Mendes, 1910) Klimesch, 1948a: 170
Ectoedemia ladaniphila (Mendes, 1910) Gómez Bustillo, 1981: 19
Parafomoria liguricella (Klimesch, 1948) van Nieukerken, 1983b: 466[29] WP
Stigmella liguricella Klimesch, 1948a: 170[29]

Parafomoria tingitella (Walsingham, 1904) van Nieukerken, 1983b: 469 WP
Nepticula tingitella Walsingham, 1904: 8
Stigmella tingitella (Walsingham, 1904) Gerasimov, 1952: 264

***Parafomoria helianthemella* group** (new)
Parafomoria cistivora (Peyerimhoff, 1871) van Nieukerken, 1983b: 458 WP
Nepticula cistivora Peyerimhoff, 1871: 414
Stigmella cistivora (Peyerimhoff, 1871) Suire, 1951: 71
Parafomoria fumanae A. Laštuvka & Z. Laštuvka, 2005b: 15 WP
Parafomoria halimivora van Nieukerken, 1985a: 24 WP
Parafomoria helianthemella (Herrich-Schäffer, 1860) Borkowski, 1975: 498 (Fig. 34) WP
Nepticula helianthemella Herrich-Schäffer, 1860: 60
Stigmella helianthemella (Herrich-Schäffer, 1860) Klimesch, 1948a: 171
Trifurcula helianthemella (Herrich-Schäffer, 1860) Leraut, 1980: 49
Parafomoria pseudocistivora van Nieukerken, 1983b: 460 WP

Etainia Beirne, 1945: 208 (TS/OD: *Lyonetia sericopeza* Zeller, 1839)
Obrussa Braun, 1915: 196 JH of *Obrussa* Heyden, 1891 (Lepidoptera: Geometridae) (TS/M: *Nepticula ochrefasciella* Chambers, 1873) (syn: van Nieukerken, 1986a: 16)
Etainia albibimaculella (Larsen, 1927) Puplesis & Diškus, 1996a: 5 WP,NEA
Nepticula albibimaculella Larsen, 1927: 5
Stigmella albibimaculella (Larsen, 1927) Hering, 1957: 112
Trifurcula albibimaculella (Larsen, 1927) Johansson, 1971: 246
Ectoedemia albibimaculella (Larsen, 1927) van Nieukerken, 1986a: 16
Etainia biarmata Puplesis, 1994: 233 WP
Ectoedemia biarmata (Puplesis, 1994) van Nieukerken & Laštůvka, 2002: 89
Etainia decentella (Herrich-Schäffer, 1855) Beirne, 1945: 207 WP
Nepticula decentella Herrich-Schäffer, 1855a: 358
Nepticula monspessulanella Jäckh, 1951: 171 (syn: van Nieukerken, 1986a: 16)
Stigmella decentella (Herrich-Schäffer, 1855) Gerasimov, 1952: 234
Trifurcula decentella (Herrich-Schäffer, 1855) Johansson, 1971: 246
Ectoedemia decentella (Herrich-Schäffer, 1855) van Nieukerken, 1986a: 16
Stigmella monspessulanella (Jäckh, 1951) Hering, 1957: 19
Etainia leptognathos Puplesis & Diškus, 1996a: 44 WP
Ectoedemia leptognathos (Puplesis & Diškus, 1996) van Nieukerken & Laštůvka, 2002: 89
Etainia louisella (Sircom, 1849) Bradley et al., 1972: 3 WP
Nepticula louisella Sircom, 1849: XIII
Nepticula sphendamni Hering, 1937: 561 (syn: van Nieukerken, 1986a: 16)
Ectoedemia louisella (Sircom, 1849) van Nieukerken, 1986a: 16
Stigmella sphendamni (Hering, 1937) Klimesch, 1951b: 64

Trifurcula sphendamni (Hering, 1937) Johansson, 1971: 246
Etainia sphendamni (Hering, 1937) Bradley et al., 1972: 3
Etainia obtusa Puplesis & Diškus, 1996a: 46 WP
Ectoedemia obtusa (Puplesis & Diškus, 1996) Krenek, 2000: 36
Etainia sericopeza (Zeller, 1839) Beirne, 1945: 207 (Fig. 35) WP,[NEA]
Lyonetia sericopeza Zeller, 1839: 215
Oecophora sericopezella Duponchel, 1843: 344 UE
Tinea maryella Duponchel, 1843: 464 (syn: Frey, 1857: 402)
Nepticula acerella Goureau, 1860: xxiii (syn: Joannis, 1915: 131)
Nepticula sericopeza (Zeller, 1839) Heyden, 1843: 209
Stigmella sericopeza (Zeller, 1839) Walsingham, 1916: 160
Trifurcula sericopeza (Zeller, 1839) Johansson, 1971: 246
Obrussa sericopeza (Zeller, 1839) Wilkinson & Scoble, 1979: 101
Ectoedemia sericopeza (Zeller, 1839) van Nieukerken, 1986b: 16
Lyonetia sericopezella (Duponchel, 1843) Duponchel, 1844: 378
‡ *Stigmella sericopeza* f. *palliolella* Le Marchand, 1944: 358
Etainia capesella (Puplesis in Puplesis & Ivinskis, 1985) Puplesis, 1994: 232 EP
Obrussa capesella Puplesis in Puplesis & Ivinskis, 1985: 39
Ectoedemia capesella (Puplesis in Puplesis & Ivinskis, 1985) Hirano, 2013: 93
Etainia peterseni (Puplesis in Puplesis & Ivinskis, 1985) Puplesis, 1994: 231 EP
Obrussa peterseni Puplesis in Puplesis & Ivinskis, 1985: 41
Ectoedemia peterseni (Puplesis in Puplesis & Ivinskis, 1985) Hirano, 2013: 92
Etainia sabina (Puplesis in Puplesis & Ivinskis, 1985) Puplesis, 1994: 231 EP
Obrussa sabina Puplesis in Puplesis & Ivinskis, 1985: 43
Ectoedemia sabina (Puplesis in Puplesis & Ivinskis, 1985) online comb.
Etainia trifasciata (Matsumura, 1931) Diškus & Puplesis, 2003: 408 [59] EP
Nepticula trifasciata Matsumura, 1931: 1114
Obrussa tigrinella Puplesis in Puplesis & Ivinskis, 1985: 40 **syn. n.** [59]
Stigmella trifasciata (Matsumura, 1931) Kuroko, 1982a: 448
Ectoedemia trifasciata (Matsumura, 1931) Hirano, 2013: 93
Ectoedemia tigrinella (Puplesis in Puplesis & Ivinskis, 1985) Hirano, 2013: 92
Etainia tigrinella (Puplesis in Puplesis & Ivinskis, 1985) Puplesis, 1994: 232
Etainia crypsixantha (Meyrick, 1918) Vári & Kroon, 1986: 153 AFR
Nepticula crypsixantha Meyrick, 1918a: 43
Obrussa crypsixantha (Meyrick, 1918) Scoble, 1983: 17
Ectoedemia crypsixantha (Meyrick, 1918) online comb.
Etainia krugerensis (Scoble, 1983) Vári & Kroon, 1986: 153 AFR
Obrussa krugerensis Scoble, 1983: 19
Ectoedemia krugerensis (Scoble, 1983) online comb.
Etainia nigricapitella (Janse, 1948) Vári & Kroon, 1986: 153 AFR
Nepticula nigricapitella Janse, 1948: 170
Obrussa nigricapitella (Janse, 1948) Scoble, 1983: 18
Ectoedemia nigricapitella (Janse, 1948) online comb.

Etainia zimbabwiensis (Scoble, 1983) Vári & Kroon, 1986: 153 AFR
Obrussa zimbabwiensis Scoble, 1983: 18
Ectoedemia zimbabwiensis (Scoble, 1983) online comb.
Etainia ochrefasciella (Chambers, 1873) Puplesis & Diškus, 1996a: 4 NEA
Nepticula ochrefasciella Chambers, 1873: 128
Obrussa ochrefasciella (Chambers, 1873) Braun, 1915: 196
Ectoedemia ochrefasciella (Chambers, 1873) van Nieukerken, 1986a: 19

Acalyptris Meyrick, 1921a: 410 (TS/OD,M: *Acalyptris psammophricta* Meyrick, 1921: 410)
Microcalyptris Braun, 1925b: 224 (TS/OD,M: *Microcalyptris scirpi* Braun, 1925: 225) (syn: van Nieukerken, 1986a: 14)
Weberia Müller-Rutz, 1934a: 122 JH of *Weberia* Robineau-Desvoidy, 1830 (Diptera: Tachinidae) (TS/OD,M: *Weberia platani* Müller-Rutz, 1934: 122) (syn: van Nieukerken, 1986a: 14)
Niepeltia Strand, 1934: 241; RN for *Weberia* Müller-Rutz, 1934 (syn: van Nieukerken, 1986a: 14)
Weberina Müller-Rutz, 1934b: Errata, 148; RN for *Weberia* Müller-Rutz, 1934 (syn: van Nieukerken, 1986a: 14)

***Acalyptris scirpi* group** (new)
Acalyptris bicornutus (Davis, 1978) Puplesis & Robinson, 2000: 53 NEA
Microcalyptris bicornutus Davis, 1978: 212
Acalyptris bipinnatellus (Wilkinson, 1979) van Nieukerken, 1986a: 16 NEA
Microcalyptris bipinnatellus Wilkinson, 1979: 75
Acalyptris lotella (Wagner, 1987) Diškus & Puplesis, 2003: 397 NEA
Microcalyptris lotella Wagner, 1987: 278
Acalyptris punctulata (Braun, 1910) Diškus & Puplesis, 2003: 393 NEA
Nepticula punctulata Braun, 1910: 174
Microcalyptris punctulata (Braun, 1910) Wilkinson, 1979: 71
Acalyptris scirpi (Braun, 1925) Diškus & Puplesis, 2003: 393 NEA
Microcalyptris scirpi Braun, 1925b: 225
Acalyptris thoracealbella (Chambers, 1873) Diškus & Puplesis, 2003: 393 NEA
Microcalyptris thoracealbella (Chambers, 1873) Davis, 1978: 214
Nepticula thoracealbella Chambers, 1873: 127
Nepticula badiocapitella Chambers, 1876: 160 (syn: Braun, 1917: 189)
Acalyptris tenuijuxtus (Davis, 1978) Puplesis & Robinson, 2000: 51 NEA,NEO
Microcalyptris tenuijuxtus Davis, 1978: 216
Acalyptris basicornis Remeikis & Stonis in Stonis et al., 2013d: 102 NEO
Acalyptris basihastatus Puplesis & Diškus in Puplesis et al., 2002: 29 NEO
Acalyptris bifidus Puplesis & Robinson, 2000: 50 NEO
Acalyptris bovicorneus Puplesis & Robinson, 2000: 45 NEO
Acalyptris caribbicus Diškus & Stonis in Stonis et al., 2013d: 106 NEO
Acalyptris dominicanus Remeikis & Stonis in Stonis & Remeikis, 2015: 85 NEO

Acalyptris fortis Puplesis & Robinson, 2000: 47 NEO
Acalyptris hispidus Puplesis & Robinson, 2000: 48 NEO
Acalyptris janzeni van Nieukerken & Nishida in van Nieukerken et al., 2016: 55 NEO
Acalyptris lascuevella Puplesis & Robinson, 2000: 49 NEO
Acalyptris laxibasis Puplesis & Robinson, 2000: 52 NEO
Acalyptris martinheringi Puplesis & Robinson, 2000: 46 NEO
Acalyptris nigrisignum Remeikis & Stonis in Stonis & Remeikis, 2015: 79 NEO
Acalyptris novenarius Puplesis & Robinson, 2000: 48 NEO
Acalyptris paradividua Šimkevičiūtė & Stonis in Šimkevičiūtė et al., 2009: 272 NEO
Acalyptris peteni Diškus & Stonis in Stonis et al., 2013d: 102 NEO
Acalyptris pseudohastatus Puplesis & Diškus in Puplesis et al., 2002: 30 NEO
Acalyptris statuarius Diškus & Stonis in Stonis et al., 2013d: 109 NEO
Acalyptris terrificus Šimkevičiūtė & Stonis in Šimkevičiūtė et al., 2009: 275 NEO
Acalyptris trifidus Puplesis & Robinson, 2000: 50 NEO
Acalyptris trigonijuxtus Remeikis & Stonis in Stonis & Remeikis, 2015: 83 NEO
Acalyptris unicornis Puplesis & Robinson, 2000: 51 NEO

***Acalyptris staticis* group** (van Nieukerken, 2007a: 17)
Acalyptris lesbia van Nieukerken & Hull in van Nieukerken, 2007a: 22 WP
Acalyptris limoniastri van Nieukerken, 2007a: 23 WP
Acalyptris limonii Z. Laštuvka & A. Laštuvka, 1998: 314 WP
Acalyptris maritima A. Laštuvka & Z. Laštuvka, 1997: 119 WP
Acalyptris pyrenaica A. Laštuvka & Z. Laštuvka, 1993: 158 WP
Acalyptris staticis (Walsingham, 1908) van Nieukerken, 1986b: 14 WP
Stigmella staticis Walsingham, 1908a: 1009 [1]
Nepticula staticis (Walsingham, 1908) Rebel, 1910: 373

***Acalyptris psammophricta* group** (new)
Acalyptris repeteki group (Puplesis, 1988: 509)
Acalyptris falkovitshi (Puplesis, 1984) van Nieukerken, 1986a: 14 [60] WP,EP
Microcalyptris falkovitshi Puplesis, 1984c: 499
Microcalyptris turanicus Puplesis, 1984c: 497 (syn: van Nieukerken, 2010: 501)
Microcalyptris vittatus Puplesis, 1984c: 491 **syn. n.** [60]
Microcalyptris arenosus Falkovitsh, 1986: 168 **syn. n.** [60]
Acalyptris turanicus (Puplesis, 1984) van Nieukerken, 1986a: 14
Acalyptris vittatus (Puplesis, 1984) van Nieukerken, 1986a: 14
Acalyptris arenosus (Falkovitsh, 1986) Puplesis, 1990: 66
Acalyptris galinae (Puplesis, 1984) van Nieukerken, 1986a: 14 WP,EP
Microcalyptris galinae Puplesis, 1984c: 502
Microcalyptris galinae mesasiaticus Puplesis, 1984c: 503
Acalyptris galinae mesasiaticus (Puplesis, 1984) Puplesis, 1990: 84

Acalyptris pallens (Puplesis, 1984) van Nieukerken, 1986a: 14 WP,EP
Microcalyptris pallens Puplesis, 1984c: 501
Acalyptris psammophricta Meyrick, 1921a: 410 WP,EP,OR
Microcalyptris lvovskyi Puplesis, 1984c: 494 (syn: van Nieukerken, 2010: 501)
Acalyptris lvovskyi (Puplesis, 1984) van Nieukerken, 1986a: 14
Acalyptris repeteki (Puplesis, 1984) van Nieukerken, 1986a: 14 WP
Microcalyptris repeteki Puplesis, 1984c: 494
Acalyptris turcomanicus (Puplesis, 1984) van Nieukerken, 1986a: 14 WP
Microcalyptris turcomanicus Puplesis, 1984c: 499

***Acalyptris shafirkanus* group** (Puplesis, 1988: 506)
Acalyptris brevis Puplesis, 1990: 86 WP
Acalyptris desertellus (Puplesis, 1984) van Nieukerken, 1986a: 14 WP
Microcalyptris desertellus Puplesis, 1984c: 493
Acalyptris egidijui Puplesis, 1990: 87 WP
Acalyptris kizilkumi (Falkovitsh, 1986) Puplesis, 1990: 86 WP,EP
Microcalyptris kizilkumi Falkovitsh, 1986: 167
Acalyptris piculus Puplesis, 1990: 85 EP
Acalyptris shafirkanus (Puplesis, 1984) van Nieukerken, 1986a: 14 WP
Microcalyptris shafirkanus Puplesis, 1984c: 493
Acalyptris vannieukerkeni Puplesis, 1994: 218 WP

***Acalyptris platani* group** (van Nieukerken, 2007a: 7)
Acalyptris gielisi van Nieukerken, 2010: 500 WP
Acalyptris loranthella (Klimesch, 1937) van Nieukerken, 1986a: 14 WP
Nepticula loranthella Klimesch, 1937: 33
Stigmella loranthella (Klimesch, 1937) Klimesch, 1948b: 78
Weberina loranthella (Klimesch, 1937) Szőcs, 1978: 268
Niepeltia loranthella (Klimesch, 1937) van Achterberg, 1983: 30
Acalyptris minimella (Rebel, 1926) van Nieukerken, 1986a: 14 WP
Trifurcula minimella Rebel, 1926: (110)
Weberina lentiscella Groschke, 1944: 117 (syn: Klimesch, 1978a: 256)
Nepticula minimella (Rebel, 1926); Klimesch, 1953a: 162 JSH of *Nepticula minimella* Chambers, 1873
Niepeltia lentiscella (Groschke, 1944) Hering, 1957: 781
Niepeltia minimella (Rebel, 1926) Scoble, 1980a: 207
Acalyptris pistaciae van Nieukerken, 2007a: 14 WP
Acalyptris platani (Müller-Rutz, 1934) van Nieukerken, 1986a: 14 (Fig. 36) WP
Weberia platani Müller-Rutz, 1934a: 122
Niepeltia platani (Müller-Rutz, 1934) Strand, 1934: 241
Weberina platani (Müller-Rutz, 1934) Müller-Rutz, 1934b: slip
Trifurcula platani (Müller-Rutz, 1934) Klimesch, 1978a: 253

Acalyptris acontarcha (Meyrick, 1926) Diškus & Puplesis, 2003: 393 OR
Nepticula acontarcha Meyrick, 1926: 295
Stigmella acontarcha (Meyrick, 1926) Fletcher, 1933: 82
Acalyptris auratilis Puplesis & Diškus, 2003a: 219 OR
Acalyptris clinomochla (Meyrick, 1934) Diškus & Puplesis, 2003: 394 OR
Nepticula clinomochla Meyrick, 1934a: 468
Trifurcula clinomochla (Meyrick, 1934) Gustafsson, 1976: 49
Niepeltia clinomochla (Meyrick, 1934) Scoble, 1980a: 216
Acalyptris heteranthes (Meyrick, 1926) Diškus & Puplesis, 2003: 393 OR
Nepticula heteranthes Meyrick, 1926: 296
Acalyptris melanospila (Meyrick, 1934) Puplesis & Diškus, 2003a: 218 OR
Nepticula melanospila Meyrick, 1934a: 468
Acalyptris nigripexus Puplesis & Diškus, 2003: 220 OR
Acalyptris acumenta (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia acumenta Scoble, 1980b: 213
Acalyptris bispinata (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia bispinata Scoble, 1980b: 213
Acalyptris combretella (Vári, 1955) Vári et al., 2002: 8 AFR
Stigmella combretella Vári, 1955: 332
Niepeltia combretella (Vári, 1955) Scoble, 1980a: 206
Acalyptris fagarivora (Vári, 1955) Vári et al., 2002: 8 AFR
Stigmella fagarivora Vári, 1955: 334
Niepeltia fagarivora (Vári, 1955) Scoble, 1980a: 209
Acalyptris fulva (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia fulva Scoble, 1980b: 214
Acalyptris fuscofascia (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia fuscofascia Scoble, 1980b: 210
Acalyptris krooni (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia krooni Scoble, 1980b: 212
Acalyptris krugeri (Vári, 1963) Diškus & Puplesis, 2003: 394 AFR
Stigmella krugeri Vári, 1963: 71
Acalyptris lanneivora (Vári, 1955) Vári et al., 2002: 8 AFR
Stigmella lanneivora Vári, 1955: 332
Niepeltia lanneivora (Vári, 1955) Scoble, 1980a: 215
Acalyptris lorantivora (Janse, 1948) Vári et al., 2002: 8 AFR
Nepticula lorantivora Janse, 1948: 169
Niepeltia lorantivora (Janse, 1948) Scoble, 1980a: 211
Acalyptris lundiensis (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia lundiensis Scoble, 1980b: 214
Acalyptris mariepsensis (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia mariepsensis Scoble, 1980b: 214
Acalyptris molleivora (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia molleivora Scoble, 1980b: 207

Acalyptris obliquella (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia obliquella Scoble, 1980b: 209
Acalyptris pundaensis (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia pundaensis Scoble, 1980b: 211
Acalyptris rubiaevora (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia rubiaevora Scoble, 1980b: 208
Acalyptris sellata (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia sellata Scoble, 1980b: 213
Acalyptris umdoniensis (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia umdoniensis Scoble, 1980b: 210
Acalyptris vacuolata (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia vacuolata Scoble, 1980b: 215
Acalyptris vepricola (Vári, 1963) Vári et al., 2002: 8 AFR
Stigmella vepricola Vári, 1963: 68
Niepeltia vepricola (Vári, 1963) Scoble, 1983: 44
Acalyptris vumbaensis (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia vumbaensis Scoble, 1980b: 207
Acalyptris zeyheriae (Scoble, 1980) Vári et al., 2002: 8 AFR
Niepeltia zeyheriae Scoble, 1980b: 208

***Acalyptris latipennata* group** (Puplesis et al., 2002: 66)
Acalyptris dividua Puplesis & Robinson, 2000: 54 NEO
Acalyptris ecuadoriana Puplesis & Diškus in Puplesis et al., 2002: 27 NEO
Acalyptris latipennata (Puplesis & Robinson, 2000) Puplesis et al., 2002: 66 NEO
Fomoria latipennata Puplesis & Robinson, 2000: 45
Acalyptris onorei Puplesis & Diškus in Puplesis et al., 2002: 28 NEO

unplaced *Acalyptris* [61]
Acalyptris distaleus (Wilkinson, 1979) Diškus & Puplesis, 2003: 395 [61] NEA
Microcalyptris distaleus Wilkinson, 1979: 78
Acalyptris postalatratus (Wilkinson, 1979) Diškus & Puplesis, 2003: 395 NEA
Microcalyptris postalatratus Wilkinson, 1979: 77
Acalyptris amazonius Puplesis & Diškus in Puplesis et al., 2002: 32 NEO
Acalyptris articulosus Puplesis & Diškus in Puplesis et al., 2002: 30 NEO
Acalyptris insolentis Puplesis & Diškus in Puplesis et al., 2002: 33 NEO
Acalyptris platygnathos Puplesis & Robinson, 2000: 54 NEO
Acalyptris rotundus Puplesis & Diškus in Puplesis et al., 2002: 31 NEO
Acalyptris yucatani Remeikis & Stonis in Stonis et al., 2013b: 227 NEO

Zimmermannia Hering, 1940: 266 (TS/OD,M: *Ectoedemia liebwerdella* Zimmermann, 1940) [62]
Ectoedemia castaneae group (Wilkinson & Newton, 1981: 72)
Zimmermannia amani (Svensson, 1966), **comb. n.** WP,EP
Ectoedemia amani Svensson, 1966: 200

Ectoedemia emendata Puplesis, 1985c: 69 (syn: Puplesis, 1994: 15)
Trifurcula amani (Svensson, 1966) Johansson, 1971: 245

Zimmermannia atrifrontella (Stainton, 1851), **comb. n.** (Fig. 37) WP
Trifurcula atrifrontella Stainton, 1851: 11
Zimmermannia heringiella Doets, 1947: 504 (syn: Klimesch, 1953a: 191)
Ectoedemia atrifrontella (Stainton, 1851) Klimesch, 1953b: 191
Ectoedemia heringiella (Doets, 1947) Klimesch, 1953b: 191

Zimmermannia hispanica (van Nieukerken, 1985), **comb. n.** WP
Ectoedemia hispanica van Nieukerken, 1985b: 22

Zimmermannia liebwerdella (Zimmermann, 1940) Hering, 1940: 266 WP
Ectoedemia liebwerdella Zimmermann, 1940: 264

Zimmermannia liguricella (Klimesch, 1953), **comb. n.** WP
Ectoedemia liguricella Klimesch, 1953b: 194

Zimmermannia longicaudella (Klimesch, 1953), **comb. n.** WP
Ectoedemia longicaudella Klimesch, 1953b: 193
Stigmella peiuii Nemeş, 1972: 153 (syn: van Nieukerken, 1985b: 21)
Trifurcula longicaudella (Klimesch, 1953) Johansson, 1971: 245

Zimmermannia monemvasiae (van Nieukerken, 1985), **comb. n.** WP
Ectoedemia monemvasiae van Nieukerken, 1985b: 23

Zimmermannia reichli (Z. Laštuvka & A. Laštuvka, 1998), **comb. n.** WP
Ectoedemia reichli Z. Laštuvka & A. Laštuvka, 1998: 316

Zimmermannia vivesi (A. Laštuvka, Z. Laštuvka & van Nieukerken, 2010), **comb. n.** WP
Ectoedemia vivesi A. Laštuvka, Z. Laštuvka & van Nieukerken in van Nieukerken et al., 2010: 12

Zimmermannia admiranda (Puplesis, 1984), **comb. n.** EP
Ectoedemia admiranda Puplesis, 1984b: 588

Zimmermannia nuristanica (van Nieukerken, 1985), **comb. n.** EP
Ectoedemia nuristanica van Nieukerken, 1985b: 25

Zimmermannia sivickisi (Puplesis, 1984), **comb. n.** EP
Ectoedemia sivickisi Puplesis, 1984b: 590
Ectoedemia laura Puplesis, 1985c: 68 (syn: Rocienė & Stonis, 2013: 108)

Zimmermannia bosquella (Chambers, 1878), **stat rev., comb. n.** [62] NEA
Nepticula bosquella Chambers, 1878a: 106
Nepticula bosqueella Chambers, 1878b: 157 ISS
Ectoedemia castaneae Busck, 1913: 103 **syn. n.** [62]
Ectoedemia heinrichi Busck, 1914: 149 **syn. n.** [62]
Ectoedemia helenella Wilkinson, 1981: 105 **syn. n.** [62]
Ectoedemia bosquella (Chambers, 1878) Braun, 1917: 200
Opostega bosqueella (Chambers, 1878) Dyar et al., 1903: 547
Ectoedemia obrutella sensu Wilkinson & Newton, 1981: 72 [misapplied]

Zimmermannia grandisella (Chambers, 1880), **comb. n.** [62] NEA
Nepticula grandisella Chambers, 1880a: 193

Ectoedemia chloranthis Meyrick, 1928b: 462 **syn. n.** [62]
Ectoedemia acanthella Wilkinson & Newton, 1981: 75 **syn. n.** [62]
Ectoedemia grandisella (Chambers, 1880) Wilkinson, 1981: 96

Zimmermannia mesoloba (Davis, 1978), **comb. n.** [62] NEA
Ectoedemia mesoloba Davis, 1978: 209
Ectoedemia coruscella Wilkinson, 1981: 99 **syn. n.** [62]

Zimmermannia obrutella (Zeller, 1873), **comb. n.** [62] NEA
Trifurcula obrutella Zeller, 1873: 316
Ectoedemia piperella Wilkinson & Newton, 1981: 77 **syn. n.** [62]
Ectoedemia reneella Wilkinson, 1981: 104 **syn. n.** [62]
Ectoedemia obrutella (Zeller, 1873) Busck, 1913: 103

Zimmermannia phleophaga (Busck, 1914), **comb. n.** NEA
Ectoedemia phleophaga Busck, 1914: 3

Ectoedemia Busck, 1907: 97 (TS/OD,M: *Ectoedemia populella* Busck, 1907: 98)
Dechtiria Beirne, 1945: 204 (TS/OD: *Tinea subbimaculella* Haworth, 1828: 583) (syn: Svensson, 1966: 200)

***Ectoedemia commiphorella* group** (Doorenweerd et al., 2015: 9)

Ectoedemia commiphorella Scoble, 1978a: 82 AFR
Ectoedemia expeditionis Mey, 2004: 30 AFR
Ectoedemia mauni Scoble, 1979: 36 AFR
Ectoedemia nigrimacula (Janse, 1948) Scoble, 1978a: 84 AFR
Nepticula nigrimacula Janse, 1948: 171
Ectoedemia tersiusi Mey, 2004: 31 AFR

***Ectoedemia terebinthivora* group** (van Nieukerken, 1985b: 63)

Ectoedemia terebinthivora (Klimesch, 1975) van Nieukerken, 1985b: 63 WP
Trifurcula terebinthivora Klimesch, 1975c: 19

***Ectoedemia populella* group** (Wilkinson & Newton, 1981: 41)

Ectoedemia intimella (Zeller, 1848) Bradley et al., 1972: 3 WP
Nepticula intimella Zeller, 1848: 323
Stigmella intimella (Zeller, 1848) Fletcher & Clutterbuck, 1945: 61
Dechtiria intimella (Zeller, 1848) Beirne, 1945: 205
Trifurcula intimella (Zeller, 1848) Johansson, 1971: 245
Ectoedemia insularis Puplesis, 1985c: 68 [63] EP
Ectoedemia sinevi Puplesis, 1985c: 67 [64] EP
Ectoedemia populella Busck, 1907: 98 NEA
Ectoedemia hannoverella (Glitz, 1872) Borkowski, 1972a: Fig. 7 WP,EP
Nepticula hannoverella Glitz, 1872: 25
Stigmella hannoverella (Glitz, 1872) Klimesch, 1951b: 64
Trifurcula hannoverella (Glitz, 1872) Johansson, 1971: 245

Ectoedemia canutus Wilkinson & Scoble, 1979: 81 NEA

Ectoedemia turbidella (Zeller, 1848) Bradley et al., 1972: 3 [65] WP

Nepticula argyropeza var. *turbidella* Zeller, 1848: 321

‡ [no genus] *argyropeza* Herrich-Schäffer, 1853: pl106 NN

Nepticula argyropezella Herrich-Schäffer, 1855a: 357 UE

Nepticula populi-albae Hering, 1935: 7 (syn: van Nieukerken, 1985b: 31)

Stigmella marionella Ford, 1950: 39 (syn: Bradley et al., 1972: 3)

Ectoedemia similigena Puplesis, 1994: 180 **syn. n.** [65]

Dechtiria turbidella (Zeller, 1848) Vári, 1950: 182

Stigmella turbidella (Zeller, 1848) Klimesch, 1951b: 64

Trifurcula turbidella (Zeller, 1848) Johansson, 1971: 245

Stigmella populialbae (Hering, 1935) Gerasimov, 1952: 252

Ectoedemia populialbae (Hering, 1935) Borkowski, 1975: 495

Ectoedemia albida Puplesis, 1994: 179 WP

Ectoedemia klimeschi (Skala, 1933) Borkowski, 1975: 495 (Fig. 38) WP

Nepticula klimeschi Skala, 1933a: 31

Stigmella niculescui Nemeş, 1970: 33 (syn: van Nieukerken, 1985b: 34)

Stigmella klimeschi (Skala, 1933) Gerasimov, 1952: 244

Ectoedemia argyropeza (Zeller, 1839) Bradley et al., 1972: 3 WP,EP,[NEA]

Lyonetia argyropeza Zeller, 1839: 215

Lyonetia argyropezella Duponchel, 1844: 378 UE

Nepticula apicella Stainton, 1854: 300 (syn: Heinemann & Wocke, [1876]: 768) [6]

Nepticula argyropezella Doubleday, 1859: 36 UE

Nepticula turbulentella Wocke, 1861: 129 URN

Nepticula simplicella Heinemann, 1862b: 319 (syn: van Nieukerken, 1985b: 35)

Ectoedemia argyropeza downesi Wilkinson & Scoble, 1979: 80

Nepticula argyropeza (Zeller, 1839) Zeller, 1848: 320

Stigmella argyropeza (Zeller, 1839) Fletcher & Clutterbuck, 1945: 61

Dechtiria argyropeza (Zeller, 1839) Emmet, 1971a: 243

Trifurcula argyropeza (Zeller, 1839) Johansson, 1971: 245

‡ *Nepticula argyropeza* ab. *houzeaui* Dufrane, 1942: 11

‡ *Nepticula argyropeza* ab. *morosella* Steudel in Steudel & Hoffmann, 1882: 244

***Ectoedemia subbimaculella* group - satellite taxa** (Doorenweerd et al., 2015)

Ectoedemia preisseckeri group (van Nieukerken, 1985b: 37)

Ectoedemia arisi Puplesis, 1984a: 120 EP

Ectoedemia scoblei Puplesis, 1984a: 122 EP

Ectoedemia christopheri Puplesis, 1985c: 69; RN for *E. wilkinsoni* Puplesis, 1984a EP

Ectoedemia wilkinsoni Puplesis, 1984a: 122; JPH of *Ectoedemia wilkinsoni* Scoble, 1983

Ectoedemia trinotata (Braun, 1914) Wilkinson & Newton, 1981: 46 NEA

Nepticula trinotata Braun, 1914: 18

Ectoedemia philipi Puplesis, 1984b: 590 EP
Ectoedemia preisseckeri (Klimesch, 1941) Borkowski, 1975: 493 WP,EP
Nepticula preisseckeri Klimesch, 1941: 162
Stigmella preisseckeri (Klimesch, 1941) Hering, 1957: 1092
Ectoedemia quadrinotata (Braun, 1917) Wilkinson & Scoble, 1979: 95 NEA
Nepticula quadrinotata Braun, 1917: 168

***Ectoedemia subbimaculella* group** (van Nieukerken, 1985b: 43)
Ectoedemia gilvipennella (Klimesch, 1948) van Nieukerken, 1985b: 45[29] WP
Stigmella gilvipennella Klimesch, 1948a: 168[29]
Nepticula gilvipennella (Klimesch, 1948) Szőcs, 1968: 228
Ectoedemia quinquella (Bedell, 1848) Bradley et al., 1972: 2 WP
Microsetia quinquella Bedell, 1848: 1986
Nepticula quinquella (Bedell, 1848) Stainton, 1849: 29
Dechtiria quinquella (Bedell, 1848) Beirne, 1945: 206
Stigmella quinquella (Bedell, 1848) Gerasimov, 1952: 255
Trifurcula quinquella (Bedell, 1848) Johansson, 1971: 245
Ectoedemia coscoja van Nieukerken, A. Laštuvka & Z. Laštuvka, 2010: 45 WP
Ectoedemia algeriensis van Nieukerken, 1985b: 44 WP
Ectoedemia leucothorax van Nieukerken, 1985b: 46 WP
Ectoedemia haraldi (Soffner, 1942) Klimesch, 1975a: 864 WP
Nepticula haraldi Soffner, 1942: 56
Stigmella prinophyllella Le Marchand, 1946: 285 (syn: Le Marchand, 1948: 298)
Stigmella haraldi (Soffner, 1942) Hering, 1957: 867
Trifurcula haraldi (Soffner, 1942) Leraut, 1980: 49
Ectoedemia pseudoilicis Z. Laštuvka & A. Laštuvka, 1998: 317 WP
Ectoedemia ilicis (Mendes, 1910b) van Nieukerken, 1985b: 48 WP
Nepticula ilicis Mendes, 1910b: 164
Stigmella ilicis (Mendes, 1910b) Gerasimov, 1952: 243
Ectoedemia heringella (Mariani, 1939) van Nieukerken, 1985b: 49 WP
Nepticula heringella Mariani, 1939: 5
Stigmella heringella (Mariani, 1939) Hering, 1957: 868
‡ *Nepticula heringella* f. *alliatae* Mariani, 1939: 7
Ectoedemia alnifoliae van Nieukerken, 1985b: 50 WP
Ectoedemia aligera Puplesis, 1985c: 67 EP
Ectoedemia ermolaevi Puplesis, 1985c: 68 EP
Ectoedemia cerviparadisicola Sato in Shinozaki et al., 2012: 578 EP
Ectoedemia maculata Puplesis, 1987: 11 EP
Ectoedemia rufifrontella (Caradja, 1920) van Nieukerken, 1987a: 142 WP
Trifurcula rufifrontella Caradja, 1920: 161
Nepticula nigrosparsella Klimesch, 1940b: 91 (syn: van Nieukerken, 1987a: 142)
Stigmella nigrosparsella (Klimesch, 1940b) Klimesch, 1948a: 170
Ectoedemia nigrosparsella (Klimesch, 1940b) Kasy, 1983: 5

Ectoedemia albifasciella complex (van Nieukerken, 1985b: 52) (next 4 species)

Ectoedemia pubescivora (Weber, 1937) van Nieukerken, 1985b: 55 WP

Nepticula pubescivora Weber, 1937: 212

Stigmella pubescivora (Weber, 1937) Klimesch, 1948b: 73

Trifurcula pubescivora (Weber, 1937) Kasy, 1979: 4

Ectoedemia albifasciella (Heinemann, 1871) Bradley et al., 1972: 3 WP

Nepticula albifasciella Heinemann, 1871: 222

Nepticula subapicella Stainton, 1886: 238 (syn: Emmet, 1974b: 274)

Dechtiria albifasciella (Heinemann, 1871) Beirne, 1945: 205

Stigmella albifasciella (Heinemann, 1871) Klimesch, 1951b: 66

Trifurcula albifasciella (Heinemann, 1871) Johansson, 1971: 245

Ectoedemia contorta van Nieukerken, 1985b: 55 WP

Ectoedemia cerris (Zimmermann, 1944) Szőcs, 1978: 266 WP

Nepticula cerris Zimmermann, 1944: 121

Nepticula montissancti Skala, 1948: 121 (syn: van Nieukerken, 1985b: 54)

Stigmella cerris (Zimmermann, 1944) Hering, 1957: 866

Ectoedemia subbimaculella complex (van Nieukerken, 1985b: 56) (next 4 species)

Ectoedemia subbimaculella (Haworth, 1828) Bradley et al., 1972: 3 WP

Tinea subbimaculella Haworth, 1828: 583

‡ *Microsetia nigrociliella* Stephens, 1829: 208 NN (syn: van Nieukerken, 1985b: 57)

Microsetia nigrociliella Stephens, 1834: 267 (syn: van Nieukerken, 1985b: 57)

‡ *Nepticula cursoriella* Heyden, 1843: 209 NN (syn: Herrich-Schäffer, 1855a: 356)

Nepticula cursoriella Zeller, 1848: 326 (syn: Herrich-Schäffer, 1855a: 356)

Nepticula bistrimaculella Heyden, 1861: 40 (syn: van Nieukerken & Johansson, 1987: 462)

Microsetia subbimaculella (Haworth, 1828) Stephens, 1829: 208

Nepticula subbimaculella (Haworth, 1828) Stainton, 1849: 29

Stigmella subbimaculella (Haworth, 1828) Fletcher & Clutterbuck, 1945: 61

Dechtiria subbimaculella (Haworth, 1828) Beirne, 1945: 206

Trifurcula subbimaculella (Haworth, 1828) Johansson, 1971: 245

Stigmella bistrimaculella (Heyden, 1861) Gerasimov, 1952: 231

Ectoedemia phyllotomella (Klimesch, 1948) van Nieukerken, 1985b: 62[29] WP

Stigmella phyllotomella Klimesch, 1948a: 166[29]

Ectoedemia heringi (Toll, 1934) Borkowski, 1975: 491 WP

Nepticula heringi Toll, 1934a: 3

Nepticula quercifoliae Toll, 1934b: 71 (syn: Borkowski, 1975: 491)

Nepticula sativella Klimesch, 1936: 208 (syn: van Nieukerken, 1985b: 59)

Nepticula zimmermanni Hering, 1942: 26 (syn: van Nieukerken, 1985b: 59)

Stigmella heringi (Toll, 1934) Hering, 1957: 867

Trifurcula heringi (Toll, 1934) Kasy, 1979: 4

Stigmella sativella (Klimesch, 1936) Klimesch, 1948b: 74

Stigmella quercifoliae (Toll, 1934) Hering, 1957: 867

Ectoedemia quercifoliae (Toll, 1934) Bradley et al., 1972: 3

Stigmella zimmermanni (Hering, 1942) Klimesch, 1951a: 65
Trifurcula zimmermanni (Hering, 1942) Kasy, 1979: 4
Ectoedemia zimmermanni (Hering, 1942) Szőcs, 1981: 210
Ectoedemia liechtensteini (Zimmermann, 1944) Szőcs, 1978: 266 WP
Nepticula liechtensteini Zimmermann, 1944: 119
Stigmella liechtensteini (Zimmermann, 1944) Hering, 1957: 866

***Ectoedemia platanella* group** (Wilkinson & Newton, 1981: 51)
Ectoedemia similella (Braun, 1917) Wilkinson & Newton, 1981: 56 NEA
Nepticula similella Braun, 1917: 188
Ectoedemia platanella (Clemens, 1861) Wilkinson & Scoble, 1979: 89 NEA
Nepticula platanella Clemens, 1861: 83
Nepticula maximella Chambers, 1873: 126 (syn: Braun, 1917: 187)
Ectoedemia clemensella (Chambers, 1873) Wilkinson & Scoble, 1979: 86 NEA
Nepticula clemensella Chambers, 1873: 125
Ectoedemia virgulae (Braun, 1927) Wilkinson & Newton, 1981: 59 NEA
Nepticula virgulae Braun, 1927: 198

***Ectoedemia ornatella* group** (Puplesis, 1984b: 584)
Ectoedemia ivinskisi Puplesis, 1984a: 120 EP
Ectoedemia olvina Puplesis, 1984a: 119 EP
Ectoedemia ornatella Puplesis, 1984a: 120 EP

***Ectoedemia suberis* group** (van Nieukerken, 1985b: 38)
Ectoedemia chasanella Puplesis, 1984a: 124 EP
Ectoedemia aegilopidella (Klimesch, 1978) van Nieukerken, 1985b: 42 WP
Trifurcula aegilopidella Klimesch, 1978b: 269
Ectoedemia caradjai (Groschke, 1944) Szőcs, 1981: 211 WP
Nepticula caradjai Groschke, 1944: 118
Stigmella caradjai (Groschke, 1944) Klimesch, 1951b: 65
Trifurcula caradjai (Groschke, 1944) Klimesch, 1978a: 250
Ectoedemia andalusiae van Nieukerken, 1985b: 41 WP
Ectoedemia suberis (Stainton, 1869) van Nieukerken, 1985b: 40 WP
Nepticula suberis Stainton, 1869b: 229
Nepticula viridella Mendes, 1910b: 165 (syn: van Nieukerken, 1985b: 40)
Stigmella suberis (Stainton, 1869) Gerasimov, 1952: 262
Stigmella viridella (Mendes, 1910) Gerasimov, 1952: 260
Ectoedemia phaeolepis van Nieukerken, A. Laštuvka & Z. Laštuvka, 2010: 38 WP
Ectoedemia hendrikseni A. Laštuvka, Z. Laštuvka & van Nieukerken in van Nieukerken et al., 2010: 31 WP
Ectoedemia heckfordi van Nieukerken, A. Laštuvka & Z. Laštuvka, 2010: 34 WP
Ectoedemia ortiva Rocienė & Stonis, 2013: 76 EP
Ectoedemia paraortiva Rocienė & Stonis in Stonis & Rocienė, 2013: 210 EP

***Ectoedemia angulifasciella* group** (Wilkinson et al., 1983: 211)
Ectoedemia rubifoliella group (Wilkinson & Newton, 1981: 61)
Ectoedemia occultella group (van Nieukerken, 1985b: 78)
Ectoedemia hexapetalae (Szőcs, 1957) van Nieukerken, 1985b: 68 WP
Nepticula utensis var. *hexapetalae* Szőcs, 1957: 322
Nepticula hexapetalae Szőcs, 1957 (Szőcs, 1965:79)
Trifurcula hexapetalae (Szőcs, 1957) Kasy, 1980: 47
Ectoedemia rosae van Nieukerken & Berggren, 2011: 182 WP
Ectoedemia rosiphila Puplesis in Puplesis et al., 1992: 55 EP
Ectoedemia marmaropa (Braun, 1925) Wilkinson & Newton, 1981: 49 NEA
Nepticula marmaropa Braun, 1925b: 225
Ectoedemia spiraeae Gregor & Povolny, 1983: 174 [66] WP,EP?
‡ *Stigmella spireae* Gregor & Povolny, 1955: 124 NNLM
‡ *Nepticula spireae* (Gregor & Povolny, 1955) Szőcs, 1968: 229 NNLM
Ectoedemia jacutica Puplesis, 1988: 26 [66] EP
Ectoedemia agrimoniae (Frey, 1858) Bradley et al., 1972: 2 WP
Nepticula agrimoniae Frey, 1858c: 44
Nepticula agrimoniae Hofmann, 1858: 188
Nepticula agrimoniella Herrich-Schäffer, 1860: 60 UE
Dechtiria agrimoniae (Frey, 1858) Beirne, 1945: 205
Stigmella agrimoniae (Frey, 1858) Gerasimov, 1952: 224
Trifurcula agrimoniae (Frey, 1858) Johansson, 1971: 245
Stigmella agrimoniella (Herrich-Schäffer, 1860) Le Marchand, 1946a: 217
‡ *Nepticula agrimomella* Rössler, 1881: 337 ISS
Ectoedemia nyssaefoliella (Chambers, 1880) Wilkinson & Newton, 1981: 67 NEA
Nepticula nyssaefoliella Chambers, 1880b: 66
Ectoedemia pilosae Puplesis, 1984a: 123 EP
Ectoedemia picturata Puplesis, 1985c: 65 EP
Ectoedemia minimella (Zetterstedt, 1839) van Nieukerken, 1985b: 80 [67] WP,EP,NEA
Elachista minimella Zetterstedt, 1839: 1011
Nepticula woolhopiella Stainton, 1887: 262 (syn: van Nieukerken, 1985b: 80)
Nepticula canadensis Braun, 1917: 185 **syn. n.** [67]
Nepticula viridicola Weber, 1938: 211 (syn: van Nieukerken, 1985b: 80)
Nepticula vividicola Weber, 1938: 211 IOS
Stigmella woolhopiella (Stainton, 1887) Fletcher & Clutterbuck, 1945: 60
Dechtiria woolhopiella (Stainton, 1887) Beirne, 1945: 205
Trifurcula woolhopiella (Stainton, 1887) Johansson, 1971: 245
Ectoedemia woolhopiella (Stainton, 1887) Borkowski, 1975: 493
Ectoedemia mediofasciella auct. [misapplied] Bradley et al., 1972: 2
Trifurcula mediofasciella auct. [misapplied] Karsholt & Nielsen, 1976: 18
Stigmella viridicola (Weber, 1938) Klimesch, 1948b: 70
Stigmella canadensis (Braun, 1917) Davis & Wilkinson, 1983: 3
Ectoedemia canadensis (Braun, 1917) Wilkinson, 1981: 94

Ectoedemia occultella (Linnaeus, 1767) Robinson & Nielsen, 1983: 221 [68] WP,EP,NEA

Phalaena occultella Linnaeus, 1767: 899
Tinea strigilella Thunberg, 1794: 87 (syn: Robinson & Nielsen, 1983: 221)
Tinea mucidella Hübner, 1817: pl. 65: Fig. 435 (syn: Zeller, 1839: 215)
Tinea mediofasciella Haworth, 1828: 584 (syn: van Nieukerken, 1985b: 78)
Lyonetia argentipedella Zeller, 1839: 215 (syn: Robinson & Nielsen, 1983: 221)
Nepticula flexuosella Fologne, 1859: 140 (syn: van Nieukerken & Johansson, 1987: 462)
Nepticula lindquisti Freeman, 1962: 899 (syn: van Nieukerken, 1985b: 80) [68]
Elachista mucidella (Hübner, 1817) Treitschke, 1833: 179
Lyonetia mucidella (Hübner, 1817) Duponchel, 1844: 378
Nepticula argentipedella (Zeller, 1839) Heyden, 1843: 209
Stigmella argentipedella (Zeller, 1839) Fletcher & Clutterbuck, 1945: 60
Dechtiria argentipedella (Zeller, 1839) Beirne, 1945: 205
Ectoedemia argentipedella (Zeller, 1839) Bradley et al., 1972: 2
Trifurcula argentipedella (Zeller, 1839) Johansson, 1971: 245
Ectoedemia lindquisti (Freeman, 1962) Wilkinson & Scoble, 1979: 83
Microsetia mediofasciella (Haworth, 1828) Stephens, 1829: 208

Ectoedemia angulifasciella (Stainton, 1849) Bradley et al., 1972: 2 (Fig. 38) WP

Nepticula angulifasciella Stainton, 1849: 29
Nepticula schleichiella Frey, 1870: 286 (syn: van Nieukerken, 1985b: 69)
Nepticula brunniella Sauber, 1904: 55 (syn: van Nieukerken, 1985b: 69)
Nepticula utensis Weber, 1937: 669 (syn: van Nieukerken, 1985b: 69)
Nepticula minorella Zimmermann, 1944: 118 (syn: van Nieukerken, 1985b: 69)
Stigmella angulifasciella (Stainton, 1849) Vári, 1944b: xxv
Dechtiria angulifasciella (Stainton, 1849) Beirne, 1945: 205
Trifurcula angulifasciella (Stainton, 1849) Johansson, 1971: 245
Stigmella schleichiella (Frey, 1870) Gerasimov, 1952: 259
Stigmella utensis (Weber, 1937) Klimesch, 1948b: 72
Stigmella minorella (Zimmermann, 1944) Klimesch, 1961: 739

Ectoedemia rubivora complex (van Nieukerken et al., 2012a: 7) (next 3 species)

Ectoedemia arcuatella (Herrich-Schäffer, 1855) Bradley et al., 1972: 2 WP,EP

Nepticula arcuatella Herrich-Schäffer, 1855a: 354
Nepticula arcuosella Doubleday, 1859: 36 UE
Stigmella arcuatella (Herrich-Schäffer, 1855) Fletcher & Clutterbuck, 1945: 60
Dechtiria arcuatella (Herrich-Schäffer, 1855) Beirne, 1945: 206
Trifurcula arcuatella (Herrich-Schäffer, 1855) Johansson, 1971: 245

Ectoedemia atricollis (Stainton, 1857) Bradley et al., 1972: 2 WP,EP

Nepticula atricollis Stainton, 1857a: 112
Nepticula atricolella Doubleday, 1859: 36 UE
Nepticula aterrima Wocke, 1865: 270 (syn: van Nieukerken, 1985b: 71)
Nepticula staphyleae Zimmermann, 1944: 117 (syn: van Nieukerken, 1985b: 71)
Stigmella atricollis (Stainton, 1857) Vári, 1944b: xxv

Dechtiria atricollis (Stainton, 1857) Vári, 1951: 197
Trifurcula atricollis (Stainton, 1857) Johansson, 1971: 245
Stigmella aterrima (Wocke, 1865) Gerasimov, 1952: 228
Stigmella staphyleae (Zimmermann, 1944) Hering, 1957: 1027
Ectoedemia staphyleae (Zimmermann, 1944) Borkowski, 1975: 493
‡ *Nepticula malivora* Toll, 1934b: 70 NNLM (syn: Skala, 1948: 121)
‡ *Nepticula atricollis* var. *aterrimoides* Skala, 1940: 143 NNLM (syn: Skala, 1948: 121)
‡ *Nepticula atricollis* var. *prunivora* Skala, 1941b: 77 NNLM

Ectoedemia rubivora (Wocke, 1860) Bradley et al., 1972: 2 WP
Nepticula rubivora Wocke, 1860: 132
Stigmella rubivora (Wocke, 1860) Fletcher & Clutterbuck, 1945: 60
Dechtiria rubivora (Wocke, 1860) Beirne, 1945: 205
Trifurcula rubivora (Wocke, 1860) Johansson, 1971: 245

Ectoedemia spinosella (Joannis, 1908) Bradley et al., 1972: 2 WP
Nepticula spinosella Joannis, 1908: 328
Ectoedemia albiformae Puplesis & Diškus, 2003a: 186 (syn: van Nieukerken et al., 2010: 70)
Stigmella spinosella (Joannis, 1908) Klimesch, 1951b: 62
Dechtiria spinosella (Joannis, 1908) Emmet, 1971b: 244
Trifurcula spinosella (Joannis, 1908) Johansson, 1971: 245

Ectoedemia mahalebella (Klimesch, 1936) Szőcs, 1978: 266 WP
Nepticula mahalebella Klimesch, 1936: 207
Stigmella mahalebella (Klimesch, 1936) Lhomme, 1945: 155

Ectoedemia erythrogenella (Joannis, 1908) Emmet, 1974a: 129 WP
Nepticula erythrogenella Joannis, 1908: 327
Stigmella erythrogenella (Joannis, 1908) Gerasimov, 1952: 238
Trifurcula erythrogenella (Joannis, 1908) Leraut, 1980: 49
‡ *Stigmella erythrogenella* ab. *juncta* Dufrane, 1949: 9

Ectoedemia rubifoliella (Clemens, 1860) Wilkinson & Scoble, 1979: 90 NEA
Nepticula rubifoliella Clemens, 1860: 214

Ectoedemia ulmella (Braun, 1912) Wilkinson & Scoble, 1979: 91 [69] NEA
Nepticula ulmella Braun, 1912: 87
Ectoedemia andrella Wilkinson, 1981: 102 **syn. n.** [69]

Ectoedemia ingloria Puplesis, 1988: 280 EP

Ectoedemia insignata Puplesis, 1988: 281 EP

Ectoedemia petrosa Puplesis, 1988: 282 EP

Ectoedemia tadshikiella Puplesis, 1988: 25 WP,EP

***Ectoedemia* - unplaced species**

Ectoedemia fuscivittata Puplesis & Robinson, 2000: 42 NEO

Stigmellites Kernbach, 1967: 104 (TS/OD,M: *Stigmellites heringi* Kernbach, 1967)
Ophiheliconoma Krassilov, 2008: 100 (TS/OD,M: *Ophiheliconoma resupinata* Krassilov, 2008) (syn: Doorenweerd et al., 2015a: 309)
Stigmellites almeidae (Martins-Neto, 1989) Doorenweerd et al., 2015a: 315 NEO†
Nepticula almeidae Martins-Neto, 1989: 381
Stigmella almeidae (Martins-Neto, 1989) Sohn et al., 2012: 22
Stigmellites baltica Kozlov, 1988: 30 WP†
Stigmellites carpiniorientalis Straus, 1977: 60 WP†
Stigmellites centennis Jarzembowski, 1989: 448 WP†
Stigmellites fossilis (Heyden, 1862) Kozlov, 1988: 31 WP†
Nepticula fossilis Heyden, 1862: 77
Stigmellites gossi Jarzembowski, 1989: 448 WP†
Stigmellites heringi Kernbach, 1967: 104 WP†
Stigmellites kzyldzharica Kozlov, 1988: 32 EP†
Stigmellites messelensis Straus, 1976: 445 WP†
Stigmellites pliotityrella Kernbach, 1967: 105 WP†
Stigmellites resupinata (Krassilov, 2008) Doorenweerd et al., 2015a: 309 WP†
Ophiheliconoma resupinata Krassilov, 2008: 100
Stigmellites samsonovi Kozlov, 1988: 33 EP†
Stigmellites serpentina Kozlov, 1988: 32 EP†
Stigmellites sharovi Kozlov, 1988: 33 EP†
Stigmellites tyshchenkoi Kozlov, 1988: 33 EP†
Stigmellites zelkovae Straus, 1977: 61 WP†

Nomina dubia et oblita

Nepticula alpinella Herrich-Schäffer, 1863b: 170 [70] NO WP
Nepticula alticolella Herrich-Schäffer, 1863c: 182 [70] NO WP
Nepticula reuttiella Herrich-Schäffer, 1863c: 182 [70] NO WP
Nepticula oritis Meyrick, 1910: 229 [71] ND OR
Nepticula xuthomitra Meyrick, 1921b: 140 [72] ND AFR
Nepticula anguinella Clemens, 1861: 85 [73] ND NEA
Ectoedemia anguinella (Clemens, 1861) Wilkinson, 1981: 98 ND
Nepticula platea Clemens, 1861: 85 [73] ND NEA
Ectoedemia platea (Clemens, 1861) Wilkinson, 1981: 98 ND

Unplaced unavailable names [74]

‡ *Nepticula brunensis* Skala, 1939g: 144 NNLM WP
‡ *Nepticula buhri* Skala, 1938: 43 NNLM WP
‡ *Nepticula sorbifoliella* Skala, 1939g: 144 NNLM WP
‡ *Nepticula tentationis* Hoffmann, 1893: 215 NN WP
‡ *Nepticula ulmi* Skala, 1934a: 51 NNLM WP
‡ *Stigmella acernella* Dovnar-Zapolski & Tomilova, 1978: 27 NNLM EP
‡ *Stigmella amygdaliella* Dovnar-Zapolski, 1969: 23 NNLM EP
‡ *Stigmella apocynella* Gerasimov, 1937: 284 NNLM EP

‡ *Stigmella atraphaxidella* Dovnar-Zapolski, 1969: 29 NNLM EP
‡ *Stigmella betulivora* Dovnar-Zapolski, 1969: 32 NNLM EP
‡ *Stigmella crataegifolia* Dovnar-Zapolski, 1969: 49 NNLM EP
‡ *Stigmella loniceraefolia* Dovnar-Zapolski, 1969: 67 NNLM EP
‡ *Stigmella loniceraevora* Dovnar-Zapolski, 1969: 67 NNLM EP
‡ *Stigmella prunivora* Dovnar-Zapolski, 1969: 90 NNLM WP
‡ *Stigmella pseudoanomalella* Dovnar-Zapolski, 1969: 94 NNLM EP
‡ *Stigmella roseifolia* Dovnar-Zapolski, 1969: 94 NNLM EP
‡ *Stigmella roseivora* Dovnar-Zapolski, 1969: 94 NNLM EP
‡ *Stigmella rosella* Dovnar-Zapolski, 1969: 95 NNLM EP

FAMILY **OPOSTEGIDAE** Meyrick, 1893: 479 (TG: *Opostega* Zeller, 1839)
Family Opostegides Meyrick, 1893: 479 (TG: *Opostega* Zeller, 1839)
Subfamily Opostegoidinae Kozlov, 1987: 856 (TG: *Opostegoides* Kozlov, 1985), **syn. n.**
Subfamily Oposteginae Meyrick, 1893 (TG: *Opostega* Zeller, 1839)

Notiopostega Davis, 1989: 30 (TS/OD,M: *Notiopostega atrata* Davis, 1989)
Notiopostega atrata Davis, 1989: 32 NEO

Eosopostega Davis, 1989: 41 (TS/OD,M: *Eosopostega issikii* Davis, 1989)
Eosopostega issikii Davis, 1989: 42 EP
Eosopostega armigera Puplesis & Robinson, 1999: 29 OR

Neopostega Davis & Stonis, 2007: 34 (TS/OD: *Neopostega petila* Davis & Stonis, 2007: 38)
Neopostega asymmetra Davis & Stonis, 2007: 37 NEO
Neopostega distola Davis & Stonis, 2007: 39 NEO
Neopostega falcata Davis & Stonis, 2007: 36 NEO
Neopostega longispina Davis & Stonis, 2007: 36 NEO
Neopostega nigrita Heppner & Davis, 2009: 31 NEO
Neopostega petila Davis & Stonis, 2007: 38 NEO

Paralopostega Davis, 1989: 52 (TS/OD: *Opostega callosa* Swezey, 1921)
Paralopostega callosa (Swezey, 1921) Davis, 1989: 72 AUS
Opostega callosa Swezey, 1921: 532
Paralopostega dives (Walsingham, 1907) Davis, 1989: 72 AUS
Opostega dives Walsingham, 1907: 711
Paralopostega filiforma (Swezey, 1921) Davis, 1989: 72 AUS
Opostega filiforma Swezey, 1921: 534
Paralopostega maculata (Walsingham, 1907) Davis, 1989: 72 AUS
Opostega maculata Walsingham, 1907: 711
Paralopostega peleana (Swezey, 1921) Davis, 1989: 73 AUS
Opostega peleana Swezey, 1921: 534

Paralopostega serpentina (Swezey, 1921) Davis, 1989: 73 AUS
Opostega serpentina Swezey, 1921: 533

Opostegoides Kozlov, 1985: 54 (TS/OD: *Opostega minodensis* Kuroko, 1982) [75]
Opostegoides menthinella (Mann, 1855) Davis, 1989: 72 WP
Opostega menthinella Mann, 1855: 568
Opostega snelleni Nolcken, 1882: 197 (syn: van Nieukerken, 1996: 300)
Opostegoides albella Sinev, 1990: 102 EP
Opostegoides bicolorella Sinev, 1990: 105 EP
Opostegoides minodensis (Kuroko, 1982) Kozlov, 1985: 54 EP
Opostega minodensis Kuroko, 1982: 50, 448
Opostegoides omelkoi Kozlov, 1985: 57 EP
Opostegoides padiensis Sinev, 1990: 105 EP
Opostegoides sinevi Kozlov, 1985: 55 EP
Opostegoides argentisoma Puplesis & Robinson, 1999: 22 OR
Opostegoides auriptera Puplesis & Robinson, 1999: 28 OR
Opostegoides cameroni Puplesis & Robinson, 1999: 27 OR
Opostegoides epistolaris (Meyrick, 1911b) Puplesis & Robinson, 1999: 20 OR
Opostega epistolaris Meyrick, 1911b: 108
Opostegoides flavimacula Puplesis & Robinson, 1999: 27 OR
Opostegoides gorgonea Puplesis & Robinson, 1999: 22 OR
Opostegoides index (Meyrick, 1922) Puplesis & Robinson, 1999: 20 OR
Opostega index Meyrick, 1922: 557
Opostegoides longipedicella Puplesis & Robinson, 1999: 26 OR
Opostegoides malaysiensis Davis, 1989: 52 OR
Opostegoides nephelozona (Meyrick, 1915) Puplesis & Robinson, 1999: 19 OR
Opostega nephelozona Meyrick, 1915b: 352
Opostegoides pelorrhoa (Meyrick, 1915) Puplesis & Robinson, 1999: 18 OR
Opostega pelorrhoa Meyrick, 1915b: 352
Opostegoides spinifera Puplesis & Robinson, 1999: 26 OR
Opostegoides tetroa (Meyrick, 1907) Puplesis & Robinson, 1999: 18 OR
Opostega tetroa Meyrick, 1907: 986
Opostegoides thailandica Puplesis & Robinson, 1999: 23 OR
Opostegoides uvida (Meyrick, 1915) Puplesis & Robinson, 1999: 19 OR
Opostega uvida Meyrick, 1915b: 352
Opostegoides granifera (Meyrick, 1913), **comb. n.** [75] AFR
Opostega granifera Meyrick, 1913: 327
Opostegoides melitardis (Meyrick, 1918), **comb. n.** [75] AFR
Opostega melitardis Meyrick, 1918a: 41
Opostegoides pelocrossa (Meyrick, 1928), **comb. n.** [75] AFR
Opostega pelocrossa Meyrick, 1928a: 396
Opostegoides praefusca (Meyrick, 1913), **comb. n.** [75] AFR
Opostega praefusca Meyrick, 1913: 327

Opostegoides gephyraea (Meyrick, 1880) Davis, 1989: 72 AUS
Opostega gephyraea Meyrick, 1880: 176
Opostegoides scioterma (Meyrick, 1920) Kozlov, 1985: 55 NEA
Opostega scioterma Meyrick, 1920c: 358

Opostega Zeller, 1839: 214 (TS/SD (Walsingham, 1914: 349): *Elachista salaciella* Treitschke, 1833)
Opostega cretatella Chrétien, 1915: 364 [76] WP,EP
Opostega rezniki Kozlov, 1985: 51 **syn. n.** [76]
Opostega kuznetzovi Kozlov, 1985: 53 WP,EP
Opostega salaciella (Treitschke, 1833) Zeller, 1939: 214 WP
Elachista salaciella Treitschke, 1833: 180
Opostega reliquella Zeller, 1848: 282
Opostega spatulella Herrich-Schäffer, 1855a: 360 (Fig. 39) WP,EP
Opostega nepticulella Bruand, 1859: 691 (syn: Leraut, 1997: 80)
Opostega bimaculatella N.R. Rothschild, 1912: 29 (syn: van Nieukerken, 1990a: 368)
Opostega costantiniella Costantini in Turati, 1923: 70 (syn: van Nieukerken, 1990a: 368)
Opostega angulata Gerasimov, 1930: 45 (syn: Puplesis et al., 1996: 192)
Opostega heringella Mariani, 1937: 12 (syn: van Nieukerken, 1990a: 368)
Opostega stekolnikovi Kozlov, 1985: 53 WP
Opostega afghani Davis, 1989: 62 EP
Opostega chalcophylla Meyrick, 1910: 229 OR

"*Opostega*" (unplaced African species) [77]
"Opostega" cirrhacma Meyrick, 1911a: 237 AFR
"Opostega" diplardis Meyrick, 1921b: 123 AFR
"Opostega" radiosa Meyrick, 1913: 327 AFR

"*Opostega*" (unplaced Australian species) [77]
"Opostega" arthrota Meyrick, 1915b: 352 AUS
"Opostega" atypa Turner, 1923: 179 AUS
"Opostega" basilissa Meyrick, 1893: 606 AUS
"Opostega" brithys Turner, 1923: 179 AUS
"Opostega" chalcoplethes Turner, 1923: 178 AUS
"Opostega" chalinias Meyrick, 1893: 607 AUS
"Opostega" chordacta Meyrick, 1915b: 351 AUS
"Opostega" diorthota Meyrick, 1893: 607 AUS
"Opostega" horaria Meyrick, 1921d: 457 AUS
"Opostega" luticilia Meyrick, 1915b: 351 AUS
"Opostega" monotypa Turner, 1923: 179 AUS
"Opostega" nubifera Turner, 1900: 23 AUS
"Opostega" orestias Meyrick, 1880: 175 AUS

"Opostega" phaeospila Turner, 1923: 179 AUS
"Opostega" scoliozona Meyrick, 1915b: 351 AUS
"Opostega" stiriella Meyrick, 1880: 175 AUS
"Opostega" xenodoxa Meyrick, 1893: 608 AUS

Pseudopostega Kozlov, 1985: 53 (TS/OD: *Tinea auritella* Hübner, 1813)
Palearctic species
Pseudopostega auritella (Hübner, 1813) Davis, 1989: 76 WP,EP
Tinea auritella Hübner, 1813: Pl. 57: Fig. 387
Leucoptera auritella (Hübner, 1813) Hübner, 1825: 426
Opostega auritella (Hübner, 1813) Zeller, 1939: 214
Pseudopostega chalcopepla (Walsingham, 1908) van Nieukerken, 1996: 27 [78] WP
Opostega chalcopepla Walsingham, 1908b: 228
Pseudopostega cyrneochalcopepla Nel & Varenne, 2012: 11 **syn. n.** [78]
‡ *Opostega rosmarinella* Staudinger, 1894 (syn: Walsingham, 1908b: 228) NN
Pseudopostega crepusculella (Zeller, 1839) Davis, 1989: 76 (Fig. 40) WP,EP
Opostega crepusculella Zeller, 1839: 214
Oecophora crepusculella Duponchel, 1843: 337 JSH
Opostega crepusculella lvovskyi Kozlov, 1985: 54

Oriental species
Pseudopostega alleni Puplesis & Robinson, 1999: 40 OR
Pseudopostega amphivittata Puplesis & Robinson, 1999: 39 OR
Pseudopostega brevicaudata Remeikis & Stonis in Stonis et al., 2013a: 183 OR
Pseudopostega epactaea (Meyrick, 1907) Puplesis & Robinson, 1999: 32 OR
Opostega epactaea Meyrick, 1907: 985
Pseudopostega euryntis (Meyrick, 1907) Puplesis & Robinson, 1999: 44 OR
Opostega euryntis Meyrick, 1907: 985
Pseudopostega frigida (Meyrick, 1906) Puplesis & Robinson, 1999: 32 OR
Opostega frigida Meyrick, 1906a: 416
Pseudopostega fungina Puplesis & Robinson, 1999: 42 OR
Pseudopostega indonesica Puplesis & Robinson, 1999: 41 OR
Pseudopostega javae Puplesis & Robinson, 1999: 39 OR
Pseudopostega machaerias (Meyrick, 1907) Puplesis & Robinson, 1999: 30 OR
Opostega machaerias Meyrick, 1907: 986
Pseudopostega myxodes (Meyrick, 1916) Puplesis & Robinson, 1999: 34 OR
Opostega myxodes Meyrick, 1916a: 619
Pseudopostega nepalensis Puplesis & Robinson, 1999: 37 OR
Pseudopostega nigrimaculella Puplesis & Robinson, 1999: 40 OR
Pseudopostega parvilineata Puplesis & Robinson, 1999: 31 OR
Pseudopostega saturella Puplesis & Robinson, 1999: 38 OR
Pseudopostega similantis Puplesis & Robinson, 1999: 33 OR
Pseudopostega spilodes (Meyrick, 1915b) Puplesis & Robinson, 1999: 45 OR
Opostega spilodes Meyrick, 1915b: 351

Pseudopostega strigulata Puplesis & Robinson, 1999: 45 OR
Pseudopostega subviolacea (Meyrick, 1920) Puplesis & Robinson, 1999: 45 OR
Opostega subviolacea Meyrick, 1920c: 357
Pseudopostega sumbae Puplesis & Robinson, 1999: 37 OR
Pseudopostega velifera (Meyrick, 1920) Puplesis & Robinson, 1999: 34 OR
Opostega velifera Meyrick, 1920c: 357
Pseudopostega zelopa (Meyrick, 1905) Puplesis & Robinson, 1999: 43 OR
Opostega zelopa Meyrick, 1905: 613

African species [79]

Pseudopostega amphimitra (Meyrick, 1913), **comb. n.** [79] AFR
Opostega amphimitra Meyrick, 1913: 328
Pseudopostega bellicosa (Meyrick, 1911a) Davis, 1989: 76 AFR
Opostega bellicosa Meyrick, 1911a: 236
Pseudopostega clastozona (Meyrick, 1913) Davis, 1989: 76 AFR
Opostega clastozona Meyrick, 1913: 327
Pseudopostega idiocoma (Meyrick, 1918), **comb. n.** [79] AFR
Opostega idiocoma Meyrick, 1918a: 42
Pseudopostega orophoxantha (Meyrick, 1921), **comb. n.** [79] AFR
Opostega orophoxantha Meyrick, 1921b: 124
Pseudopostega phaeosoma (Meyrick, 1928), **comb. n.** [79] AFR
Opostega phaeosoma (Meyrick, 1928a): 396
Pseudopostega symbolica (Meyrick, 1914), **comb. n.** [79] AFR
Opostega symbolica Meyrick, 1914: 203
Pseudopostega tincta (Meyrick, 1918), **comb. n.** [79] AFR
Opostega tincta Meyrick, 1918a: 41

Nearctic species

Pseudopostega acidata (Meyrick, 1915) Davis, 1989: 75 NEA,NEO
Opostega acidata Meyrick, 1915a: 240
Pseudopostega albogaleriella (Clemens, 1862) Davis, 1989: 76 NEA
Opostega albogaleriella Clemens, 1862: 131
Opostega napaeella Clemens, 1872: 42 (syn: Davis, 1983: 3)
Opostega bistrigulella Braun, 1918: 245 (syn: Davis & Stonis, 2007: 71)
Opostega nonstrigella Chambers, 1881: 296 (syn: Forbes, 1923: 161)
Pseudopostega napaeella (Clemens, 1872) Davis, 1989: 76
Pseudopostega bistrigulella (Braun, 1918) Davis, 1989: 76
Pseudopostega nonstrigella (Chambers, 1881) Davis, 1989: 76
Pseudopostega cretea (Meyrick, 1920) Davis, 1989: 76 NEA
Opostega cretea Meyrick, 1920c: 358
Pseudopostega floridensis Davis & Stonis, 2007: 57 NEA
Pseudopostega kempella (Eyer, 1967) Davis, 1989: 76 NEA,NEO
Opostega kempella Eyer, 1967: 39
Pseudopostega parakempella Davis & Stonis, 2007: 100 NEA,NEO

Pseudopostega quadristrigella (Chambers, 1875) Davis, 1989: 77 NEA
Opostega quadristrigella Chambers, 1875b: 106
Opostega accessoriella Frey & Boll, 1876: 216 (syn: McDunnough, 1939: 100)
Pseudopostega accessoriella (Frey & Boll, 1876) Davis, 1989: 75
Pseudopostega texana Davis & Stonis, 2007: 115 NEA
Pseudopostega venticola (Walsingham, 1897) Davis, 1989: 77 NEA,NEO
Opostega venticola Walsingham, 1897: 140

Neotropic species
Pseudopostega abrupta (Walsingham, 1897) Davis, 1989: 75 NEO
Opostega abrupta Walsingham, 1897: 139
Pseudopostega acrodicra Davis & Stonis, 2007: 122 NEO
Pseudopostega acuminata Davis & Stonis, 2007: 89 NEO
Pseudopostega adusta (Walsingham, 1897) Davis, 1989: 76 NEO
Opostega adusta Walsingham, 1897: 140
Pseudopostega apoclina Davis & Stonis, 2007: 131 NEO
Pseudopostega latifurcata apoclina Davis & Stonis, 2007: 131
Pseudopostega apotoma Davis & Stonis, 2007: 65 NEO
Pseudopostega attenuata Davis & Stonis, 2007: 76 NEO
Pseudopostega beckeri Davis & Stonis, 2007: 136 NEO
Pseudopostega bicornuta Davis & Stonis, 2007: 138 NEO
Pseudopostega bidorsalis Davis & Stonis, 2007: 127 NEO
Pseudopostega brachybasis Davis & Stonis, 2007: 142 NEO
Pseudopostega breviapicula Davis & Stonis, 2007: 85 NEO
Pseudopostega brevifurcata Davis & Stonis, 2007: 120 NEO
Pseudopostega brevivalva Davis & Stonis, 2007: 121 NEO
Pseudopostega caulifurcata Davis & Stonis, 2007: 123 NEO
Pseudopostega clavata Davis & Stonis, 2007: 105 NEO
Pseudopostega colognatha Davis & Stonis, 2007: 90 NEO
Pseudopostega concava Davis & Stonis, 2007: 119 NEO
Pseudopostega congruens (Walsingham, 1914) Davis, 1989: 76 NEO
Opostega congruens Walsingham, 1914: 350
Pseudopostega conicula Davis & Stonis, 2007: 78 NEO
Pseudopostega constricta Davis & Stonis, 2007: 141 NEO
Pseudopostega contigua Davis & Stonis, 2007: 129 NEO
Pseudopostega crassifurcata Davis & Stonis, 2007: 117 NEO
Pseudopostega curtarama Davis & Stonis, 2007: 116 NEO
Pseudopostega denticulata Davis & Stonis, 2007: 74 NEO
Pseudopostega didyma Davis & Stonis, 2007: 109 NEO
Pseudopostega diskusi Davis & Stonis, 2007: 67 NEO
Pseudopostega divaricata Davis & Stonis, 2007: 128 NEO
Pseudopostega dorsalis Davis & Stonis, 2007: 98 NEO
Pseudopostega dorsalis dorsalis Davis & Stonis, 2007: 98 NEO
Pseudopostega duplicata Davis & Stonis, 2007: 108 NEO

Pseudopostega ecuadoriana Davis & Stonis, 2007: 134 NEO
Pseudopostega elachista (Walsingham, 1914) Davis, 1989: 76 NEO
Opostega elachista Walsingham, 1914: 350
Pseudopostega fasciata Davis & Stonis, 2007: 99 NEO
Pseudopostega dorsalis fasciata Davis & Stonis, 2007: 99
Pseudopostega ferruginea Davis & Stonis, 2007: 54 NEO
Pseudopostega fumida Davis & Stonis, 2007: 62 NEO
Pseudopostega galapagosae Davis & Stonis, 2007: 93 NEO
Pseudopostega gracilis Davis & Stonis, 2007: 63 NEO
Pseudopostega lateriplicata Davis & Stonis, 2007: 59 NEO
Pseudopostega latiapicula Davis & Stonis, 2007: 133 NEO
Pseudopostega latifurcata Davis & Stonis, 2007: 130 NEO
Pseudopostega latifurcata latifurcata Davis & Stonis, 2007: 130 NEO
Pseudopostega latiplana Remeikis & Stonis in Remeikis et al., 2009: 283 NEO
Pseudopostega latisaccula Davis & Stonis, 2007: 75 NEO
Pseudopostega lobata Davis & Stonis, 2007: 104 NEO
Pseudopostega longifurcata Davis & Stonis, 2007: 141 NEO
Pseudopostega longipedicella Davis & Stonis, 2007: 102 NEO
Pseudopostega mexicana Remeikis & Stonis in Remeikis et al., 2009: 282 NEO
Pseudopostega microacris Davis & Stonis, 2007: 61 NEO
Pseudopostega microlepta (Meyrick, 1915) Davis, 1989: 76 NEO
Opostega microlepta Meyrick, 1915a: 239
Pseudopostega mignonae Davis & Stonis, 2007: 86 NEO
Pseudopostega monosperma (Meyrick, 1931) Davis, 1989: 76 NEO
Opostega monosperma Meyrick, 1931b: 162
Pseudopostega monstruosa Davis & Stonis, 2007: 68 NEO
Pseudopostega obtusa Davis & Stonis, 2007: 91 NEO
Pseudopostega ovatula Davis & Stonis, 2007: 52 NEO
Pseudopostega paraplicatella Davis & Stonis, 2007: 82 NEO
Pseudopostega paromias (Meyrick, 1915a) Davis, 1989: 77 NEO
Opostega paromias Meyrick, 1915a: 240
Pseudopostega perdigna (Walsingham, 1914) Davis, 1989: 77 NEO
Opostega perdigna Walsingham, 1914: 349
Pseudopostega pexa (Meyrick, 1920) Davis, 1989: 77 NEO
Opostega pexa Meyrick, 1920c: 358
Pseudopostega plicatella Davis & Stonis, 2007: 82 NEO
Pseudopostega pontifex (Meyrick, 1915) Davis, 1989: 77 NEO
Opostega pontifex Meyrick, 1915a: 240
Pseudopostega protomochla (Meyrick, 1935) Davis, 1989: 77 NEO
Opostega protomochla Meyrick, 1935: 567
Pseudopostega pumila (Walsingham, 1914) Davis, 1989: 77 NEO
Opostega pumila Walsingham, 1914: 350
Pseudopostega resimafurcata Davis & Stonis, 2007: 124 NEO
Pseudopostega robusta Remeikis & Stonis in Remeikis et al., 2009: 281 NEO

Pseudopostega rotunda Davis & Stonis, 2007: 51 NEO
Pseudopostega sacculata (Meyrick, 1915) Davis, 1989: 77 NEO
Opostega sacculata Meyrick, 1915a: 240
Pseudopostega saltatrix (Walsingham, 1897) Davis, 1989: 77 NEO
Opostega saltatrix Walsingham, 1897: 140
Pseudopostega sectila Davis & Stonis, 2007: 113 NEO
Pseudopostega serrata Davis & Stonis, 2007: 52 NEO
Pseudopostega spatulata Davis & Stonis, 2007: 70 NEO
Pseudopostega sublobata Davis & Stonis, 2007: 107 NEO
Pseudopostega subtila Davis & Stonis, 2007: 88 NEO
Pseudopostega suffuscula Davis & Stonis, 2007: 139 NEO
Pseudopostega tanygnatha Davis & Stonis, 2007: 90 NEO
Pseudopostega tenuifurcata Davis & Stonis, 2007: 112 NEO
Pseudopostega triangularis Davis & Stonis, 2007: 79 NEO
Pseudopostega trinidadensis (Busck, 1910) Davis, 1989: 77 NEO
Opostega trinidadensis Busck, 1910: 245
Pseudopostega truncata Davis & Stonis, 2007: 67 NEO
Pseudopostega tucumanae Davis & Stonis, 2007: 64 NEO
Pseudopostega turquinoensis Davis & Stonis, 2007: 119 NEO
Pseudopostega uncinata Davis & Stonis, 2007: 60 NEO

Taxa excluded from Nepticuloidea

See further van Nieukerken & Johansson (1987), Davis (1989), Puplesis & Robinson (1999) and for the fossils Doorenweerd et al. (2015b).

FAMILY **ARGYRESTHIIDAE**

Argyresthia abdominalis Zeller, 1839: 205 WP
Nepticula abdominalella (Duponchel, [1845]) Bruand, 1859: 686

FAMILY **BUCCULATRICIDAE**

Bucculatrix cristatella (Zeller, 1839) Zeller, 1848: 300 WP
Lyonetia concolorella Tengström, 1848 (syn: Rebel, 1901: 220)
Nepticula concolorella (Tengström, 1848) Heydenreich, 1851: 92
Bucculatrix frangutella (Goeze, 1783) WP
Elachista rhamnifoliella Treitschke, 1833: 183
Opostega rhamnifoliella (Treitschke, 1833) Bruand, [1851]: 86
Bucculatrix centrospila (Turner, 1923) Davis, 1989: 2 AUS
Opostega centrospila Turner, 1923: 179

FAMILY **COSMOPTERIGIDAE**

Stagmatophora heydeniella (Fischer von Röslerstamm, 1841) WP
Oecophora heydeniella Fischer von Röslerstamm, 1841: 256
‡ *Opostega torquillaepennella* Bruand, [1851]: 86 NN (syn: Bruand, [1851]: 86)

FAMILY **GELECHIIDAE**

Nepticula belfrageella Chambers, 1875a: 75 [80] NO NEA

Stigmella belfrageella (Chambers, 1875) Newton & Wilkinson, 1982: 456 NO

FAMILY **GRACILLARIIDAE**

Metriochroa latifoliella (Millière, 1886) Vári, 1961: 196 WP

Nepticula latifoliella Millière, 1886: 220

Phyllocnistis saligna (Zeller, 1839) WP

Tinea cerasifoliella Hübner, 1796: pl 28: 190 NO (syn: Stainton, 1848: 2158)

Opostega saligna Zeller, 1839: 214

Opostega salicifoliella Duponchel, 1844: 377

Opostega salignatella Bruand, [1851]: 86 UE

Opostega lugdunensella Bruand, 1859: 691

Opostega cerasifoliella (Hübner, 1796) Bruand, 1859: 691

Phyllocnistis unipunctella (Stephens, 1834) WP

Argyromyges unipunctella Stephens, 1834: 260

Opostega suffusella Zeller, 1847: 894 [type species of *Phyllocnistis*]

‡ *Opostega tremulella* Fischer von Röslerstamm in Zeller, 1843: 21 NN (syn: Zeller, 1848: 266)

Opostega tremulella Heeger, 1852: 278

Phyllocnistis argentella (Bradley, 1957) Puplesis & Robinson, 1999: 18 AUS

Opostega argentella Bradley, 1957: 108

Phyllonorycter populifoliella (Treitschke, 1833) WP

Elachista populifoliella Treitschke, 1833: 188

Nepticula pilosissimella Bruand, 1859 : 686

FAMILY **HELIOZELIDAE**

Heliozela sericiella (Haworth, 1828) WP

Tinea sericiella Haworth, 1828: 585

Aechmia saltatricella Fischer von Röslerstamm, 1841: 249

Nepticula saltatricella (Fischer von Röslerstamm, 1841) Bruand, 1859: 687

Heliozela lithargyrellum (Zeller, 1850) WP

Tinagma lithargyrellum Zeller, 1850: 158

Nepticula lithargyrella (Zeller, 1850) Bruand, 1859: 687

FAMILY **LYONETIIDAE**

Leucoptera malifoliella (O. Costa, 1836) WP

Elachista malifoliella O. Costa, 1836: [239] Elachista 3

Opostega scitella Zeller, 1839: 214 (syn: Stainton, 1869a: 269)

Leucoptera sinuella (Reutti, 1853) WP

Cemiostoma sinuella Reutti, 1853: 208

Cerniostoma susinella Herrich-Schäffer, 1855: 342

Opostega susinella (Herrich-Schäffer, 1855) Bruand, 1859: 691

Leucoptera spartifoliella (Hübner, 1813) Hübner, 1825: 426 WP
Tinea spartifoliella Hübner, 1813: 49
Opostega spartifoliella (Hübner, 1813) Zeller, 1839: 214
Leucoptera phaeopasta (Turner, 1923) Davis, 1989: 2 AUS
Opostega phaeopasta Turner, 1923: 180
Lyonetia clerkella (Linnaeus, 1758) WP,EP
Phalaena clerkella Linnaeus, 1758: 542
Opostega magnimaculella Bruand, 1859: 691 (syn: Leraut, 1997: 101)
Lyonetia leucoprepes (Bradley, 1961) Davis, 1989: 2 AUS
Opostega leucoprepes Bradley, 1961: 160
Petasobathra ischnophaea (Meyrick, 1930) Davis, 1989: 2 OR
Opostega ischnophaea Meyrick, 1930: 7

FAMILY **TISCHERIIDAE**

Coptotriche angusticollella (Duponchel, 1843) Diškus & Puplesis, 2003: 430 WP,EP
Nepticula suberoidella Walsingham, 1891: 152 (syn: Diškus & Puplesis, 2003: 430)
Stigmella suberoidella (Walsingham, 1891) Le Marchand, 1946b: 284
(see van Nieukerken, 2004: 112)

FAMILY **UNKNOWN**

Tinea minimella O.G. Costa, 1836: [230] Tinea 18, JH of *Tinea minimella* [Denis & Schiffermüller], 1775, now *Nemophora minimella* (Adelidae) WP
Nepticula minimella (O.G. Costa, 1836) Stainton, 1869a: 267

FAMILY **UNKNOWN, may be Trichoptera, Hydroptilidae**

Tinea commatella Schrank, 1802: 133 ND WP
Nepticula commatella (Schrank, 1802) Stainton et al., 1855: 264

UNPLACED FOSSILS

Tinea araliae Fritsch, 1882: 6 [may be Gracillariidae] WP†
Stigmellites araliae (Fritsch, 1882) Kozlov, 1988: 30
Foliofossor cranei Jarzembowski, 1989: 448 WP†
Troponoma curvitracta Krassilov, 2008: 101 WP†
Troponoma festunata Krassilov, 2008: 102 WP†

Notes

1 The genus *Manoneura* Davis, 1979 had been synonymised with *Enteucha* Meyrick, 1915 by van Nieukerken (1986a), but resurrected as separate genus by Puplesis and Robinson (2000) on the basis of its aberrant genitalia. In two independent molecular analyses, the type species *Manoneura basidactyla* (Davis, 1978) clearly

groups inside the genus *Enteucha* (Regier et al. 2015, Doorenweerd et al. 2016a) and we thus regard it here again as synonym.

2 The paper by Walsingham on the Tenerife fauna (Walsingham 1908a), where also the validity of the name *Stigmella* was established, has often been cited as Walsingham, 1907. This paper is published in the last issue of the Entomologist's Monthly Magazine for 1907, but issued on June, 4th, 1908 as can be seen on the wrappers of volume 1908 (1), page iv (http://biodiversitylibrary.org/page/31209657) (see also Sattler 1973), thus the citations should be Walsingham 1908.

3 The genus *Stigmella* is divided into two large clades (Doorenweerd et al. 2016a) that are termed respectively "core *Stigmella*" for the clade containing the type species of both *Stigmella* and *Nepticula* (*S. anomalella* and *S. aurella*) and non-core *Stigmella* for the other clade (containing the type species of *Astigmella*: *S. naturnella*). We refrain from recognising different genera, since recognising these clades morphologically is not always possible. Several species groups could not be placed due to lack of molecular information, they are listed at the end of *Stigmella*.

4 *Stigmella resplendensella* (Chambers, 1875) was placed as species with uncertain affinities due to the lacking abdomen in the lectotype (Newton and Wilkinson 1982). We re-examined the lectotype and could match the characteristic metallic forewing colour pattern to two males and one female we had on loan, of which the female had been barcoded (see also BugGuide: http://bugguide.net/node/view/391014/bgpage). Barcode and genitalia confirm that *S. resplendensella* is closely related to *S. unifasciella* (Chambers, 1875), as earlier suggested by Braun (1917). Further details to be published elsewhere. Material: Lectotype female, MCZ-ENT00014954 (designated by Newton & Wilkinson, 1982: 456, [USA: Kentucky, Covington], captured May 23rd, under hackberry trees (Celtis occidentalis), V.T. Chambers, Type14954 [head and abdomen missing].

5 We prefer the name *prunifoliella* group rather than the older "*prunetorum* group", since the *prunifoliella* group contains several North American species, whereas the name "*prunetorum* group" was based on a single European species only.

6 The date of publication of "Die Schmetterlinge Deutschlands und der Schweiz. Zweite Abtheilung. Kleinschmetterlinge. Band 2. Die Motten und Federmotten. Heft 2" has previously often been cited as 1877, the date that also figures on the Title page (https://archive.org/details/dieschmetterlin01heingoog). However, already Kirby concluded in the Zoological Record 13 (published 1878) on page 187: "Band ii, Die Motten ..... . though bearing date 1877, was published not later than November, 1876" (see also Sattler 1973). We therefore cite the paper as Heinemann and Wocke ([1876]).

7 *Stigmella acerna* Puplesis, 1988. We synonymise the unavailable name *Stigmella acerifoliella* Dovnar-Zapolski, 1969, which was collected from *Acer turcomanicum* in Turmenistan, Kopet-Dag, the same type locality and host as for *S. acerna* and with a similar mine form. The paper by Dovnar-Zapolski (1969) is poorly known, but contains a number of new names for Nepticulidae and other leafminers, all

based on the mine alone. They are therefore not available (ICZN art. 13.6.2), and in many cases it is impossible to determine the identity of these names with the little information provided, but they do provide some interesting records.

8 *Stigmella ulmivora* (Fologne, 1860). Hofmann (1858) named a species *Nepticula ulmella* HS. from Regensburg. This name should be regarded as an unavailable name (*nomen nudum*), since there is no description nor indication. This name is also a *nomen oblitum*, never cited again until Segerer (1997) synonymised it with *S. ulmivora*. Thus no further action needs to be taken to reverse precedence to avoid rejecting the junior synonym *S. ulmivora* or junior homonym *Nepticula ulmella* Braun, 1917 (now *Ectoedemia ulmella*).

9 Stonis and Rocienė (2014) placed *Stigmella multispicata* Rocienė & Stonis, 2014 in the *Stigmella malella* group (that previously also contained the species of our *S. rhamnella* group and in their vision also the *prunifoliella* group), but in fact the species is in all aspects extremely similar to *Stigmella ulmivora*, that only can be separated by details in the genitalia. It is therefore moved here, and it is highly likely that it feeds on *Ulmus*.

10 *Stigmella palionisi* Puplesis, 1984. We synonymise *Stigmella nakamurai* Kemperman & Wilkinson, 1985 from Japan with *S. palionisi* from Russia: Primorye, on the basis of a comparison of descriptions, a male paratype slide of *S. nakamurai* (Fig. 41, 42) and detailed photos of *S. palionisi* genitalia (Rocienė and Stonis 2013; Stonis and Rocienė 2013) (also Fig. 43). This also confirms *Ulmus* as host for *S. palionisi*. This species is also morphologically very similar to European *S. viscerella* (Stainton, 1853), but the mines are somewhat different. Material: 1♂, Paratype *S. nakamurai*, Hokkaido, Sapporo, em. 20.viii.1981, S. Nakamura, Host 0360 Ulmus davidiana v. japonica, slide VU no. 0790 (collection Sapporo).

11 *Stigmella macrocarpae* (Freeman, 1967). We use the junior name for this North American oak mining species, earlier known as *S. latifasciella* (Chambers, 1878), because its original combination *Nepticula latifasciella* is a junior primary homonym of *Nepticula latifasciella* Herrich-Schäffer, 1855, a junior synonym of *Stigmella hybnerella* (Hübner, 1796).

12 *Stigmella birgittae* Gustafsson, 1985. We place the unavailable name *Nepticula amseli* Skala, 1941 as synonym under *S. birgittae*. *N. amseli* was described from mines from *Zizyphus spina-christi* in Jericho, Palestine. *Stigmella birgittae* is a common and widespread species in the Middle East on this host (van Nieukerken 2010).

13 *Stigmella abaiella* Klimesch, 1979 and *S. ficulnea* Puplesis & Krasnilnikova, 1994. The illustrated genitalia of both species are extremely similar. Study of the holotype and some paratypes of *S. abaiella* confirmed this (Figs 44–47). It is possible that this represents a single species, but we hesitate to synonymize them, since *S. abaiella* allegedly was reared from *Pyrus* and *S. ficulnea* clearly is a *Ficus* miner. Since these hosts are totally unrelated, it is not very likely that one species feeds on both. Dr. Mansour Abaii (Teheran, e-mail 26.xii.2015 to EvN) confirmed that the identification of *Pyrus* must be correct, but no mines are kept. Awaiting

Figures 41–47. *Stigmella* species. **41–43** *S. palionisi* Puplesis, 1984 **41, 42** Male genitalia of paratype of *Stigmella nakamurai* Kemperman & Wilkinson, Japan, slide VU 0790 **43** Male genitalia of specimen from Russia, Primorye, slide JCK8111. **44–47** *Stigmella abaiella* Klimesch, 1979 **44** Holotype male with labels **45, 46** Male genitalia of holotype, slide EvN4759 **47** Female genitalia, paratype, slide Klimesch 866.

further information from freshly collected larvae, we keep the species separate, but a synonymy is not excluded.
Material: Holotype ♂ *S. abaiella*, Iran, Teheran, 20.ix.1978, Abai, *Pyrus communis*, Genitalia slide EvN4759 (Staatliches Museum für Naturkunde Karlsruhe).

14 *Stigmella paliurella* Gerasimov, 1937. Previously the author for this species was given as (Klimesch, 1941), because Gerasimov (1937) based the name on the leafmine (van Nieukerken 1986a), following the rule that names based on the work of an animal published after 1930 are excluded from zoological nomenclature (ICZN art. 13.6.2). However, since Gerasimov also describes characters of the larva, the description fulfils the code (article 13.1), despite the brief description (van Nieukerken 2013) and the name is available and valid.

15 *Stigmella turbatrix* Puplesis, 1994. We synonymise the unavailable name *Stigmella celtivora* Dovnar-Zapolski, 1969, which was described from leafmines on *Celtis* in Kazakhstan, Alma-Ata. See also note 4.

16 *Stigmella microtheriella* (Stainton, 1854) was recorded from China (van Nieukerken and Liu 2000) and Japan (Hirano 2013). During our recent fieldwork in Korea and Japan we found larvae on *Carpinus*, *Corylus* and *Betula* (Hokkaido: RMNH. INS.29748; new host record, Fig. 56) (in Japan also found on *Ostrya*, N. Hirano pers. comm.) of which the DNA barcode shows a very short distance (1.11%) to European *S. microtheriella*, still clustering together (BIN Korea/Japan: BOLD:ACU7085, Europe: BOLD:AAI0007) (see Fig. 57). We reared both males and females from these Asian specimens, whereas in most of Europe *S. microtheriella* is parthenogenetic (but we reared males from Greece and Laštůvka and Laštůvka 1997 recorded males from Croatia). Originally we considered the presence of *S. microtheriella* in East Asia as an introduction, but the separate, but similar barcode, suggests the species is an indigenous element of the East Asian fauna. By comparing male genitalia to the type of *S. cathepostis* Kemperman & Wilkinson, 1985 (Figs 48–50), we realised that the latter is nothing else than male *S. microtheriella* and hence synonymise it here, even though the uncus of east Palearctic specimens seems to have a deeper indentation than European males (compare Figs 48–53 with 54 and 55). *S. cathepostis* was also recorded from Russia: Primorye (Rocienė and Stonis 2013), and we here also illustrate a male from Primorye (Fig. 53). It is unclear whether *Stigmella microtheriella* is a trans Palearctic species, since its best known hosts, *Corylus, Carpinus* and *Ostrya* do not have a continuous distribution, but show a gap between the Urals and the middle of China (Sokolov et al. 1980; Fang et al. 2011). A continuous distribution, however, might still be a possibility, since apparently *Betula*, that occurs throughout Siberia, can be an alternative host (see above and Fig. 56). Recently, *S. microtheriella* was also recorded from North America (Eiseman and van Nieukerken 2015), but DNA barcodes show that here indeed it is an introduced species, as it is in New Zealand (Donner and Wilkinson 1989).
Material: Holotype ♂: [Japan, Kyushu], Hikosan, Buzen, 30.vii.1954, H. Kuroko, Host: *Carpinus tschonoskii* Maxim., genitalia slide VU no. 0783 (Entomological Laboratory, University of Osaka Prefecture).



Figures 48–56 *Stigmella microtheriella*, male genitalia and leafmine (56). **48–50** Holotype of *Stigmella cathepostis* Kemperman & Wilkinson, 1985, slide VU0783 **51–52** slide EvN4629, Korea (Jeollanam-do) Wando Island, Hwaheung-ri, from *Carpinus tschonoskii*, slide EvN4629 **53** Russia, Primorye, slide JCK8117 **54–55** Greece (Messinía) Taygetos Mts, from *Ostrya carpinifolia*, slide EvN4430 **56** larva in leafmine on *Betula platyphylla*, Japan, Hokkaido, RMNH.INS.29748.



Figure 57. Neighbor Joining tree of DNA Barcodes of Palearctic members of the *Stigmella betulicola* group. East Palearctic records are presented in red, West Palearctic and imported North American records in black. Barcode Identification numbers are given at the right.

17 *Stigmella nivenburgensis* (Preissecker, 1942). We here synonymise *Stigmella populnea* Kemperman & Wilkinson, 1985. We found *Stigmella nivenburgensis* commonly on *Salix* in South Korea and China and DNA barcodes (BOLD:ACU6572) are rather close to European specimens (BOLD:AAV7002) (see Fig. 57). Although we did not yet collect mines on *Salix* in Japan, we consider it more likely that the mine on *Populus* from which the single specimen of *S. populnea* was reared is *S. nivenburgensis*, rather than a separate species, as in many other leafminers that feed both on *Salix* and *Populus*. The female genitalia are not different from European specimens, but are not very diagnostic in this group. The male genitalia figured as *Stigmella betulicola* from Russia: Primorye (Rocienė and Stonis 2013) belong in our opinion in fact also to *S. nivenburgensis*.

18 *Stigmella cornuta* Rocienė & Stonis, 2013. The authors of this Chinese *Quercus* feeding species erected a separate species group for it (Stonis et al. 2013e), but we think that the morphology fits well in the somewhat enlarged *betulicola* group as we define it here. Also *S. xystodes* has relatively large cornuti, comparable to *S. cornuta*.

19 *Stigmella caryaefoliella* (Clemens, 1861). Wilkinson and Scoble (1979) synonymized *S. caryaefoliella* and *S. obscurella* (Braun, 1912) (see note 17) with *ostryaefoliella* on the basis of the male genitalia. We remove these synonyms here and recognize three good species feeding on different hosts and having quite different DNA barcodes and morphologies. Details to be published elsewhere.

20 *Stigmella myricafoliella* (Busck, 1900) was described from Florida, Palm Beach, and reared from leafmines on *Morella cerifera* (as *Myrica cerifera*). We revise the synonymy of *N. obscurella* Braun, 1912 as a synonym of *S. myricafoliella*, on the basis of the host plant and genitalia. Details to be published elsewhere.

21 *Stigmella ostryaefoliella* (Clemens, 1861). Wilkinson and Scoble (1979) synonymized *S. caryaefoliella* (Clemens, 1861) and *S. obscurella* (Braun, 1912) (see notes 16 and 17) with *ostryaefoliella* on the basis of the male genitalia. We remove these synonyms here and recognize three good species feeding on different hosts and having quite different DNA barcodes and morphologies. Details to be published elsewhere.

22 *Stigmella pelanodes* (Meyrick, 1920). The holotype was examined by EvN (Figs 58–61): the genitalia are rather similar to *Stigmella xystodes*, we therefore place this species in the *betulicola*-group, together with several other African species that had been placed there before (Diškus and Puplesis 2003).
Material: Holotype ♂: [Kenya, Kwale] *Côte d'Afrique or. angl.*, Rivière RAMISI, ALLUAUD ET JEANNEL, Nov. 1911, St. 8;TYPE; "M.647"[Meyrick's hand]; MUSÉUM PARIS, Ch. ALUAUD & R. JEANNEL, 1914; Nepticula pelanodes Meyr. type; RMNH Lepidoptera Genitalia Slide EvN4795 ♂ (MNHN).

23 *Stigmella tropicatella* Legrand, 1965 is the only nepticulid species known from the islands in the Indian Ocean off the African continent (Madagascar excluded) as found on the atoll of Aldabra. The holotype, unfortunately, is almost completely destroyed, only a head and part of the thorax remain on the minuten pin. There are

Figures 58–62. African *Stigmella* species. **58–61** *S. pelanodes* (Meyrick, 1920), male holotype, labels and genitalia, slide EvN4795. **62** *Stigmella tropicatella* Legrand, 1965, male paratype and labels.

several paratypes, three of which EvN examined. Two are males of a *Stigmella*, the third is a male of an unnamed *Acalyptris*. Comparing the remains of the holotype and the description, we are convinced that the *Stigmella* paratypes are the real *S. tropicatella*, and hence base the identity on these (Figs 62–65). It is a typical tropical



Figures 63–69. *Stigmella tropicatella* Legrand, 1965, paratypes. **63–65** Male genitalia of male in Fig. 62, slide EvN4794, on which identity of species is based **66–69** *Acalyptris* sp., misidentified paratype of *S. tropicatella*, male, labels and male genitalia, slide EvN4846.

Stigmella species in the *betulicola* group, rather similar to *S. satarensis* Scoble, 1978. The unnamed *Acalyptris* is illustrated in Figs 66–69.

Material: Adult Holotype, [specimen almost completly disappeared, only a head on pin, [Seychelles], Aldabra, 11.xi.1959, M. Gerber (Muséum National d'Histoire Naturelle, Paris). Further specimens see S1.



Figures 70–75. *Stigmella wollofella* (Gustafsson, 1972), male genitalia. **70–71** *Nepticula mandingella* Gustafsson, 1972, holotype, slide NHRS4874 **72–75** *Nepticula wollofella* Gustafsson, 1972, holotype, slide NHRS5103.

24 *Stigmella wollofella* (Gustafsson, 1972). Study of the slides of the holotypes of *Nepticula wollofella* Gustafsson, 1972 (Figs 72–75) and *N. mandingella* Gustafsson, 1972 (Figs 70–71), both collected together in Gambia, showed that the genitalia are in fact identical, but the slide of *S. mandingella* is more squashed, obscuring some of the characters. Gustafsson (1985) gave the shape of the juxta as distinguishing character. The juxta is arrow shaped, with many small spines on the lateral apexes of the arrow head. These were not figured in the original drawing of *N. mandingella*, but are present in the slide, although difficult to separate from spines on the phallus tube. Because both *N. mandingella* and *N. wollofella* were described in the same publication, we determine here as first reviser the relative priority of *N. wollofella* (ICZN article 24.2) and synonymise *N. mandingella*. The much better genitalia preparation of *N. wollofella* was the first criterion and the second one is that the name *wollofella* has been used again by Gustafsson (1985) when describing the biology of the species as leafminer of *Zizyphus*.

Material: Holotype ♂ *Nepticula wollofella* Gustafsson, Gambia: Gambia river between Bathurst and Basse Santa Su., on riverboat M/S Lady Wright in flash light, 5.xii.1970, B. Gustafsson Genitalia slide 5103 (NHRS). Holotype ♂ *Nepticula mandingella* Gustafsson, Gambia: Gambia river between Bathurst and Basse Santa Su., on riverboat M/S Lady Wright in flash light, 5.xii.1970, B. Gustafsson Genitalia slide 4874 (NHRS).

25 *Stigmella rosaefoliella* (Clemens, 1862). The subspecies *S. rosaefoliella pectocatena* Wilkinson & Scoble, 1979 is removed as a synonym to *S. centifoliella*, see there.

26 *Stigmella ogygia* group. Since our studies show that all New Zealand *Stigmella* species – as far as studied - belong to one monophyletic clade, we group them as the *Stigmella ogygia* group, based on the widespread and common *S. ogygia* (Meyrick, 1889), which is readily recognizable on host-plant alone (herbaceous *Senecio*, Asteraceae).

27 *Stigmella epicosma* group. Puplesis and Robinson (2000) placed many Neotropical species in an enlarged *salicis* group because of the similarity in male genitalia. Our phylogeny (Doorenweerd et al. 2016a) gives the two examined Neotropical species of this group (both Asteraceae feeders) as sister to the *salicis* group s.str., whereas the only Neotropical *Salix* feeder *S. molinensis* van Nieukerken & Snyers, 2016 is sister to the remaining Holarctic *salicis* group members. Because of the strong difference in host plant choice, all *salicis* group members but one feed on Salicaceae, whereas the Neotropic species feed on Asteraceae and various other families, and because of the strong apomorphy of the signa band in the female genitalia for the *salicis* group, we separate all Neotropic species (apart from *molinensis*) and place them in their own *epicosma* group.

28 *Stigmella costalimai* (Bourquin, 1962) and *S. guittonae* (Bourquin, 1962) are tentatively moved to the *epicosma* group because their host plants belong to Asteraceae, resp. *Tessaria integrifolia* Ruiz & Pav. and *Senecio bonariensis* Hook. & Arn. (Bourquin 1961) and so far all known Neotropic Asteraceae feeders belong to this group. Puplesis and Robinson (2000) overlooked the original host plant data for *S. costalimai*. The record of *Ludwigia longifolia* (DC.) H.Hara (Onagraceae) as one of the host plants for *S. guittonae* by Bourquin (as *Jussiaea longifolia*) is most likely an error: just after this description he wrote the description of a Momphid, *Psacophora orfilai*, making leafmines on that host. Momphidae frequently feed on Onagraceae.

29 Species described by Klimesch in his paper "Neue *Stigmella*-Arten (Lep., Stigmellidae)" in the "Zeitschrift der Wiener Entomologischen Gesellschaft vol 31(9–12) have been incorrectly cited by us and others as Klimesch, 1946 (van Nieukerken 1986a; 2013; Diškus and Puplesis 2003). However, this issue, 9–12 of volume 31, for the year 1946 was published on 15 March 1948 as can be seen on the first page of that issue, 129 (http://www.zobodat.at/pdf/ZOEV_31_0129.pdf) ("Ausgegeben 15.März 1948").

30 *Stigmella salicis* (Stainton, 1854) forms a complex of species that requires revision: some synonyms may be one of the constituent species, others have to be described as new (van Nieukerken et al. 2012a).

31 *Stigmella centifoliella* (Zeller, 1848). *Nepticula centifoliella* was first named by Heyden (1843) when describing *Nepticula*. Since this was a short message during a meeting, there is no description, and only a vague indication to unspecified earlier papers ("Von einer Art (der *Centifoliella*) kannten schon de Geer und Goeze die eigenthümlich gebildete Raupe"). We consider this insufficient to make the name available, and thus keep Zeller, 1848, who described the species in detail, as author (see also Wilkinson 1978). We synonymise *Stigmella rosaefoliella pectocatena* Wilkinson & Scoble, 1979 from Canada with *S. centifoliella*. The authors overlooked the fact that this represents in fact the introduced European species, which is rather different from North American *S. rosaefoliella*, that only externally superficially resembles it and has very different genitalia. Details will be published elsewhere.
Material: Holotype ♀ *Stigmella rosaefoliella pectocatena* Wilkinson & Scoble: Canada, Ontario, Ottawa, emerged 24.ix.1962, leafmines on *Rosa* sp., 62-11, Freeman & Lewis, genitalia slide CNC3292 [not found] (Canadian National Collection of Insects, Arachnids, and Nematodes).

32 *Stigmella azaroli* (Klimesch, 1978) is very similar to *S. perpygmaeella*, but the moth is usually much paler. DNA barcodes of *S. perpygmaeella*, *incognitella* and *azaroli* form a tangled cluster, where *S. perpygmaeella* is paraphyletic, and island forms have very different barcodes from continental populations (Fig. 76). In this unclear situation we rather do not synonymise *S. azaroli*, until more reared material and DNA analyses are available (BIN's: *S. perpygmaeella*, most of Europe: BOLD:AAI0008, Greece, Holotype *S. azaroli*: BOLD:ACG8759, *S. azaroli* Cyprus: BOLD:ACU6096, *S. incognitella*: BOLD:AAF3364, "*S. perpygmaeella*" Malta: BOLD:ACG8760, Mallorca: BOLD:ACG8761).
Material: Holotype ♂ *Nepticula azaroli*: Greece, Rodos, Rodini, 22.ix.1972, mines on *Crataegus azarolus*, em. 11.x.1972, J. Klimesch, Genitalia slide EvN4235, DNA barcode RMNH.INS.24235 (Zoologische Staatssammlung München).

33 *Stigmella magdalenae* (Klimesch, 1950). This species was known as *Stigmella nylandriella* or *Nepticula nylandriella* in most literature prior to Borkowski (1975) and in Britain prior to Emmet (1979). See Schoorl et al. (1985).

34 *Stigmella nylandriella* (Tengström, 1848). Before the type of *Lyonetia nylandriella* was re-examined in the 1970's, this species was known as *S. aucupariae* (Frey, 1857) and *Stigmella* or *Nepticula nylandriella* referred to the species now known as *S. magdalenae* (Borkowski 1975; Schoorl et al. 1985).

35 *Stigmella oxyacanthella* (Stainton, 1854). The finding of larvae on *Crataegus*, *Malus* and *Amelanchier* in North America apparently having the same DNA barcode as European *S. oxyacanthella* (BOLD:AAF3421) led EvN to re-examine specimens identified as *S. pomivorella* (Packard, 1870) in CNC and USNM. All show genitalia inseparable from *S. oxyacanthella*, and we thus synonymise *Stigmella pomivorella*. This leafminer apparently was already introduced in the USA during the 19th century. Details will be published elsewhere.

36 *Stigmella micromelis* Puplesis, 1985. After the finding that Siberian larvae on *Crataegus* share the barcode with Korean and Japanese larvae found on *Aria alnifolia*



Figure 76. Neighbor Joining tree of DNA Barcodes of the *Stigmella incognitella* group, showing the paraphyly with regards to *S. azaroli* and *S. incognitella* and large distances for specimens identified as *Stigmella perpygmaeella* from Malta and Mallorca. Specimens from Cyprus are identified as *S. azaroli*, of which the holotype from Rhodos was also barcoded.

(= *Sorbus alnifolia* or *Micromelis alnifolia*) (BOLD:ACK9547), and the fact that genitalia of *S. micromelis* and *S. crataegivora* Puplesis, 1985 are indistinguishable, we synonymise both species here. Since both species were named in the same publication, we here determine as First Reviser (ICZN art 24.2) that *S. micromelis* has priority over *crataegivora*. Photos of the genitalia of the types were given by Stonis and Rocienė (2013). A paper with more details is in preparation.

37 *Stigmella amelanchierella* (Clemens, 1862), **stat. rev**. *Stigmella amelanchierella* was described on the basis of leafmines only. Subsequent authors have been unable to rear or to identify the species and Newton and Wilkinson (1982) were even uncertain about its generic status. We found several times leafmines with green larvae on *Amelanchier*, that fit Clemens' description well. DNA barcodes of these showed two different clusters, both closely related to other members of the *oxyacanthella* group, such as *S. crataegifoliella*, one occurring more west in Colorado and Can-

ada (BOLD:ACG5879), another east in Tennessee, Virginia and Massachusetts (BOLD:ACG8835). We consider it likely that the last group represents the real *S. amelanchierella*, since Clemens described it from Pennsylvania. Even though we have not yet studied a single adult, we consider it likely to be a separate species in the *S. oxyacanthella* group, of which the identity is based on the DNA barcode.

38 *Stigmella purpuratella* (Braun, 1917). After study of the holotype slides we synonymise *S. scinanella* Wilkinson & Scoble, 1979 with *S. purpuratella*. Already Newton and Wilkinson (1982) suggested both species cannot be distinguished, except by the paler and less iridescent forewings of *S. purpuratella*. This difference can easily be explained by the much older age of the *purpuratella* specimens, but this text did not seem to be supported by the very different drawings of the genitalia of both species in the same paper. After checking the slides it can be concluded that the valvae of *S. scinanella* were drawn incorrectly, and the drawing of the phallus of *S. purpuratella*, which is a reconstruction of a broken phallus, is turned upside down, making the characteristic longer cornuti pointing posteriorly rather than anteriorly, and the phallus tube is an incorrect reconstruction. The species makes leafmines on *Crataegus* that are inseparable from those of *S. crataegifoliella*. Figures will be published elsewhere.
Material: Holotype ♂ *Nepticula purpuratella*: (United States), Pennsylvania, Pittsburgh, 30.v.1906, Engel, Genitalia slide CNC3467 P. Newton (USNM).
Holotype ♂ *Stigmella scinanella*: [Canada], Ontario, Normandale, mines 26.vii.1956 on *Malus*, 56-154, Freeman & Lewis, Genitalia slide MIC7102 (=CNC2972) (Canadian National Collection of Insects, Arachnids, and Nematodes).

39 *Stigmella aurella/ruficapitella* cluster. The remaining *Stigmella* species form a well supported clade in our multi-gene molecular analyses (Doorenweerd et al. 2016a; Doorenweerd et al. 2016b), that is also relatively well recognisable by genitalia characters, such as the presence of a manica (phallocrypt) in the male genitalia and a large accessory sac in the female genitalia. However, within this cluster the relationships are confusing and many unsupported clades appear in various analyses, often at different places. Some supported groups stand out, including the complete *aurella* group (including *S. lediella*), the *S. floslactella* group and the core *S. ruficapitella* group s.s. Also part of the *lemniscella* group (formerly *marginicolella* group) is well supported: viz. *S. continuella, S. lemniscella* + *S. zelkoviella*, usually with poorer support linked to *S.apicialbella* and an unnamed *Betula* feeding North American species, but always never with *S. gimmonella* included (only one specimen sequenced). The other Fagaceae mining species, including the North American *S. procrastinella* and *S. alba* almost always form a poorly supported clade together with the core *ruficapitella* group, which we therefore together classify as the *S. ruficapitella* group s.l., that is here restricted to Fagaceae feeding species only. *Stigmella speciosa*, *lonicerarum* and relatives, previously placed with the *ruficapitella* group, always group outside. A *Stigmella sorbi* group is never recognised in molecular analyses and even the morphologically similar *S. sorbi* and *S. plagicolella* are never placed close to each other. On the other hand, *S. amygdali*, previously placed

in a monotypic species group, always groups with moderate support together with *S. plagicolella*, forming a *Prunus* feeding clade. This situation can currently not be easily translated into monophyletic species groups, and apart from the well supported ones, we recognise tentatively a *S. speciosa* group for the non-Fagaceae miners previously placed in the *ruficapitella* group (or the *hemargyrella* group) and an enlarged *sorbi* group, at the moment a kind of waste bin for mainly Rosaceae feeding species, but also including *S. hamamelella* on *Hamamelis*. We leave *S. zagulaevi* from the Caucasus and *S. talassica* from the Tyan Shan in the *lemniscella* group, although we are not convinced that this is the correct placement.

40 *Stigmella monticulella* Puplesis, 1984. We synonymise here *Stigmella gracilipae* Hirano, 2014 from Japan. Both taxa make linear leafmines on *Lonicera*, occur in East Asia and have inseparable genitalia (Stonis and Rocienė 2013; Hirano 2014). The species is also close to *S. lonicerarum* (Frey, 1857) from Europe.

41 *Stigmella filipendulae* (Wocke, 1871). After *S. ulmariae* (Wocke, 1879) had earlier been synonymized (van Nieukerken et al. 2012a), we also synonymise *S. palmatae* Puplesis, 1984 from Russia: Primorye, feeding on *Filipendula palmatae*, on the basis of the similar genitalia (Stonis and Rocienė 2013). *S. filipendulae* is apparently a widespread Palearctic *Filipendula* feeding species.

42 *Stigmella lediella* (Schleich, 1867). Puplesis (1994) already synonymized the two *Rhododendron* feeding species from Russia: Primorye, and here we also add the Japanese *Rhododendron* feeder to the synonymy: *Stigmella sesplicata* Kemperman & Wilkinson, 1985. The original description only dealt with females, but meanwhile male genitalia have been described (Hirano 2013) and are identical to *S. lediella*. We have no DNA barcodes yet from Japan, but those from Korea are almost identical to those from Europe and have the same BIN (BOLD:AAL6954). *Stigmella rhododendrifolia* Dovnar-Zapolski & Tomilova, 1978 is an unavailable name, based on leafmines from Siberia. Considering the fact that all *Rhododendron* leafmines belonging to *Stigmella* from eastern Europe to Japan apparently are all *S. lediella*, we conclude that this name also belongs here.

43 *Stigmella longispina* Puplesis, 1994 is here moved to the *aurella* group on the basis of similarity in its male genitalia as shown in a single studied specimen from Tajikistan (RMNH.INS.15381) (Figs 77–79).

44 *Stigmella spiculifera* Kemperman & Wilkinson, 1985. *Stigmella oa* Kemperman & Wilkinson, 1985 was described on the basis of a single female from Japan. After carefully comparing descriptions and figures, we can only conclude that it is indistinguishable from the *Rubus* feeding *S. spiculifera* and hence synonymize it here.

45 *Stigmella lurida* Puplesis, 1994. On the basis of DNA analysis of one female from Altai, that consistently groups with *Stigmella sorbi*, we place *S. lurida* in the *S. sorbi* group (RMNH.INS.24818).

46 *Stigmella humboldti* Remeikis & Stonis, 2015 was described from a single female reared from *Quercus humboldti* in Colombia. The authors could not place this species in a known group, due to the spiny signa on the female bursa. Since we see some resemblance with the spiny signa of *S. procrastinella* and *S. alba*, both shown

Figures 77–79. *Stigmella longispina* Puplesis, 1994, male genitalia, Tajikistan, 30 km N Dushanbe, Kondara, 20.viii.1989, R. Puplesis, slide JCK8318, RMNH.INS.15381.

to be *Quercus* feeders as well (DNA barcode data), we place *S. humboldti* close to these species.

47 *Stigmella samiatella* (Zeller, 1839). We list here two previously overlooked Herrich-Schäffer names: *Nepticula quercella* Herrich-Schäffer, 1863a was merely published as a name in a checklist (abbreviated as *Nepticula querc.*: Herrich-Schäffer followed the rule of fixed endings, -ella for Tineina, so that everybody knew this to be written as *quercella*). This unavailable nomen nudum was synonymised by Segerer (1997). *Nepticula chaoniella* Herrich-Schäffer, 1863b was described in a paper on the Lepidoptera of Engadin, in comparison with another new species, *N. alpinella* (see note 67). Herrich-Schäffer wrote: "Durch letzteres Merkmal unterscheidet sie sich auf den ersten Blick von der eichenbewohnenden *chaoniella* m. (früher unter *samiatella*, aber ohne verdickte Schuppen der männlichen Hfl.)" [By the last character it is separated on first sight from the oak feeding *chaoniella* m[ihi] (earlier under *samiatella*, but without thicker scales in the male hindwing)]. Like many early authors Herrich-Schäffer mixed several oak mining *Stigmella* species and separates his *chaoniella* from male *atricapitella* or *ruficapitella* [named by him *samiatella*], that both show these thicker (androconial) scales. From this it is clear that *chaoniella* can only be a synonym of *S. samiatella*. Further the name *N. chaoniella* can be considered a nomen oblitum, we are not aware of any later use of this name.

48 *Bohemannia pulverosella* (Stainton, 1849). We synonymise *Bohemannia piotra* Puplesis, 1984 from Russia: Primorye, because it is indistinguishable in genitalia from *pulverosella*, and makes similar mines on *Malus* as European *pulverosella*. We expect that the species has a continuous distribution throughout Siberia. In Europe most populations of *B. pulverosella* seem to be parthenogenetic without any males, and only a few males are known (van Nieukerken and Johansson 1990).

49 *Bohemannia quadrimaculella* (Boheman, 1853). The publication year for the original paper has been cited as 1851, 1852 or 1853. Boheman's paper was published in the volume for the year 1851 that according to the title page was published in 1853. On the title page of the article is printed: "Inlemnad den 6 Mars 1852 [submitted on March 6, 1852], probably causing the incorrect references to 1852. The synonymy of *Bucculatrix antispilella* Meess, 1907 established by van Nieukerken and Johansson (1990), who incorrectly cited 1910 as year of description, in fact was already spotted shortly after its description by Disqué (1912) (see Hausenblas 2009).

50 *Bohemannia manschurella* Puplesis, 1984. We synonymize here *Bohemannia nipponicella* Hirano, 2010 from Japan with *B. manschurella* from Russia: Primorye. Hirano (2010) described *B. nipponicella* as different in the cornuti, comparing it to the original description of *B. manschurella*, but by comparing the photos of the types of both species (Hirano 2010; Stonis and Rocienė 2013) and the study of material from Russia and Japan, no difference can be seen at all.

51 *Glaucolepis oishiella* (Matsumura, 1931), **comb. n.** The female holotype of *Trifurcula oishiella* Matsumura 1931 was examined, and compared to the description of *Sinopticula sinica* Yang 1989; both species were described as gall maker on *Prunus*. Further, material collected in China and Japan helped to establish that this concerns a single species, with gall making larvae on *Prunus*. Further details will be published elsewhere.

Translation of the original description in Japanese by Matsumura:

"Head orange yellow, setae on both sides grey-white. Antenna grey, base of several segments with dark rings. Body and forewing dark brown with a little purple colour ribbon and also with reflection. Scales large, tip of wing spoon-like with long hairscales. Hindwing dark. Legs brown. Hindtibia dorsally with half erect setae. Larvae on *Prunus*, boring into branches, making a gall. In mid-June to late-June in Japan, Honshu". [*Translated by Liu Youqiao, August 1996*]

Material: Holotype *Trifurcula oishiella* ♀, Japan, Honshu, reared from galls on *Prunus*, T. Oishi, Genitalia slide EvN2916 (Entomological Institute, Hokkaido University).

52 *Glaucolepis hamirella* (Chrétien) and *G. saturejae* (Parenti). The holotype of *G. hamirella* is very similar to male *G. saturejae* (Figs 80–85), and a synonymy was considered. However, DNA barcodes of Italian populations are very different from those from Morocco, Cyprus and Turkey together, and both clusters show again a lot of variation. A detailed morphological and molecular study is required to solve the status of this complex and until then we keep the species separate. Since the type of *G. hamirella* is from North Africa and *saturejae* from northern Italy, we consider the Turkish/Moroccan cluster as *G. hamirella*, the others as *G. saturejae*. Some specimens of *G. saturejae* studied by EvN earlier may carry labels with the name *Trifurcula hamirella*.

Material: Holotype *Nepticula hamirella*: ♂, Algeria: Biskra, v.1909, Chrétien, Genitalia slide KL0718 (MNHN).



Figures 80–85. *Glaucolepis* species, male genitalia. 80–82 *G. hamirella* (Chrétien, 1915), lectotype, slide KL0718 **83–85** *G. saturejae* (Parenti), Italy (Asti) fraz. Valmanera, Oasi WWF, 13.vii.2006, G. Baldizzone, slide EvN3749.

53 *Glaucolepis magna* (A. Laštůvka & Z. Laštůvka, 1997). The recently described *Trifurcula collinella* Nel, 2012 from France is here synonymised with *G. magna*. The holotype was examined (Figs 86–89), and the relatively large moth shows clearly the typical dorsal spot of this species and the genitalia are similar (e.g. Laštůvka and Laštůvka 1997; van Nieukerken et al. 2006). The long processes of the tegumen, as described by Nel (2012), are in fact the arms of the uncus, that is broken in the middle (Fig. 87). Material: Holotype *Trifurcula collinella* ♂, France, Var, Tourves, D64, Rte de Mazaugues, Malausse, 25.iv.2011, Jacques Nel, Genitalia slide JN25662 (J. Nel, personal collection, later to be deposited in the Tiroler Landesmuseum Ferdinandeum).

Figures 86–89. *Glaucolepis magna* (Laštuvka, A. & Z., 1997), holotype of *Trifurcula collinella* Nel, 2012, adult, labels and male genitalia.

54 *Glaucolepis*: "unassigned to group". The two Neotropical species placed in *Glaucolepis* (Puplesis and Robinson 2000) almost certainly do not belong there. The male genitalia show some resemblance to species in the *Glaucolepis raikhonae* group. However, we examined three males of *G. argentosa* Puplesis & Robinson, 2000, which show none of the apomorphies of *Glaucolepis* + *Trifurcula* (van Nieukerken 1986b), the trifurculate hindwing (Fig. 90), "velvet patch on underside hindwing" or the three pairs of abdominal tufts. The hindwing does show two remarkable groups of special scales on the humeral lobe (Fig. 90). It is possible that this species forms yet another Neotropical generic group. Unfortunately we have only fragments of the DNA barcode, that do not provide enough information on its taxonomic place. *Glaucolepis aerifica* again has very different genitalia and may belong to an entirely different clade.

55 *Trifurcula bicolorella* (Chrétien, 1915), **comb. n.** This species was described as *Bucculatrix* and not recognized as nepticulid, until Zdeno Tókar (Slovakia) identified it as a *Trifurcula* when studying sketches of Western Palearctic *Bucculatrix* genitalia made by Gerfried Deschka. Unfortunately the holotype (Figs 91, 92) is a



Figure 90. *Glaucolepis argentosa* Puplesis & Robinson, 2000, paratype, venation of right wings, slide 4502, RMNH.INS. 24502. Note the absence of Cu in forewing, absence of trifurculate Rs+M in hindwing and presence of two groups of special scales on humeral lobe.

female, and female genitalia are not very diagnostic in this group, and no males are available. The wing markings resemble *T. aurella* Rebel, 1933, that feeds on *Spartium junceum*, and one unnamed species from Spain that feeds on *Retama* species. On the type locality most likely only *Retama* species occur, more North African material is needed to establish the identity of this species firmly.
Material: Holotype ♀, [Tunisia, Gafsa, Mai 1907–1912], "169", "Type", "Eukitt-Präp. Nr. 1219 G. Deschka, "Bucculatrix bicolorella" (Muséum National d'Histoire Naturella, Paris).

56 *Ectoedemia scobleella* Minet, 2004 (replacement name for the junior homonym *E. scoblei* Minet, 1990) is here recombined with *Fomoria* on the basis of examination of the type series (Figs 93–97).
Material: Holotype ♂, Madagascar, Forêt de Befasy, 40 km S. de Morondava, 1–10.i.1956, R. Paulian, Genitalia slide EvN4799 (Muséum National d'Histoire Naturelle, Paris).

57 *Fomoria argyraspis* (Puplesis & Diškus, 1995), **comb. n.** We here tentatively move this species from *Acalyptris* to *Fomoria* on the basis of the genitalia and externals that are very similar to a number of *Fomoria* species and not to any other *Acalyptris* species.

Figures 91–92. *Trifurcula bicolorella* (Chrétien, 1915), holotype female, labels and genitalia, slide Deschka 1219.



Figures 93–97. *Fomoria scobleella* (Minet, 2004), holotype male, labels and male genitalia.

58 *Fomoria sporadopa* (Meyrick, 1911), **comb. n.** This species from Sri Lanka is very difficult to place. EvN studied the holotype, the only known specimen, in 1986 (Figs 98–101). The venation, studied in the intact specimen shows a curved main trunk of R+Rs+M but further details were hardly visible; the abdomen has a triangular S2a (van Nieukerken 1986b), both speaking against placement in *Acalyptris* where Diškus and Puplesis (2003) placed it. The genitalia are rather peculiar, without transtilla, and elaborate carinal processes. For the time being we place this in *Fomoria*, that comes closest, but it may well belong to a separate clade.
Material: Holotype ♂, [Sri Lanka, Eastern Province] Ceylon, Trincomali, 8.vi.1907, BF [T. Bainbrigge Fletcher], Genitalia slide BM24103 (Natural History Museum, London).

59 *Etainia trifasciata* (Matsumura, 1931). Although the holotype of *Nepticula trifasciata* Matsumura is badly damaged, the forewing pattern is clearly identical to that of *Obrussa tigrinella* Puplesis, 1985. Since this species is common in Japan, and no other species with the same colour pattern occur there, we can safely conclude there is only one trifasciate *Etainia* in Japan, hence we synonymise *O. tigrinella* here.

Figures 98–101. *Fomoria sporadopa* (Meyrick, 1911), holotype male, labels and male genitalia.

Translation of the original description in Japanese by Matsumura:

"Body and wing with grey-yellow and white with reflection. Head orange yellow, both sides silvery white. Antenna grey white. Forewing darker with three bands. One near wing base, one in middle and a little one outwards. Third fascia at the apex. Terminal and dorsal cilia grey-yellow white. Costal fringe darker. Hindwing darker. Legs dark grey. Seems to be the smallest moth in Japan. Early September. Hokkaido." [*Translated by Liu Youqiao, August 1996*]

60 *Acalyptris falkovitshi* (Puplesis, 1984). We synonymize here *Microcalyptris arenosus* Falkovitsh, 1986 and *M. vittatus* Puplesis, 1984, after earlier *M. turanicus* Puplesis, 1984 was synonymized with *A. falkovitshi* (van Nieukerken 2010). We act as First Reviser (ICZN article 24.2) to identify *M. falkovitshi* as having priority to the other species that were published in the same paper (*vittatus* and *turanicus*). The slight differences given earlier (Puplesis 1990; 1994) do not hold when a larger sample is studied: the sclerotization of the T-shaped sclerites on segments 4–8 may vary, probably depending on age. This is the larger species in this group, usually with a longitudinally brown stripe on forewings, a pointed gnathos and conspicuous tufts on tergites 4–8.

61 Unplaced *Acalyptris*. Several of these are morphologically rather different from other *Acalyptris* species. *Acalyptris distaleus* almost certainly does not belong in this genus, it is part of a group of species, occurring all in southern California and Arizona.

62 *Zimmermannia*. The North American species of *Zimmermannia*, previously recognised as the *Ectoedemia castaneae* group, were revised within the revision of *Ectoedemia* (Wilkinson 1981; Wilkinson and Newton 1981). They recognised in all 12 species, but failed to provide sufficient diagnostic characters or keys to separate all these often extremely similar species, several of which were based on a single specimen only. Whereas some lectotypes were selected, a major shortcoming was that the types of the oldest name in this group, *Trifurcula obrutella* Zeller, had not been examined, and the synonymy of *Nepticula bosquella* Chambers with *obrutella*, earlier suggested by Busck (1903) was simply accepted. Comparing the descriptions it is clear that Zeller described the pale headed species ("Kopf bleich lehmgelblich"= Head pale loam yellowish), whereas Chambers described the black headed species. Examination of the *obrutella* types confirms this and leads to a change of the interpretation of *obrutella* as the species currently known as *Ectoedemia piperella* Wilkinson & Newton, 1981, thus reviving *obrutella's* former synonym *bosquella* as the valid name for the black headed species. Also the identity of the Lectotype of *Nepticula grandisella* Chambers, 1880 (indicated as Holotype by Wilkinson 1981) lead to a re-interpretation of this name, here synonymised with *E. chloranthis* Meyrick, 1928 and *E. acanthella* Wilkinson & Newton, 1981.

To establish the identities of these old names firmly, we designate lectotypes below for *Trifurcula obrutella* and *Nepticula grandisella*. As a result of our revision we reduce the number of valid species in Eastern North America to five, synonymising eight names, and reviving two old names. We will more extensively diagnose and illustrate these species in a forthcoming publication. There are more North American species in this genus, particularly in the West and South, but these are all unnamed species.
Material: *Nepticula grandisella* Chambers: Lectotype ♂ (designated here), MCZ-ENT00001302, [United States], Texas, [Bosque Co., Norse, Gustav W. Belfrage], Chambers, Type 1302, Genitalia slide CNC 3495 (Museum of Comparative Zoology, Harvard University).
Trifurcula obrutella Zeller: Lectotype ♂ (designated here), MCZ-ENT00014248, [United States], Texas, Dallas, Boll, Type 14248, Genitalia slide DRD 2936 (Museum of Comparative Zoology, Harvard University). For the data of the other types in this group see Wilkinson and Newton (1981).

63 *Ectoedemia insularis* Puplesis, 1985 was described from two specimens from Sakhalin. We transfer the species here to the *populella* group. In the male genitalia it resembles *E. intimella* (Zeller, 1848) closely, a species also recorded from east Asia, and even Sakhalin. We compared genitalia of specimens reared from *intimella*-like leafmines on *Salix* in South Korea, and also examined light collected adults from Korea and an earlier reported female from Japan (van Nieukerken et al. 2012b). The male genitalia of these are identical to those of the *E. insularis* holotype, illustrated by Stonis and Rocienė (2013), and show a slight, but constant

difference to European populations: European *E. intimella* has a wider gnathos and the ventral carinae are wider apart than in *insularis*. Adult externals, larva and biology of East Asian and European populations are identical, but there is a large DNA barcoding gap (more than 5%). On the base of these findings we consider all East Asian "*intimella*" as *E. insularis* and keep both tentatively as separate species. Study of populations and their phylogeography between East Asia and Europe is needed to establish the status of these populations more firmly. BIN's: *E. insularis*: BOLD:AAD0468, *E. intimella*: BOLD:AAD0467.

64 *Ectoedemia sinevi* Puplesis, 1985. This species is placed in the *populella* group on the basis of its male genitalia, and DNA barcodes of material that we consider conspecific. This species will be redescribed in a forthcoming paper on Japanese *Ectoedemia* (Hirano et al. in prep.).

65 *Ectoedemia turbidella* (Zeller, 1848). *Ectoedemia similigena* Puplesis, 1994 was described from a single series from Yalta on the Crimea. Earlier we found the slight differences in genitalia just sufficient to keep it tentatively as a separate species (van Nieukerken et al. 2010). Meanwhile, we have been able to obtain a partial DNA barcode of the dissected and illustrated paratype male (RMNH.INS.23924, genitalia slide EvN3924). The 420 basepairs are almost the same as in Western European specimens of *E. turbidella*, with just 2 basepairs difference, and BOLD classifies the barcode in the same BIN (BOLD:AAD4374). This small difference over a large geographic gap, together with the weak differences observed earlier, prompt us to reverse our earlier opinion, and synonymize *E. similigena* with *E. turbidella*.

66 *Ectoedemia spiraeae* Gregor & Povolný *& jacutica* Puplesis, 1988. Recently, Stonis et al. (2015b) reinstated *Ectoedemia jacutica* Puplesis, 1988 as valid species, after we had synonymised it earlier with *E. spiraeae* (van Nieukerken et al. 2010). In China and Japan we (and Japanese collectors) collected a species on *Spiraea* and *Aruncus* that we hitherto identified as *E. spiraeae*, although it has a large barcode gap with European specimens (van Nieukerken et al. 2012b). Considering the distribution being rather close to the type locality Yakutsk and some morphological similarities, we consider it likely that the Japanese and NE Chinese populations belong to *E. jacutica* (with BIN BOLD:AAI9354). The situation is comparable to that of *E. insularis* and *E. intimella* (above) and thus studies of populations between these are needed for more final conclusions. For now we consider specimens from the much more western Altai (van Nieukerken et al. 2010) to belong to *E. spiraeae*.

67 *Ectoedemia minimella* (Zetterstedt, 1839). The North American *Ectoedemia canadensis* (Braun, 1917) is here synonymised with *E. minimella*. Details will be published elsewhere.

68 *Ectoedemia occultella* (Linnaeus, 1767). This is one of the most widespread Nepticulidae, occurring from westernmost Europe to Japan and throughout northern North America, extending south in the mountain ranges to Colorado and Tennessee. The north American form was described as *Ectoedemia lindquisti* (Freeman, 1962), but we see no reason to treat North American populations

differently from Palearctic ones. DNA barcodes: Europe to Mongolia: BOLD:AAD0469; Japan: BOLD:ACU6927; North America: BOLD:AAH4532, distance 2.73%.

69 *Ectoedemia ulmella* (Braun, 1912). We synonymize *E. andrella* Wilkinson, 1981 on the basis of the description and illustrations of the holotype. The genitalia are indistinguishable, and also the characteristic androconial scales on the hindwing are identical.

70 *Nepticula alpinella* Herrich-Schäffer, 1863b, *N. alticolella* Herrich-Schäffer, 1863c and *N. reuttiella* Herrich-Schäffer, 1863c. These three names were described in a paper dealing with a collecting trip to Engadin, in the Swiss Alps, and are available names. However, they have never appeared again in the literature, apart from Snellen (1873), who listed the names as missing from the Staudinger & Wocke catalogue (Wocke 1871), and these names can thus easily be considered Nomina Oblita (ICZN art. 23.9) whenever they are threatening stability of any junior synonyms. Only the name *reuttiella* was picked up in the cardindex of the London Natural History Museum (Beccaloni et al. 2005), and as a consequence appears on some websites that are copying these names uncritically. Looking at the descriptions and localities, *N. alpinella* may refer to *Stigmella thuringiaca*, *N. alticolella* could be *S. stelviana* or *dryadella* and *N. reuttiella* an abberant *S. prunetorum*.

71 *Nepticula oritis* Meyrick, 1910. This species was described from a single specimen from India, Himachal Pradesh, Simla Hills. It is probably a fasciate *Stigmella* species, of which there are many, that can only be identified with genitalia. Meyrick (1910) stated in the introduction that the material was from the Indian Museum (in Kolkota), but we failed to get any information on the whereabouts of the holotype, and therefore this species is better treated as a *Nomen oblitum*, until a type turns up.

72 *Nepticula xuthomitra* Meyrick, 1921. Meyrick described this species on the basis of one specimen from Pretoria, that unfortunately is no longer present in TM or BMNH. The description is too vague to identify the species with any certainty, and since it is very well possible that *N. xuthomitra* is in fact a senior synonym of one of the currently recognized valid species, it is best regarded as a *Nomen oblitum*. Once a synonymy is established, it is best to follow up with an action as described in ICZN 23.9, if this name threatens the stability of nomenclature of another species.

73 *Nepticula anguinella* Clemens, 1861 and *N. platea* Clemens, 1861. These names were given to larvae in incomplete leafmines on oaks. We think that the description can be best interpreted as *Stigmella* species, even though Wilkinson (1981) recombined them with *Ectoedemia*. He did this on the base of the larva having "ten square brown or blackish spots", indeed a character of several *Ectoedemia* species. However, we do not know of any oak feeding *Ectoedemia* in North America with this character, certainly the common *E. similella* Braun does not have such plates. *Stigmella* larvae that feed with the venter upwards, such as those in the *saginella* group, also show dots: the often conspicuous ganglia. Since North American *Stigmella* oak mines are all very similar, there is no way to identify these species on

the basis of the very short descriptions, and by absence of any reared adults or other type material we think it is better to leave these names as *Nomina oblita*. If at any time, a synonymy of these species can be established, these names almost certainly have priority to the currently valid names; we therefore strongly advise taxonomists then to reverse priority according to ICZN 23.9, declare these names as *Nomen oblitum*, and declare the junior synonym as *Nomen protectum.*

74 *Unplaced unavailable names.* These names are partly copied from van Nieukerken (1986a) and we add a whole series of names by Dovnar-Zapol'skij from Russia and Central Asia, based on leafmines only, in two poorly known publications (Dovnar-Zapolski 1969; Dovnar-Zapolski and Tomilova 1978). Several of these almost certainly belong to known species, but *Stigmella atraphaxidella* refers to an interesting new host association (*Atraphaxis*, Polygonaceae). The few names that we could identify have been listed under the respective valid names (*S. acerna, S. turbatrix, S. lediella*, see notes 4, 12 and 39).

75 *Opostegoides.* We recombine four South African species, now in *Opostega*, with this genus after study of genitalia slides of type material, prepared by L. Vári: *Opostegoides granifera* (Meyrick, 1913), **comb. n.** (Fig. 102), *O. melitardis* (Meyrick, 1918), **comb. n.** (Fig. 103), *O. pelocrossa* (Meyrick, 1928), **comb. n.** (Fig. 104), *O. praefusca* (Meyrick, 1913), **comb. n.** (Fig. 105).

76 *Opostega cretatella* Chrétien, 1915. Unfortunately no type material of this species could be found. The species has been described from Algeria, Biskra, in April, on the northern edge of the Sahara. It is a very large species, with a wingspan of 14.5 mm. When comparing the description with that of *O. rezniki* Kozlov, 1985, the similar large size (16–17 mm) and uniform colour pattern with a single dorsal dot is striking. Also *O. rezniki* is described from a desert/steppe habitat, in Kazakhstan, 150 km NNE of Almaty, Sarytaukum, flying in mid-May. We consider it very likely that they represent the same species, with a similar distribution as many desert dwelling species, such as *Acalyptris psammophricta* (see van Nieukerken 2010) and thus synonymise them here. Obviously a confirmation from new material collected in North Africa would be welcome.

77 *"Opostega".* Until the eighties of the last century all Opostegids were placed in the single genus *Opostega* Zeller. After the generic revision (Davis 1989) most species have been moved to *Pseudopostega* or *Opostegoides*, but a revision of the African and Australian species has not yet taken place. Since *Opostega* s.str. is really a rather small Palearctic genus, and probably most Australian and African belong to *Pseudopostega* or *Opostegoides*, we prefer to place the species that have not yet been examined in detail tentatively in a separate "*Opostega*". We have been able to place several South African species correctly, since Lajos Vári had already dissected many types, present in the Ditsong Museum of Natural History in Pretoria (former Transvaal Museum).

78 *Pseudopostega chalcopepla* (Walsingham, 1908). We synonymize *Pseudopostega cyrneochalcopepla* Nel & Varenne, 2012 on the basis of virtual identical genitalia (Figs 113, 114) and external characters. We examined and barcoded the paratype. The

Figures 102–105. South African *Opostegoides* species, male genitalia. **102** *O. granifera* (Meyrick, 1913), holotype, slide TM4044 **103** *O. melitardis* (Meyrick, 1918), holotype, slide TM4037 **104** *O. pelocrossa* (Meyrick, 1928), holotype, slide TM4042 **105** *O. praefusca* (Meyrick, 1913), holotype, slide TM4043.

barcodes of the two specimens from Mediterranean islands (Corsica and Cyprus: RMNH.INS.24603) show large barcode distances to the mainland populations, but otherwise we find no differences. For now we prefer to keep these all in one

Figures 106–114. *Pseudopostega* species, female (106–108) and male genitalia. **106, 107** *P. idiocoma* (Meyrick, 1918), syntype, slide TM4036 **108** *P. orophoxantha* (Meyrick, 1921), holotype, slide TM4034 **109** *P. amphimitra* (Meyrick, 1913), syntype, slide TM4039 **110** *P. phaeosoma* (Meyrick, 1928), holotype, slide TM4046 **111** *P. symbolica* (Meyrick, 1914), holotype, slide TM4038 **112** *P. tincta* (Meyrick, 1918), holotype, slide TM4035 **113, 114** *Pseudopostega chalcopepla* (Walsingham, 1908): **113** paratype of *Pseudopostega cyrneochalcopepla* Nel & Varenne, 2012, slide J Nel 24980 **114** France, Var, Mazaugues, slide EvN4279.

variable species, until more is known of these and other island populations. Island populations often have large barcoding gaps to mainland populations, which taken alone is in our opinion not sufficient for species status.

Material: *Pseudopostega cyrneochalcopepla*: 1♂ paratype, [France, Corsica] Pertusato, Bonifacio, 24.v.2011, P. Varenne, DNA extracted from 1 leg, RMNH.INS.550071, collection J. Nel (later to be deposited in the Tiroler Landesmuseum Ferdinandeum).

79 *Pseudopostega* (African species). We recombine six South African species, now in *Opostega*, with this genus after studying genitalia slides of type material, prepared by L. Vári: *Pseudopostega amphimitra* (Meyrick, 1913), **comb. n.** (Fig. 109), *P. idiocoma* (Meyrick, 1918), **comb. n.** (Fig. 106, 107), *P. orophoxantha* (Meyrick, 1921), **comb. n.** (Fig. 108), *P. phaeosoma* (Meyrick, 1928), **comb. n.** (Fig. 109), *P. symbolica* (Meyrick, 1914), **comb. n.** and *P. tincta* (Meyrick, 1918), **comb. n.**

80 *Nepticula belfrageella* Chambers, 1875a. The slide labelled "Type" is not a nepticulid, but belongs to an unidentified Gelechiidae (personal communication J.F. Landry, D. Adamski). Also the description most likely does not refer to a nepticulid species, so we exclude it here.

Material: Genitalia slide ♂ examined, CNC3496, *Nepticula belfrageella* Cham, #99, #1555, no specimen [note in pencil], TYPE, G.L.[ewis], remounted by PJN.[ewton] (USNM).

Acknowledgements

This catalogue, summarising our taxonomic work on Nepticuloidea for more than 30 years, would never have been possible without the help of numerous people, of which we can only name those who were more directly involved in this paper. We are grateful for the support of many people who lend or donated us important material, or provided information on various matters: Mansour Abaii, Ian Barton, Alison Brown, Jeremy deWaard, Charley Eiseman, Alessandro Giusti, Bert Gustafsson, Rachel Hawkins, Nagao Hirano, Toshiya Hirowatari, Roland Johansson, Ole Karsholt, Natalia Kirichenko, Shigeki Kobayashi, Martin Krüger, Jean-François Landry, Aleš Laštůvka, Zdeněk Laštůvka, Bong-Woo Lee, David Lees, Wolfram Mey, Joël Minet, Vazrick Nazari, Jacques Nel, Masahiro Ohara, Philip Perkins, Gregory Pohl, Yagi Sadahisa, Klaus Sattler, Andreas Segerer, Sergey Sinëv, Jonas R. Stonis, Angela Telfer, Zdeno Tókar, Robert Trusch, Kevin Tuck, Hectór Vargas and the late Liu Youqiao.

Sjaak Koster made many genitalia preparations and Kees van den Berg assisted us in many ways.

The following persons provided information on the chapter on taxonomic history: Arūnas Diškus, David Lees, Steph Menken, Andrius Remeikis, Malcolm Scoble, Jonas Rimantas Stonis and Diana Wilkinson. We are grateful to Roland Johansson for his permission to publish scans of his original watercolours, and to Bengt Å. Bengtsson for his assistance in this matter. Henk Caspers and Niko Korenhof are acknowledged for scanning colour art.

We are grateful to Peter Huemer, Jean-François Landry, Marko Mutanen, Andreas Segerer and Christian Wieser to allow us to add their nepticuloid barcode data to our dataset and to BOLD staff for making this possible.

Yuri Roskov and Luisa Abucay of the Catalogue of Life were very helpful in translating our database into CoL format, and Dimitri Koureas, Vincent Smith and Edward Baker kindly helped to find the best procedure to maintain and update the Scratchpad with the new classification.

We acknowledge the insights and useful comments of reviewers Malcolm Scoble and Zdeněk Laštůvka and are grateful to the ZooKeys editorial office and editor Thomas Simonsen for organising the complex simultaneous publication of three related nepticulid manuscripts.

References

Achterberg C van (1983) Revisionary notes on the subfamily Gnaptodontinae with description of eleven new species (Hymenoptera, Braconidae). Tijdschrift voor Entomologie 126: 25–57. http://www.biodiversitylibrary.org/part/66040

Aguiar AMF, Karsholt O (2006) Systematic catalogue of the entomofauna of the Madeira Archipelago and Selvages Islands, Lepidoptera. Vol. 1. Boletim do Museu Municipal do Funchal Supplement 9: 5–139.

Amsel HG (1934) Neue palästinische Lepidopteren. Mitteilungen aus dem Zoologischen Museum in Berlin 20: 271–319, pls.

Amsel HG, Hering M (1931) Beitrag zur Kenntnis der Minenfauna Palästinas. Deutsche Entomologische Zeitschrift 1931: 113–152, pls 1–2. doi: 10.1002/mmnd.193119310203

Aspöck H (2003) Erinnerungen an Dr. Josef Klimesch (1902–1997). Denisia 8: 237–277.

Baldizzone G, van der Wolf H, Landry J-F (2006) Coleophoridae, Coleophorinae (Lepidoptera). World Catalogue of Insects 8: 1–215.

Beccaloni G, Scoble M, Robinson G, Pitkin B (2005) The Global Lepidoptera names index. Natural History Museum. http://www.nhm.ac.uk/our-science/data/lepindex/index.html

Beccaloni G, Holloway J, Kitching I, Sattler K, Scoble M, Tuck K, Vane-Wright D (2009) A tribute to Dr Gaden S. Robinson (1949–2009). Systematics and Biodiversity 7: 351–355. doi: 10.1017/s1477200009990247

Bedell G (1848) Description of *Microsetia quinquella*, a new species of moth of the family Tineidae. Zoologist 6: 1986.

Beirne BP (1945) The male genitalia of the British Stigmellidae (Nepticulidae) (Lep.). Proceedings of the Royal Irish Academy Section B 50: 191–218. http://www.jstor.org/pss/20490833

Bengtsson BÅ, Johansson R, Palmqvist G (2008) Fjärilar: Käkmalar - säckspinnare. Lepidoptera: Micropterigidae - Psychidae. ArtDatabanken, SLU, Uppsala, Nationalnyckeln till Sveriges flora och fauna (= Encyclopedia of the Swedish Flora and Fauna) DE 1–13: 646 pp.

Boheman CH (1853) Entomologiska anteckningar under en resa i Södra Sverige 1851. Kongliga Svenska Vetenskaps-Akademiens Handlingar 1851: 55–210. doi: 10.5962/bhl.title.35818

Borkowski A (1969) Studien an Stigmelliden (Lepidoptera). Teil I. Zur Verbreitung, Biologie und ökologie der Stigmelliden in den polnischen Sudeten. Polskie Pismo Entomologiczne 34: 95–122.

Borkowski A (1970a) Studien an Stigmelliden (Lepidoptera). Teil II. *Fedalmia thymi* sp. n.: eine neue Art aus Mitteleuropa. Polskie Pismo Entomologiczne 40: 69–78.

Borkowski A (1970b) Studien an Stigmelliden (Lepidoptera). Teil III. Beitrag zur Kenntnis der Stigmellidenfauna Polens. Polskie Pismo Entomologiczne 40: 541–555.

Borkowski A (1972a) Studien an Nepticuliden (Lepidoptera). Teil IV. Bemerkungen zur Nomenklatur und Systematik der Familie Nepticulidae. Polskie Pismo Entomologiczne 42: 689–709, 2 pls.

Borkowski A (1972b) Studien an Nepticuliden (Lepidoptera). Teil V. Die europäischen Arten der Gattung *Nepticula* Heyden von Eichen. Polskie Pismo Entomologiczne 42: 767–799.

Borkowski A (1975) Studien an Nepticuliden (Lepidoptera). Teil VI. Die Verbreitung der Nepticuliden in Polen. Polskie Pismo Entomologiczne 45: 487–535, 10 pls.

Börner C (1925) 22 Ordn. Lepidoptera, Schmetterlinge. In: Brohmer P (Ed.) Fauna von Deutschland, Ein Bestimmungsbuch unserer heimischen Tierwelt. Quelle & Meyer, Leipzig, 358–387.

Bourquin F (1961) Microlepidopteros nuevos con sus biologias. Revista de la Sociedad Entomológica Argentina 23: 31–46.

Bradley JD (1957) Microlepidoptera from Rennell and Bellona Islands. In: Wolf T (Ed.) The Natural History of Rennell Island, British Solomon Islands. Danish Science Press, Copenhagen, 87–112.

Bradley JD (1961) Microlepidoptera from the Solomon islands: additional records and descriptions of Microlepidoptera collected in the Solomon islands by the Rennell Island Expedition 1953–1954. Bulletin of the British Museum (Natural History) Entomology 10: 111–168, pls 5–19. doi: 10.5962/bhl.part.16259

Bradley JD (1965) Microlepidoptera. Ruwenzori Expedition 1952 II (12): 81–148. http://biodiversitylibrary.org/page/43669653

Bradley JD, Fletcher DS, Whalley PES (1972) Kloet and Hincks. A checklist of British Insects. Second Edition (Revised) part 2: Lepidoptera. Handbooks for the Identification of British Insects 11: 1–153.

Braun AF (1910) New species of Tineina from California. Entomological News 21: 171–179. http://biostor.org/reference/84670

Braun AF (1912) Notes on North American species of *Nepticula*, with descriptions of new species (Lepidoptera). Journal of the Cincinnati Society of Natural History 21: 84–101. http://biodiversitylibrary.org/page/12268566

Braun AF (1914) Notes on North American species of *Nepticula* with description of new species (Lepidoptera). Canadian Entomologist 46: 17–24. doi: 10.4039/ent4617-1

Braun AF (1915) New genera and species of Tineina. Canadian Entomologist 47: 188–197. doi: 10.4039/ent47188-6

Braun AF (1916) *Nepticula rhamnicola* nom. nov. Entomological News 27: 56. http://biostor.org/reference/144453

Braun AF (1917) Nepticulidae of North America. Transactions of the American Entomological Society 43: 155–209, pls. 5–8. http://biostor.org/reference/98155

Braun AF (1918) New genera and species of Lyonetiidae (Microlepidoptera). Entomological News 29: 245–251. http://biodiversitylibrary.org/page/26262244

Braun AF (1923) Superfamily Nepticuloidea. Family 5. Nepticulidae. In: Forbes WTM, The Lepidoptera of New York and neighboring states. Memoirs of the Cornell University Agricultural Experiment Station 68: 79–98. http://biodiversitylibrary.org/page/19819738

Braun AF (1925a) Some undescribed Microlepidoptera and notes on life histories. Transactions of the American Entomological Society 51: 13–17. http://www.jstor.org/stable/25077120

Braun AF (1925b) Microlepidoptera of Northern Utah. Transactions of the American Entomological Society 51: 183–226. http://www.jstor.org/stable/25077135

Braun AF (1927a) New Microlepidoptera from Ontario. Canadian Entomologist 59: 56–59. doi: 10.4039/Ent5956-3

Braun AF (1927b) Descriptions of new Microlepidoptera. Transactions of the American Entomological Society 53: 191–199. http://www.jstor.org/stable/25077178

Braun AF (1930) Notes and new species of Microlepidoptera from the mineral springs region of Adams County, Ohio. Transactions of the American Entomological Society 56: 1–17. http://www.jstor.org/stable/25077235

Brown JW, Baixeras J, Brown R, Horak M, Komai F, Metzler EH, Razowski J, Tuck K (2003) Tortricidae (Lepidoptera). World Catalogue of Insects 5: 1–741.

Bruand T ([1851]) Catalogue systématique et synonymique des Lépidoptères du Département du Doubs. [6]Tinéides. Mémoires de la Société d'Emulation du Doubs 3: 58–92.

Bruand d' Uzelle T (1859) Classification des Tinéides et examen des caractères et de leur importance relative, d'apres la méthode naturelle (suite et fin). Annales de la Société Entomologique de France (sér. 3) 6: 601–702. http://biodiversitylibrary.org/page/8375522

Busck A (1900) New species of moths of the superfamily Tineina from Florida. Proceedings of the United States National Museum 23: 225–254, pl. 221. doi: 10.5479/si.00963801.23-1208.225

Busck A (1901) *Nepticula pomivorella* Packard; alias *Micropteryx pomivorella* Packard. Canadian Entomologist 33: 52. doi: 10.4039/ent3352-2

Busck A (1903) Notes on Brackenridge Clemens' types of Tineina. Proceedings of the Entomological Society of Washington 5: 181–220. http://www.biodiversitylibrary.org/item/19613

Busck A (1907) New american Tineina. Proceedings of the Entomological Society of Washington 8: 86–99. http://biodiversitylibrary.org/page/2345065.

Busck A (1910) List of Trinidad Microlepidoptera with descriptions of new forms. Bulletin of the Department of Agriculture, Trinidad and Tobago 66: 241–245.

Busck A (1913) Two microlepidoptera injurious to Chestnut. Proceedings of the Entomological Society of Washington 15: 102–104. http://biostor.org/reference/99133

Busck A (1914a) The chestnut bastminer (Lepidoptera, Tineidae). Insecutor Inscitiae menstruus 2: 3–4. http://biostor.org/reference/110552

Busck A (1914b) On the classification of the Microlepidoptera. Proceedings of the Entomological Society of Washington 16: 46–54. http://biodiversitylibrary.org/page/2597455

Busck A (1914c) Descriptions of new Microlepidoptera of forest trees. Proceedings of the Entomological Society of Washington 16: 143–150, pls vii-viii. http://biostor.org/reference/99128

Căpuşe I (1975) Eine neue Art der Gattung *Nepticula* aus Sambia (Lepidoptera,Nepticulidae). Reichenbachia 15: 211–213.

Caradja A (1920) Beitrag zur Kenntnis der geographischen Verbreitung der Mikrolepidopteren des palaearktischen Faunengebietes nebst Beschreibung neuer Formen. Deutsche Entomologische Zeitschrift Iris 34: 75–179. http://biodiversitylibrary.org/page/12720924

Carey P, Cameron S, Cerda ML, Gara RI (1978) Ciclo estacional de un minador subcortical de coigue (*Nothofagus dombeyi*). Turrialba 28: 151–153.

Carolsfeld-Krausé AG (1944) *Nepticula fragarivora* nov. spec. (Nepticulidae, Lepidoptera). Entomologiske Meddelelser 24: 158–159.

Carolsfeld-Krausé AG (1949) Some remarks on the synonymy of the *Fagus* feeding Nepticulae (Nepticulidae, Lepidoptera). Entomologiske Meddelelser 25: 299–310, Figs 291–296.

Chambers VT (1873) Micro-lepidoptera. Canadian Entomologist 5: 124–128. doi: 10.4039/Ent5124-7

Chambers VT (1875a) Tineina from Texas. Canadian Entomologist 7: 73–75. http://biodiversitylibrary.org/page/304808

Chambers VT (1875b) Tineina of the Central United States. Cincinnati Quarterly Journal of Science 2: 97–121. http://biodiversitylibrary.org/page/5251872

Chambers VT (1876) Tineina. Canadian Entomologist 8: 158–160. doi: 10.4039/Ent8158-8

Chambers VT (1878a) Descriptions of new Tineina from Texas, and others from more Northern localities. Bulletin of the United States Geological and Geographical Survey of the Territories 4: 79–106. http://www.biodiversitylibrary.org/part/101215

Chambers VT (1878b) Index to the described Tineina of the United States and Canada. Bulletin of the United States Geological and Geographical Survey of the Territories 4: 125–167. http://biodiversitylibrary.org/page/31549211

Chambers VT (1880a) Descriptions of some new Tineina, with notes on a few old species. Journal of the Cincinnati Society of Natural History 2: 179–194. http://biodiversitylibrary.org/page/7310943

Chambers VT (1880b) Notes upon some Tineid larvae. Psyche (Cambridge) 3: 63–68. doi: 10.1155/1880/38474

Chambers VT (1880c) Further notes upon some Tineid larvae. Psyche (Cambridge) 3: 135–137. doi: 10.1155/1880/18501

Chambers VT (1881) New species of Tineina. Journal of the Cincinnati Society of Natural History 3: 289–296. http://biodiversitylibrary.org/page/7313835

Chrétien P (1904) Les chenilles de Lins. Naturaliste Paris 26: 162–164. http://biodiversitylibrary.org/page/44538286

Chrétien P (1907) Les chenilles des Buplèvres. Naturaliste Paris 29: 89–91. http://biodiversitylibrary.org/page/44572245

Chrétien P (1908) Faune entomologique des Îles Canaries. Séjour de M.P. Lesne dans la Grande Canarie. 1. Microlépidoptères. Bulletin du Muséum National d'Histoire Naturelle Paris 14: 358–364. doi: 10.5962/bhl.part.5663

Chrétien P (1914) Descriptions de trois nouvelles espèces de *Nepticula* (Lep. Nepticulidae). Bulletin de la Société Entomologique de France 1914: 269–272. http://biostor.org/reference/144448

Chrétien P (1915) Contribution à la connaissance des Lépidoptères du Nord de l'Afrique. Annales de la Société Entomologique de France 84: 289–374. http://biostor.org/reference/97306

Chrétien P (1922) Lés Lépidoptères du Maroc. Galleriinae - Micropterygidae. In: Oberthür C (Ed) Études de Lépidoptérologie comparée. Oberthür, Rennes, 324–379. http://biodiversitylibrary.org/page/10505866

Clarke JFG (1955) Catalogue of the types specimens of Microlepidoptera in the British Museum (Natural History) described by Edward Meyrick, volume 1. Trustees of the British Museum (Natural History), London, 1: vii, 332 pp. http://www.biodiversitylibrary.org/bibliography/68439

Clemens B (1860) Contributions to American Lepidopterology - No. 5. Proceedings of the Academy of Natural Sciences of Philadelphia 12: 203–221. http://biostor.org/reference/98227

Clemens B (1861) Micro-Lepidopterous larvae. Notes on a few species, the imagos of which are probably undescribed. Proceedings of the Entomological Society of Philadelphia 1: 75–87. http://www.biodiversitylibrary.org/item/22852

Clemens B (1862a) New American Micro-Lepidoptera. Proceedings of the Entomological Society of Philadelphia 1: 131–137. http://biodiversitylibrary.org/page/3328877

Clemens B (1862b) North American Microlepidoptera. Proceedings of the Entomological Society of Philadelphia 1: 147–151. http://biodiversitylibrary.org/page/3328740

Clemens B (1865) North American Micro-Lepidoptera. Proceedings of the Entomological Society of Philadelphia 5: 133–147. http://www.biodiversitylibrary.org/part/9075

Clemens B (1872) Letters received from Dr. Brackenridge Clemens. 9. Letter of October 29th, 1860. In: Stainton HT (Ed) The Tineina of North America by (the late) Dr Brackenridge Clemens (being a collected edition of his writing on that group of insects). John van Voorst, London, 41–46. http://biodiversitylibrary.org/page/9496435

Common IFB (1990) Moths of Australia. E. J. Brill, Leiden, New York etc., 535 pp.

Costa O-G (1832–1836 [1836]) Fauna del Regno di Napoli, ossia enumerazione di tutti gli animali che abitano le diverse regioni di questo regno e le acque che le bagnano, contenente la descrizione de nuovi o poco esattamente conosciuti, Lepidotteri. Torchi del Tramater, Napoli: 370 pp. https://books.google.nl/books?id=pgRpDQFaaDwC

Crombrugghe de Picquendaele G de (1907) Observations sur quelques larves de microlépidoptères des environs de Bruxelles. Rectification. Revue mensuelle de la Société Entomologique Namuroise 7: 14. http://biodiversitylibrary.org/page/12300903

Davis DR (1978) New leaf-mining moths of the family Nepticulidae from Florida. Florida Entomologist 61: 209–224. doi: 10.2307/3494210

Davis DR (1979) *Manoneura*, a new name to replace the generic homonym *Oligoneura* Davis (Lepidoptera: Nepticulidae). Florida Entomologist 62: 276. doi: 10.2307/3494067

Davis DR (1983) Opostegidae. In: Hodges RW, Dominick T, Davis DR, Ferguson DC, Franclemont JG, Munroe EG, Powell JA (Eds) Check list of the Lepidoptera of America north of Mexico : including Greenland. Classey, London, 3.

Davis DR (1984) Nepticulidae. In: Heppner JB (Ed.) Atlas of Neotropical Lepidoptera, Checklist 1: Micropterigoidea-Immoidea. W. Junk, The Hague etc., 17–18.

Davis DR (1985) A re-examination of *Enteucha cyanochlora* Meyrick and its subsequent transfer to the Nepticulidae (Lepidoptera: Nepticuloidea). Proceedings of the Entomological Society of Washington 87: 142–145. http://biostor.org/reference/56072

Davis DR (1989) Generic revision of the Opostegidae, with a synoptic catalogue of the world's species (Lepidoptera: Nepticuloidea). Smithsonian Contributions to Zoology 478: 1–97. doi: 10.5479/si.00810282.478

Davis DR, Stonis JR (2007) A revision of the New World plant-mining moths of the family Opostegidae (Lepidoptera: Nepticuloidea). Smithsonian Contributions to Zoology 625: i-v, 1–212. doi: 10.5479/si.00810282.625

Davis DR, Wilkinson C (1983) Nepticulidae. In: Hodges RW, Dominick T, Davis DR, Ferguson DC, Franclemont JG, Munroe EG, Powell JA (Eds) Check list of the Lepidoptera of America north of Mexico: including Greenland. Classey, London, 2–3.

De Prins J, De Prins W (2016) Global Taxonomic Database of Gracillariidae (Lepidoptera). Belgian Biodiversity Platform. http://www.gracillariidae.net

De Prins W, De Prins J (2005) Gracillariidae (Lepidoptera). World Catalogue of Insects 6: 1–502.

DeGeer C (1752) Mémoires pour servir a l'histoire des insectes, 1. L'imprimerie de L.L. Grefing, Stockholm, 1: 707 pp. http://gallica.bnf.fr/ark:/12148/bpt6k97150b

DeGeer C (1771) Mémoires pour servir a l'histoire des insectes, 2. L'imprimerie de L.L. Grefing, Stockholm, 2: 1175 pp. http://gallica.bnf.fr/ark:/12148/bpt6k97151p

Deschka G (1998) Josef Klimesch (1902–1997). Nota Lepidopterologica 21: 224–227. http://www.biodiversitylibrary.org/part/85598

Diškus A, Puplesis R (2003) Catalogue of the world Nepticuloidea & Tischerioidea. In: Puplesis R, Diškus A (Eds) Nepticuloidea ir Tischerioidea (Lepidoptera) pasaulio ir lietuvos faunoje The Nepticuloidea & Tischerioidea (Lepidoptera) - a global review, with strategic regional revisions. Lutute publishers, Kaunas, 318–436. https://www.researchgate.net/publication/282577423

Disqué H (1912) Notizen über einige Kleinschmetterlinge. Entomologische Zeitschrift 26: 75. http://biodiversitylibrary.org/page/31141564

Dittrich R (1907) Dr. med. M. F. Wocke. Nachruf. Zeitschrift für Entomologie (Breslau), NF 32: xxxvi-xlvi, frontispiece. http://biodiversitylibrary.org/page/31431526, http://biodiversitylibrary.org/page/31431486

Doets C (1947) *Zimmermannia heringiella* nov. spec. Nepticulidae (Lepidoptera). Tijdschrift voor Entomologie 88: 504–506. http://biostor.org/reference/49601

Donner H, Wilkinson C (1989) Fauna of New Zealand. Nepticulidae (Insecta: Lepidoptera). Fauna of New Zealand 16: 1–88. http://www.biotaxa.org/fnz/article/view/1761

Doorenweerd C, Nieukerken EJ van, Menken SBJ (2015a) A global phylogeny of leafmining *Ectoedemia* moths (Lepidoptera: Nepticulidae): exploring host plant family shifts and allopatry as drivers of speciation PLoS ONE 10: e0119586. doi: 10.1371/journal.pone.0119586

Doorenweerd C, Nieukerken EJ van, Sohn J-C, Labandeira CC (2015b) A revised checklist of Nepticulidae fossils (Lepidoptera) indicates an Early Cretaceous origin. Zootaxa 3963: 295–334. doi: 10.11646/zootaxa.3963.3.2

Doorenweerd C, Nieukerken EJ van, Hoare RJB (2016a) Phylogeny, classification and divergence times of pygmy leafmining moths (Lepidoptera: Nepticulidae): the earliest lepidopteran radiation on Angiosperms? Systematic Entomology, early view. doi: 10.1111/syen.12212

Doorenweerd C, Nieukerken EJ van, Lopez-Vaamonde C, Menken SBJ (2016b) Increased diversification rates across lineages of leaf-mining moths after entering the new adaptive zone of the temperate region. In: Doorenweerd C, Branching out: the role of host plants in the diversification of leaf-mining moths. PhD Thesis, University of Amsterdam, Amsterdam, 185–205.

Doubleday H (1859) The zoologist synonymic list of British butterflies and moths, 2nd ed. Edward Newman, London: 40 pp.

Douglas JW, McLachlan R (1893) Henry Tibbats Stainton, F.R.S., &c. Entomologist's Monthly Magazine 29: 4, frontispiece. http://biodiversitylibrary.org/page/30359895

Dovnar-Zapolski DP (1969) Miniruyushchie nasekomye na rasteniyakh Kirgizii i sopredel'nykh territorij. [Mining insects on plants of Kirgizia and neighbourhood territory]. Akademiya Nauk Kirgizskoj SSR, Frunze: 149 pp. [In Russian]

Dovnar-Zapolski DP, Tomilova VN (1978) Miniruyushchie nasekomye Sibiri i sosednikh territorij. Mining insects of Siberia and adjacent territories. In: Zhdanova AA (Ed.) Nasekomye Vostochnoj Sibiri Mezhbuzovskij sbornik The Insects of Eastern Siberia. Ikrutskij Gozudarstvennyj Universitet, Irkutsk, 20–51. [In Russian]

Driel JW van, Nieukerken EJ van (1985) Report on a collecting trip to China, 1984, to study and collect Nepticulidae (Lepidoptera). Department of Animal Systematics, Subfaculty Biology, Vrije Universiteit, Amsterdam, 1–31. http://www.repository.naturalis.nl/record/403568

Dufrane A (1930) Les *Nepticula* des Saules en Belgique. Bulletin et Annales de la Société Entomologique de Belgique 70: 29–32.

Dufrane A (1942) Microlépidoptères de la faune Belge (1re note). Bulletin du Musée Royal d'Histoire Naturelle du Belgique 18: 1–12.

Dufrane A (1949) Microlépidoptères de la faune Belge (6e note). Bulletin de l'Institut Royal des Sciences Naturelles de Belgique 25: 1–11.

Dufrane A (1955) Microlépidoptères de la Faune paléarctique. Mémoires de la Société Royale Entomologique de Belgique 27: 184–193.

Dugdale JS (1971) Entomology of the Aucklands and other islands south of New Zealand: Lepidoptera, excluding non-crambine Pyralidae. Pacific Insects Monograph 27: 55–172. http://hbs.bishopmuseum.org/pim/pdf/pim27-55.pdf

Dugdale JS (1988) Lepidoptera : annotated catalogue, and keys to family-group taxa. Fauna of New Zealand 14: 1–262. http://www.biotaxa.org/fnz/article/view/1754

Duponchel PAJ ([1842–1845]) Nocturnes, Supplément 4. In: Godart J-B (Ed.) Histoire naturelle des Lépidoptères ou papillons de France. Méquignon-Marvis, Paris, xc + 554 pp, pls 51–90. http://www.biodiversitylibrary.org/item/39620

Duponchel PAJ (1844) Catalogue méthodique des lépidoptères d'Europe distribués en familles, tribus et genres, avec l'exposé des caractères sur lesquels ces divisions sont fondées, et l'indication des lieux et des époques où l'on trouve chaque espèce, pour servir de complément et de rectification à l'histoire naturelle des lépidoptères de France, devenue celle des

lépidoptères d'Europe par les supplémens qu'on y a ajoutés. Méquignon-Marvis fils, Paris. xxx + 523 pp. doi: 10.5962/bhl.title.24379

Durrant JH (1920) Lord Walsingham. 1843–1919 Entomologist's Monthly Magazine 56: 25–28, pl I in January issue. http://biodiversitylibrary.org/page/33378889

Dyar HG, Fernald CH, Hulst GD, Busck A (1903) A list of North American Lepidoptera and key to the literature of this order of insects. Bulletin of the United States National Museum 52: i-xix, 1–723. http://biostor.org/reference/57795

Eiseman C, Nieukerken EJ van (2015) *Stigmella microtheriella*. BugGuide. Iowa State University Entomology. http://bugguide.net/node/view/1035533

Emmet AM (1970) *Stigmella* (*Nepticula*) *paradoxa* Frey (*nitidella* Heinemann) (Lep. Nepticulidae) - a species new to Britain. Entomologist's Record and Journal of Variation 82: 3–5. http://biodiversitylibrary.org/page/30795773

Emmet AM (1971a) Notes on some of the British Nepticulidae (Lep.). Entomologist's Record and Journal of Variation 83: 75–83. http://biodiversitylibrary.org/page/29964877

Emmet AM (1971b) Notes on some of the British Nepticulidae (Lep.). Entomologist's Record and Journal of Variation 83: 240–248. http://biodiversitylibrary.org/page/29965092

Emmet AM (1973a) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 85: 77–80. http://biodiversitylibrary.org/page/29997461

Emmet AM (1973b) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 85: 176–180. http://biodiversitylibrary.org/page/29997616.

Emmet AM (1973c) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 85: 278–283. http://biodiversitylibrary.org/page/29997774

Emmet AM (1974a) The male genitalia of the British Stigmellidae illustrated by B. P. Beirne. Entomologist's Gazette 25: 217.

Emmet AM (1974b) *Ectoedemia bradfordi* sp. nov. (Lep., Nepticulidae), a hitherto undescribed species occurring in Kent. Entomologist's Gazette 25: 269–273.

Emmet AM (1974c) *Ectoedemia subapicella* (Stainton, 1886) (Lep., Nepticulidae) a synonym of *Ectoedemia albifasciella* (Heinemann, 1871). Entomologist's Gazette 25: 274–276.

Emmet AM (1974d) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 86: 75–80. http://biodiversitylibrary.org/page/39203140

Emmet AM (1974e) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 86: 103–108. http://biodiversitylibrary.org/page/39203177

Emmet AM (1974f) Notes on some of the British Nepticulidae, 2. Entomologist's Record and Journal of Variation 86: 147–153. http://biodiversitylibrary.org/page/39203239

Emmet AM (1974g) *Ectoedemia* (*Dechtiria*) *erythrogenella* (de Joannis, 1907) (Lep.: Nepticulidae). A species new to Britain. Entomologist's Record and Journal of Variation 86: 129–130, pl. 129. http://biodiversitylibrary.org/page/39203221

Emmet AM (1976) Nepticulidae. In: Heath J (Ed.) The moths and butterflies of Great Britain and Ireland, 1 Micropterigidae - Heliozelidae. Blackwell Scientific Publications Ltd, Oxford & London, 171–267.

Emmet AM (1979) Nepticulidae. In: Emmet AM (Ed.) A field guide to the smaller British Lepidoptera. The British Entomological & Natural History Society, London, 13–28.

Emmet AM (1991) The scientific names of the British Lepidoptera. Their history and meaning. Harley Books, Colchester.: 288 pp.

Emmet AM (1992) Henry Tibbats Stainton, 1822–1892. Entomologist's Gazette 43: 233–236.

Eppelsheim F (1891) *Nepticula gei* Wk. ab. *semicolorella* Epplshm. Stettiner Entomologische Zeitung 52: 351–352. http://biodiversitylibrary.org/page/9008300

Eyer JR (1967) A new species of Opostegidae from Florida. Florida Entomologist 50: 39–42. doi: 10.2307/3493205

Fabricius JC (1775) Systema entomologiae, sistens Insectorum Classes, Ordines, Genera, Species. Kortii, Flensburgi et Lipsiae, 832 pp. http://www.biodiversitylibrary.org/item/82400

Falkovich MI (1986) Cheshuekrylye (Lepidoptera) Ostancovykh gor Kul'dzhuktau i podgornoj ravniny (yugo- zapadnyj Kyzylkum). [Lepidoptera of the Monadnock Mountains of Kuldzhuktau and the subalpine plain (south west Kyzylkum).]. Trudy Vsesoyuznogo Entomologicheskogo Obshchestva 67: 131–186. [In Russian]

Fang J, Fang Z, Tang Z (2011) Atlas of woody plants in China. Distribution and climate. 2 vols. Higher Education Press, Springer, Beijing and Heidelberg etc. : xxxvi, 1972 pp. doi: 10.1007/978-3-642-15017-3

Fischer TC (2013) Pygmy moths (Lepidoptera, Nepticulidae) from Baltic Amber (Eocene). Zitteliana A 53: 85 – 92. http://www.palmuc.de/bspg/images/pdf/7_fischer.pdf

Fischer von Röslerstamm JE (1834-[1843]) Abbildungen zur Berichtigung und Ergänzung der Schmetterlingskunde, besonders der Microlepidopterologie als Supplement zu Treitschke's und Hübner's europaeischen Schmetterlingen, mit erläuterndem Text. Leipzig: 304 pp, 100 colour plates. http://hdl.handle.net/2027/chi.086851560

Fletcher TB (1933) Life-histories of Indian Microlepidoptera (second series). Cosmopterygidae to Neopseustidae. The Imperial Council of Agricultural Research, Delhi, Scientific Monograph 4: 1–85, 77 pls.

Fletcher TB, Clutterbuck CG (1945) Microlepidoptera of Gloucestershire, 7. Proceedings of the Cotteswold Naturalists' Field Club 28: 58–66.

Fologne E (1859) Supplément au catalogue des Lépidoptères de Belgique. Annales de la Société Entomologique Belge 3: 133–142. http://biodiversitylibrary.org/page/12851389

Fologne E (1860) *Nepticula ulmivora*. Entomologist's Weekly Intelligencer 8: 92. http://biodiversitylibrary.org/page/50816516

Fologne E (1862) Observations sur quelques Lépidoptères en Belgique. Annales de la Société Entomologique Belge 6: 161–169, pl. 162. http://biodiversitylibrary.org/page/12299752

Forbes WTM (1923) The Lepidoptera of New York and neighboring states. Primitive forms Microlepidoptera, Pyraloids, Bombyces. Memoirs of the Cornell University Agricultural Experiment Station 68: 1–792. doi: 10.5962/bhl.title.23875

Forbes WTM, Leonard MD (1930) A new leaf-miner of cotton in Porto Rico (*Nepticula gossypii* new species). Journal of the Department of Agriculture, Porto Rico 14: 151–157, pls. 15–16.

Ford LT (1950a) A new Stigmellid. Entomologist's Gazette 1: 39–40.

Ford LT (1950b) *Stigmella marionella* Ford. Entomologist's Gazette 1: 84.

Freeman TN (1962) A new species of *Nepticula* v. Heyd. on birch (Lepidoptera: Nepticulidae). Canadian Entomologist 94: 522–523. doi: 10.4039/Ent94522–5

Freeman TN (1967) A new species of *Nepticula* on Bur Oak in Ontario (Nepticulidae). Journal of Research on the Lepidoptera 6: 19–21. http://www.doylegroup.harvard.edu/~carlo/JRL/06/PDF/06-019.pdf

Frey H (1856) Die Tineen und Pterophoren der Schweiz. Meyer & Zeller, Zürich: xii+430 pp. http://www.mdz-nbn-resolving.de/urn/resolver.pl?urn=urn:nbn:de:bvb:12-bsb10307221-5

Frey H (1857) Revision der Nepticulen. Linnaea Entomologica 11: 351–446. http://biodiversitylibrary.org/page/12094991

Frey H (1858a) A new *Nepticula*. Entomologist's Weekly Intelligencer 4: 14. http://biodiversitylibrary.org/page/50828050

Frey H (1858b) A new *Nepticula*. Entomologist's Weekly Intelligencer 4: 27. http://biodiversitylibrary.org/page/50828063

Frey H (1858c) *Nepticula agrimoniae*. Entomologist's Weekly Intelligencer 4: 43–44. http://biodiversitylibrary.org/page/50828079

Frey H (1870) Ein Beitrag zur Kenntniss der Microlepidopteren. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 3: 277–296. http://biodiversitylibrary.org/page/49053701

Frey H (1873) Neue schweizerische Microlepidopteren. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 4: 143–144. http://biodiversitylibrary.org/page/49012774

Frey H (1880) Die Lepidopteren der Schweiz. Wilhelm Engelmann, Leipzig, xxvi+454 pp. doi: 10.5962/bhl.title.7905

Frey H, Boll J (1876) Einige Tineen aus Texas. Stettiner Entomologische Zeitung 37: 209–228. http://biodiversitylibrary.org/page/8987937

Frey H, Boll J (1878) Tineen aus Texas. Stettiner Entomologische Zeitung 39: 249–280. http://biodiversitylibrary.org/page/8988447

Fritsch A (1882) Fossile Arthropoden aus der Steinkohlen- und Kreideformation Böhmens. Beiträge zur Paläontologie Österreich-Ungarns und des Orients 2: 1–7, pls 1–2. https://archive.org/stream/beitrgezurpal02wien#page/n11/mode/2up

Fuchs A (1895) Kleinschmetterlinge der Loreley-Gegend. Stettiner Entomologische Zeitung 56: 21–52. http://biodiversitylibrary.org/page/8945071

GBIF Secretariat (2016) Global Biodiversity Information Facility. GBIF. http://www.gbif.org/

Gerasimov AM (1930) Zur Lepidopteren-Fauna Mittel-Asiens. I. Microheterocera aus dem Distrikt Kaschka-Darja (SO-Buchara). Ezhegodnik Zoologicheskago Muzeya Akademii Nauk SSSR 31: 21–48, pls. 8–17.

Gerasimov AM (1937) Minierende Motten V. Neue *Stigmella* (*Nepticula*)- und *Tischeria*-Arten. (Lepid. Stigmell. und Tischer.). Mitteilungen aus dem Zoologischen Museum in Berlin 22: 282–285. doi: 10.1002/mmnz.19370220213

Gerasimov AM (1952) Nasekomye Cheshuekrylye. I, 2, Gusenicy 1. Zoological Institute, Academy of Sciences USSR, Moskva, Leningrad, Fauna SSSR N.S. 56: 1–338 pp. [In Russian]

Gielis C (2003) Pterophoroidea & Alucitoidea (Lepidoptera). Apollo Books, Stenstrup, World Catalogue of Insects 4: 1–198 pp.

Gilligan TM, Baixeras J, Brown JW, Tuck KR (2014) T@RTS: Online World Catalogue of the Tortricidae (Ver. 3.0). http://www.tortricidae.com/catalogue.asp

Glitz CT (1865) Zweiter Nachtrag zu dem Verzeichnisse der bei Hannover vorkommenden Schmetterlinge. Jahresbericht der Naturhistorischen Gesellschaft zu Hannover 14: 42. http://biodiversitylibrary.org/page/11824310

Glitz CT (1872) Neue Microlepidopteren. Stettiner Entomologische Zeitung 33: 23–26. http://biostor.org/reference/111632

Glitz CT (1887) Ueber einige Arten *Nepticula*. Stettiner Entomologische Zeitung 48: 276–277. http://biodiversitylibrary.org/page/8940974

Goeze JAE (1783) Entomologische Beyträge zu des Ritter Linné zwölften Ausgabe des Natursystems. Dritten Theiles vierter Band. Weidmanns Erben und Reich, Leipzig, 3(4): i-xx, 1–178 pp. http://biodiversitylibrary.org/page/32010985

Gómez Bustillo MR (1981) Revalidación de Nepticulina (Börner 1939) (=Monotrysia Brnr., 1939 partim; Nannolepidoptera Krkf., 1948) como nombre del suborden y actualización de los Nepticuloidea (Forbes, 1923) de la Península Ibérica (Microlepidoptera). SHILAP Revista de Lepidopterologia 9: 15–28.

Goureau CC (1860) [no title.] (In 'Seance du 22 Février 1860'). Annales de la Société Entomologique de France (3) 8: xxii–xxiv. http://biodiversitylibrary.org/page/30253368

Gregor F, Povolný D (1955) Nové a významné nálezy Lepidopter z Ceskoslovenska. (Neue und interessante Lepidopteren aus der CSR). Casopis Moravského Musea v Brne (Vedy prírodní) 40: 114–129. [In Czech]

Gregor F, Povolny D (1983) Description of *Ectoedemia spiraeae* (Gregor & Povolny, 1955) and designation of type specimens of *Lithocolletis desertella* Gregor & Povolny, 1949. Casopis Moravského Musea v Brne (Vedy prírodní) 68: 173–180.

Grönlien N (1928) Eine neue *Nepticula* on *Betula odorata*. Norsk Entomologisk Tidsskrift 2: 217–219.

Grönlien N (1932) Notes on Norwegian nepticulids. Norsk Entomologisk Tidsskrift 3: 103–114.

Groschke F (1939) Die Kleinschmetterlinge der Grafschaft Glatz (unter besonderer Berücksichtigung des Gebietes der Reinerzer Weistritz und mit Ausnahme der Zygaenidae, Sesiidae, Cossidae, Psychidae und Hepialidae). Mitteilungen der Münchener Entomologischen Verein 29: 643–743. http://www.zobodat.at/pdf/MittMuenchEntGes_029_0643-0734.pdf

Groschke F (1944) Neues über Minierer aus dem Mittelmeergebiet. Mitteilungen der Münchner Entomologischen Gesellschaft 34: 115–124. http://biodiversitylibrary.org/page/29138549

Grossbeck JA (1917) Insects of Florida, 4. Lepidoptera. Bulletin of the American Museum of Natural History 37: 1–147. http://hdl.handle.net/2246/933

Grossenbacher JG (1910) Medullary spots: a contribution to the life history of some cambium miners. New York Agricultural Experiment Station Technical Bulletin 15: 49–65.

Gusenleitner F (1988) Dr. Josef Klimesch: ein Leben im Dienste der Wissenschaft. Stapfia 16: i-xxvi. http://www.zobodat.at/pdf/STAPFIA_0016_0001–0026.pdf

Gustafsson B (1972) Three new species of *Nepticula* from Gambia (Lep. Nepticulidae). Entomologisk Tidskrift 93: 156–159.

Gustafsson B (1973) A new species of *Nepticula* v. Heyden from Rhodes (Lep. Nepticulidae). Entomologisk Tidskrift 94: 197–198.

Gustafsson B (1976) A new species of *Nepticula* v. Heyden and redescription of *Trifurcula clinomochla* (Meyrick) from Sri Lanka (Lep. Nepticulidae). Entomologisk Tidskrift 97: 45–49.

Gustafsson B (1981) New leaf-mining moths of the family Nepticulidae from Cyprus, Greece (Lepidoptera). Entomologica Scandinavica 12: 453–469. doi: 10.1163/187631281X00553

Gustafsson B (1985) New species of *Stigmella* from the Gambia (Lepidoptera, Nepticulidae). Tijdschrift voor Entomologie 127: 165–177. http://biodiversitylibrary.org/page/43079333

Gustafsson B, Nieukerken EJ van (1990) Larvae of Nepticulidae. In: Johansson R, Nielsen ES, Nieukerken EJ van, Gustafsson B (Eds) The Nepticulidae and Opostegidae (Lepidoptera) of NW Europe. Fauna Entomologica Scandinavica 23: 323–356.

Hampson G (1918) Some small families of the Lepidoptera which are not included in the key to the families in the catalogue of Lepidoptera Phalaenae, a list of the families and subfamilies of the lepidoptera with their types and a key to the families. Novitates zoologicae 25: 366–394. doi: 10.5962/bhl.part.29772

Harley B (2001) Obituary. Lieutenant Colonel Arthur Maitland Emmet, MBE, TD, MA, Hon. FRES, FLS. 15 July 1908 - 2 March 2001. Entomologist's Record and Journal of Variation 113: 73–78. http://biodiversitylibrary.org/page/46761155

Hartig F (1941) Neue Microlepidopteren. Mitteilungen der Münchner Entomologischen Gesellschaft 31: 154–163, pl. 158. http://biodiversitylibrary.org/page/28955741

Hartig F (1949) *Stigmella babylonicae* n. sp. von *Salix babylonica* L. Zeitschrift der Wiener Entomologischen Gesellschaft 34: 94–96. http://www.zobodat.at/pdf/ZOEV_34_0094-0096.pdf

Hartig F (1964) *Nepticula vincamajorella* sp. n. (Lepid. Nepticulidae). Bollettino dell'Associazione Romana di Entomologia 19: 8–9, pl. 1.

Hauder F (1912) Beitrag zur Mikrolepidopteren-Fauna Oberösterreichs, 1. Jahresbericht des Museums Francisco-Carolinum, Linz 70: 1–80. http://www.zobodat.at/pdf/JOM_70_0001-0321.pdf

Hausenblas D (2009) Korrekturen und Ergänzungen zur Mikrolepidopterenfauna Baden-Württembergs und angrenzender Gebiete – 2. Beitrag. Mitteilungen des Entomologischen Vereins Stuttgart 44: 81–106. http://www.zobodat.at/pdf/Mitt-Ent-Ver-Stuttgart_44_2009_0081-0107.pdf

Häuser CL, Holstein J, Steiner A (2012) The Global Butterfly Information System. Museum für Naturkunde. http://www.globis.insects-online.de

Haworth AH (1828) Lepidoptera Britannica; sistens digestionem novam Insectorum Lepidopterorum quae in Magna Britannia reperiuntur, larvarum pabulo, temporeque pascendi; expansione alarum; mensibusque volandi; synonymis atque locis observationibusque variis. 4. J. Murray, London: 513–609.

Heinemann H von (1861) Ueber *Nepticula*-Raupen. Correspondenzblatt für Sammler von Insecten, insbesondere von Schmetterlingen 2: 174–175.

Heinemann H von (1862a) Einige Bemerkungen über die Arten der Gattung *Nepticula*. Wiener Entomologische Monatschrift 6: 237–268. http://www.zobodat.at/stable/pdf/WEMS_6_0237-0268.pdf

Heinemann H von (1862b) Einige Bemerkungen über die Arten der Gattung *Nepticula*. Wiener Entomologische Monatschrift 6: 301–320. http://www.zobodat.at/stable/pdf/WEMS_6_0301-0320.pdf

Heinemann H von (1871) Nachtrag zu den Bemerkungen über die Arten der Gattung *Nepticula*. Berliner Entomologische Zeitschrift 15: 209–223. doi: 10.1002/mmnd.18710150224

Heinemann H von, Wocke MF ([1876]) Die Schmetterlinge Deutschlands und der Schweiz. Zweite Abtheilung. Kleinschmetterlinge. Band 2. Die Motten und Federmotten. Heft 2. C. A. Schwetke und Sohn, Braunschweig, 389–825, v-vi, tabelle 381–102 pp. https://archive.org/details/dieschmetterlin01heingoog

Heinrich C (1918) On the lepidopterous genus *Opostega* and its larval affinities. Proceedings of the Entomological Society of Washington 20: 27–38. http://biodiversitylibrary.org/page/2591893

Hemming F (1937) Hübner: a bibliographical and systematic account of the entomological works of Jacob Hübner and of the supplements thereto by Carl Geyer, G.F. von Fröhlich and G.A.W. Herrich-Schäffer, 2 Vols. Royal Entomological Society of London, London, Vol. 1: xxxiv, 605 pp. - Vol. 2: ix, 274 pp.

Heppner JB, Davis DR (2009) A new *Neopostega* from Guatemala (Lepidoptera: Opostegidae). Lepidoptera Novae 2: 31–34. http://entomology.si.edu/StaffPages/Davis/2009_neopostega-guatemala.pdf

Hering M (1931) Minenstudien 12. Zeitschrift für Pflanzenkrankheiten (Pflanzenpathologie) und Pflanzenschutz 41: 529–551. http://www.jstor.org/stable/43526654

Hering M (1932a) Die Schmetterlinge nach ihren Arten dargestellt. Quelle & Meyer, Leipzig, Die Tierwelt Mitteleuropas Ergänzungsband 1: i-vii, 1–545 pp.

Hering M (1932b) Minenstudien 13. Zeitschrift für Pflanzenkrankheiten (Pflanzenpathologie) und Pflanzenschutz 42: 567–579. http://www.jstor.org/stable/43526091

Hering M (1935) Minenstudien 15. Zeitschrift für Pflanzenkrankheiten (Pflanzenpathologie) und Pflanzenschutz 45: 1–15. http://www.jstor.org/stable/43526713

Hering M (1937) Die Blattminen Mittel- und Nord-Europas einschliesslich Englands. Bestimmungstabellen aller von Insektenlarven der verschiedenen Ordnungen erzeugten Minen. Lief. 6. Gustav Feller, Neubrandenburg, 561–631.

Hering EM (1940) *Zimmermannia liebwerdella* (Zimmermann). Ein Nachwort. Mitteilungen aus dem Zoologischen Museum in Berlin 24: 266. doi: 10.1002/mmnz.19390240205

Hering EM (1942) Eine neue *Nepticula* der *subbimaculella*-Gruppe. *Nepticula zimmermanni* spec. nov. von *Quercus pubescens* Willd. Mitteilungen der Deutschen Entomologischen Gesellschaft 11: 26–29.

Hering EM (1943) Untersuchungen über die Weiden-Nepticuliden I. Zeitschrift der Wiener Entomologischen Gesellschaft 28: 273–278. http://www.zobodat.at/pdf/ZOEV_28_0273-0278.pdf

Hering EM (1951) Biology of the leaf miners. Junk, 's-Gravenhage: iv, 420 pp., 2 pls.

Hering EM (1957) Bestimmungstabellen der Blattminnen von Europa, 3 vols. Junk, 's-Gravenhage: 1185 pp.

Herrich-Schäffer GAW (1853) Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text, Revision und Supplement zu Jakob Hübner's Sammlung europäischer Schmetterlinge, 5, Die Schaben und Federmotten. G. J. Manz, Regensburg, Plates Tineides 87, 88, 94, 95, 105–108, 116; Pterophides 104–106. https://books.google.nl/books?id=f2oFAAAAQAAJ&hl=nl&pg=PP15#v=onepage&q&f=false

Herrich-Schäffer GAW (1855a) Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text, Revision und Supplement zu Jakob Hübner's Sammlung europäischer Schmetterlinge, 5, Die Schaben und Federmotten. G. J. Manz, Regensburg, 289–394. http://biodiversitylibrary.org/page/42586913

Herrich-Schäffer GAW (1855b) Systema Lepidopterorum Europae. Systematisches Verzeichniss der Europäischen Schmetterlinge. In: Herrich-Schäffer GAW (Ed.) Systematische Bearbeitung der Schmetterlinge von Europa, zugleich als Text, Revision und Supplement zu Jakob Hübner's Sammlung europäischer Schmetterlinge, 6. G. J. Manz, Regensburg, 61–72.

Herrich-Schäffer GAW (1860) Revision der Europäischen Schmetterlingfauna. Correspondenzblatt für Sammler von Insecten, insbesondere von Schmetterlingen 1: 59–61. https://books.google.nl/books/download/Correspondenzblatt_f%C3%BCr_sammler_von_inse.pdf?id=-eUTAAAAYAAJ&hl=nl&output=pdf&sig=ACfU3U3rk6uPTkP4a2FWEK1NeEhwrhSN9A

Herrich-Schäffer GAW (1863a) Systematisches Verzeichniss [sic!] der Schmetterlinge von Europa. Dritte Auflage; mit Angabe des Vaterlandes. Manz, Regensburg: 28 pp. http://www.mdz-nbn-resolving.de/urn/resolver.pl?urn=urn:nbn:de:bvb:12-bsb10307519-2

Herrich-Schäffer GAW (1863b) Vorläufiger Bericht über einen Ausflug ins Engadin. Correspondenz-Blatt des zoologisch-mineralogischen Vereines in Regensburg 17: 167–175. http://biodiversitylibrary.org/page/11838960

Herrich-Schäffer GAW (1863c) Vorläufiger Bericht über einen Ausflug ins Engadin. Correspondenz-Blatt des zoologisch-mineralogischen Vereines in Regensburg 17: 181–188. http://biodiversitylibrary.org/page/11838974

Heyden C von (1843) Vierte Sitzung, am 23. September. Amtlicher Bericht über die Versammlung der Gesellschaft deutscher Naturforscher und Aerzte zu Mainz 20: 208–209. http://biodiversitylibrary.org/page/10935453

Heyden C von (1858) *Nepticula* of the *Convolvulus*. Entomologist's Weekly Intelligencer 4: 175. http://biodiversitylibrary.org/page/50828211

Heyden C von (1861) Fragmente aus meinen entomologischen Tagebüchern. Stettiner Entomologische Zeitung 22: 31–42. http://biodiversitylibrary.org/page/25455298

Heyden C von (1862) Gliederthiere aus der Braunkohle des Niederrhein's, der Wetterau und der Röhn. Paleontographica 10: 62–82, pl. x. http://publikationen.ub.uni-frankfurt.de/frontdoor/index/index/docId/16069

Heydenreich (1851) Lepidopterorum Europaeorum: catalogus methodicus. Systematisches Verzeichniss der Europaeischen Schmetterlinge. Dritte Ausgabe. Julius Klinkhardt, Leipzig, 130 pp.

Hirano N (2010) Description of five new *Stigmella* and one *Bohemannia* species from Japan, with a new record of one *Bohemannia* species (Lepidoptera, Nepticulidae). Japan Heterocerists' Journal 256: 124–134.

Hirano N (2013) Nepticulidae. In: Hirowatari T, Nasu Y, Sakamaki Y, Kishida Y (Eds) Nihon-san garui hyojun zukan The standard of moths in Japan. Gakken Kyoiku Shuppan, Tokyo, 80–96, pls 3–03 – 3–06. [In Japanese]

Hirano N (2014) Six new and an unrecorded species of *Stigmella* Schrank, 1802 from Japan (Lepidoptera, Nepticulidae). Tinea 23: 19–32.

Hirowatari T (2013) Opostegidae. In: Hirowatari T, Nasu Y, Sakamaki Y, Kishida Y (Eds) Nihon-san garui hyojun zukan The standard of moths in Japan. Gakken Kyoiku Shuppan, Tokyo, 96–97, pl. 3–06. [In Japanese]

Hoare RJB (1998) Systematics of Australian Nepticulid moths (Lepidoptera: Nepticulidae). PhD Thesis, Australian National University, Canberra, 248 pp.

Hoare RJB (2000a) A new genus of primitive Nepticulidae (Lepidoptera) from eastern Australia, with a revised diagnosis of nepticulid subfamilies. Zoological Journal of the Linnean Society 128: 289–317. doi: 10.1006/zjls.1998.0189

Hoare RJB (2000b) Gondwanan Nepticulidae (Lepidoptera)? Systematics and biology of the *Ectoedemia* (*Fomoria*) *vannifera* group. Tijdschrift voor Entomologie 142: 299–316. doi: 10.1163/22119434-99900031

Hoare RJB, Nieukerken EJ van (2013) Phylogeny and host-plant relationships of the Australian Myrtaceae leafmining moth genus *Pectinivalva* (Lepidoptera, Nepticulidae), with new subgenera and species. ZooKeys 278: 1–64. doi: 10.3897/zookeys.278.4743

Hoare RJB, Johansson R, Nieukerken EJ van, Nielsen ES (1997) Australian Nepticulidae (Lepidoptera): redescription of the named species. Entomologica Scandinavica 28: 1–26. doi: 10.1163/187631297X00132

Hoffmann A (1893) Die Entomologie in der christlichen Legende. Stettiner Entomologische Zeitung 54: 214–215. http://biodiversitylibrary.org/page/8932304

Hofmann O (1858) Ueber die Nepticulen. In: Herrich-Schäffer & Hofmann, III. Nachtrag und Berichtigungen zur Lepidopteren-Fauna von Regensburg. Korrespondenz-Blatt des Zoologisch-Mineralogischen Vereines in Regensburg 12: 185–193. http://www.biodiversitylibrary.org/item/44637

Hofmann O (1868) Zwei neue Tineen. Stettiner Entomologische Zeitung 29: 28–31. http://biodiversitylibrary.org/page/33757182

Hübner J (1796-[1836]) Sammlung europäischer Schmetterlinge, 8. Tineae. Augsburg: pls. 1–71, 477 Figs http://www.biodiversitylibrary.org/item/89072#page/2/mode/2up

Hübner J (1816–1826) Verzeichniss bekannter Schmetterlinge. Augsburg: 1–431, 431–472 pp. http://biodiversitylibrary.org/page/33045094

Hudson GV (1938) Edward Meyrick. B.A., F.R.S., 1854–1938. Transactions of the Royal Society of New Zealand 68: 141–142. http://rsnz.natlib.govt.nz/volume/rsnz_68/rsnz_68_02_001340.html

Hudson GV (1939) A supplement to The Butterflies and Moths of New Zealand. Ferguson & Osborn, Wellington: pp. [iv,] 387–481, 10 pls. http://www.bugz.org.nz/WebForms/ResultDetails.aspx?CurrentDoc=4A102474-EF01-4089-A31A-A1FE7E551E52

International Commission on Zoological Nomenclature (1999) International Code of Zoological Nomenclature. Fourth edition. Adopted by the International Union of Biological Sciences. The International Trust for Zoological Nomenclature, London, xxix + 306 pp. http://iczn.org/code

International Commission on Zoological Nomenclature (2012) Amendment of Articles 8, 9, 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication. ZooKeys 219: 1–10. doi: 10.3897/zookeys.219.3944

Isaac NJB, Mallet J, Mace GM (2004) Taxonomic inflation: its influence on macroecology and conservation. Trends in Ecology & Evolution 19: 464–469. doi: 10.1016/j.tree.2004.06.004

Ivinskis P, Nieukerken EJ van, Rimsaite J (2012) *Trifurcula* (*Glaucolepis*) *lituanica* sp. n., an unexpected new stem-miner on *Salvia pratensis* occurring in eastern Europe (Lepidoptera: Nepticulidae). Zootaxa 3570: 41–55. http://www.mapress.com/zootaxa/2012/f/zt03570p055.pdf

Jäckh E (1951) Die fruchtminierenden Arten der Gattung *Nepticula* Heyd. (*Etainia* Beirne) an den vier deutschen Ahornarten (Lep. Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 36: 170–178, pls. 14–16. http://www.zobodat.at/pdf/ZOEV_36_0170-0178.pdf

Janse AJT (1948) Nepticulidae (Stigmellidae). The Moths of South Africa 4: 149–185, plates 182–194.

Jarzembowski EA (1989) A century plus of fossil insects. Proceedings of the Geologists' Association 100: 433–449. doi: 10.1016/S0016-7878(89)80019-7

Joannis Jd (1908) Deux espèces nouvelles de *Nepticula* (Lep.). Bulletin de la Société Entomologique de France 1907: 326–329. http://biostor.org/reference/98156

Joannis J de (1909) Contribution a l'étude des Lépidoptères du Morbihan. Annales de la Société Entomologique de France 77: 688–838, pl. 615, map. http://biodiversitylibrary.org/page/15213670

Joannis J de (1915) Étude synonymique des espèces de Microlépidoptères décrites comme nouvelles par Duponchel. Annales de la Société Entomologique de France 84: 62–164. http://biodiversitylibrary.org/page/8559018

Johansson R (1971) Notes on the Nepticulidae (Lepidoptera) I. A revision of the *Nepticula ruficapitella* group. Entomologica Scandinavica 2: 241–262. doi: 10.1163/187631271X00257

Johansson R, Nielsen ES (1990) Tribus Nepticulini. In: Johansson R, Nielsen ES, Nieukerken EJ van, Gustafsson B (Eds) The Nepticulidae and Opostegidae (Lepidoptera) of NW Europe. Fauna Entomologica Scandinavica 23: 111–238.

Johansson R, Nielsen ES, Nieukerken EJ van, Gustafsson B (1990) The Nepticulidae and Opostegidae (Lepidoptera) of north west Europe. Fauna Entomologica Scandinavica 23 (2 volumes): 1–739.

Jones WJ (1933) The description and biology of *Nepticula braunella* new species (Lepidoptera-Nepticulidae), a species of leaf miner on *Prunus ilicifolia* Walp. and the variety *integrifolia* Sarg. University of California Publications in Entomology 6: 49–78.

Kalkman C (1987) The two sides of a medal. In: Hovenkamp P (Ed.) Systematics and evolution: a matter of diversity. Utrecht University, Utrecht, 13–21.

Karsholt O, Nielsen ES (1976) Systematisk fortegnelse over Danmarks sommerfugle. Scandinavian Science press, Klampenborg: 1–128 pp.

Karsholt O, Nielsen ES (1986) The Lepidoptera described by C.P. Thunberg. Entomologica Scandinavica 16: 433–463. doi: 10.1163/187631285X00388

Karsholt O, Vieira V (2005) Lepidoptera. In: Borges PAV, Cunha R, Gabriel R, Martins AF, Silva L, Vieira V (Eds) Listagem da fauna (Mollusca e Arthropoda) e flora (Bryophyta, Pteridophyta e Spermatophyta) terrestres dos Açores A list of terrestrial fauna (Mollusca and

Arthropoda) and flora (Bryophyta, Pteridophyta and Spermatophyta) from the Azores. Direcção Regional do Ambiente, Governo Regional dos Açores, Horta, Faial, Açores, 207–210, 230. http://www.azoresbioportal.angra.uac.pt/files/publicacoes_Check_List_Azores.pdf

Kasy F (1979) Die Schmetterlingsfauna des Naturschutzgebietes Hackelsberg, Nordburgenland. Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 30 Supplement: 1–27. http://www.zobodat.at/pdf/ZAOE_S1_0001-0044.pdf

Kasy F (1980) Lepidopterologisch-faunistisch bemerkenswerte Neufunde aus Niederösterreich, IX. (7 für Österreich neue Kleinschmetterlingsarten). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 32: 47–48. http://www.zobodat.at/pdf/ZAOE_32_0047-0048.pdf

Kasy F (1983) Die Schmetterlingsfauna des WWF-Naturreservates "Hundsheimer Berge" in Niederösterreich. Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 34 Supplement: 1–48. http://www.zobodat.at/pdf/ZAOE_S2_0001-0048.pdf

Kearfott WD (1908) New North American Tortricidae and Tineina. Journal of the New York Entomological Society 16: 167–188, pl. 163. http://biodiversitylibrary.org/page/33456066

Kemperman TCM, Wilkinson C, Kuroko H, Kumata T (1985) Japanese species of the genus *Stigmella* (Nepticulidae: Lepidoptera). Insecta Matsumurana 32: 1–107. http://hdl.handle.net/2115/9830

Kernbach K (1967) Über die bisher im Pliozän von Willershausen gefundenen Schmetterlings- und Raupenreste. Bericht der Naturhistorischen Gesellschaft zu Hannover 111: 103–108.

Kirby WF (1897) A hand-book to the order Lepidoptera, 5. Moths, part III. Allen & Co., London, Allen's naturalist's library: xii, 332 pp., pls. 127–158.

Klimesch J (1936) Ein kleiner Beitrag zur Kenntnis der Nepticulidenfauna (Lep.) des Vintschgaues (Prov. Bozen). Stettiner Entomologische Zeitung 97: 194–211.

Klimesch J (1937) *Nepticula loranthella* spec. nov. (Lep. Nepticulidae). Zeitschrift des Österreichischen Entomologen-Vereins 22: 33–35. http://www.zobodat.at/pdf/ZOEV_22_0033-0035.pdf

Klimesch J (1940a) Beschreibung einiger neuer *Nepticula*-Arten (Lep., Nepticulidae). Zeitschrift des Wiener Entomologen-Vereines 25: 79–81, pls 14–15. http://www.zobodat.at/pdf/ZOEV_25_0079-0081.pdf

Klimesch J (1940b) Beschreibung einiger neuer *Nepticula*-Arten (Lep., Nepticulidae). Zeitschrift des Wiener Entomologen-Vereines 25: 89–94. http://www.zobodat.at/pdf/ZOEV_25_0089-0094.pdf

Klimesch J (1940c) Über eine Nepticuliden-Ausbeute von Triest sowie Beschreibung der Mine von *Leucospilapteryx cupediella* H.S. (Lep. Nepticulidae et Gracilariidae). Zeitschrift des Wiener Entomologen-Vereines 25: 176–179, pl. 121. http://www.zobodat.at/pdf/ZOEV_25_0176-0179.pdf

Klimesch J (1940d) Über eine Nepticuliden-Ausbeute von Triest sowie Beschreibung der Mine von *Leucospilapteryx cupediella* H.S. (Lep. Nepticulidae et Gracilariidae). Zeitschrift des Wiener Entomologen-Vereines 25: 188–193, pl. 121. http://www.zobodat.at/pdf/ZOEV_25_0188-0193.pdf

Klimesch J (1941) *Nepticula preisseckeri* spec. nov. (Lep., Nepticulidae). Zeitschrift des Wiener Entomologen-Vereines 26: 162–168, pl. 116. http://www.zobodat.at/pdf/ZOEV_26_0162-0168.pdf

Klimesch J (1948a) Neue *Stigmella*-Arten (Lep., Stigmellidae). Zeitschrift der Wiener Entomologischen Gesellschaft 31: 160–172, 161 pl. http://www.zobodat.at/pdf/ZOEV_31_0160-0172.pdf

Klimesch J (1948b) Zur Frage der verwantschaftlichen Beziehungen einiger *Stigmella*-Arten auf Grund des Baues des männl. Kopulationsapparates. (Lep., Stigmellidae). Zeitschrift der Wiener Entomologischen Gesellschaft 33: 49–82. http://www.zobodat.at/pdf/ZOEV_33_0049-0082.pdf

Klimesch J (1950a) Über einige Nahrungsrassen von *Nepticula*-Arten (Lep., Nepticulidae). Wiener Entomologische Rundschau 2: 25–28. http://www.zobodat.at/pdf/ZAOE_2_0025-0028.pdf

Klimesch J (1950b) Über einige Nahrungsrassen von *Nepticula*-Arten (Lep., Nepticulidae). Entomologisches Nachrichtenblatt Österreichischer und Schweizer Entomologen 2: 72–74. http://www.zobodat.at/pdf/ZAOE_2_0072-0074.pdf

Klimesch J (1951a) Contributo alla fauna Lepidotterologica del Trentino. Studi Trentini di Scienze Naturali 27: 11–68.

Klimesch J (1951b) *Nepticula* (*Levarchama*) *ortneri* spec. nov. (Lept., Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 36: 66–70. http://www.zobodat.at/pdf/ZOEV_36_0066-0070.pdf

Klimesch J (1951c) *Nepticula arbusculae* spec. nov. (Lep., Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 36: 149–152, pl. 110. http://www.zobodat.at/pdf/ZOEV_36_0149-0152.pdf

Klimesch J (1951d) Nuova specie di *Nepticula* della Romagna (Lep., Nepticulidae). *Nepticula Zangherii* spec. nov. Bollettino della Società Entomologica Italiana 81: 61–62.

Klimesch J (1953a) Die europäischen *Trifurcula*- und *Ectoedemia*-Arten (Lep., Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 38: 160–170. http://www.zobodat.at/pdf/ZOEV_38_0160-0170.pdf

Klimesch J (1953b) Die europäischen *Trifurcula*- und *Ectoedemia*-Arten (Lep., Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 38: 191–196. http://www.zobodat.at/pdf/ZOEV_38_0191-0196.pdf

Klimesch J (1956) *Nepticula szőcsi* spec. nov. (Lepidoptera). Annales Historico-Naturales Musei Nationalis Hungarici 48 (series nova 7): 423–424. http://annales.nhmus.hu/cikkreszletes.php?idhoz=733

Klimesch J (1961) Ordnung Lepidoptera. 1. Teil. Pyralidina, Tortricina, Tineina, Eriocraniina und Micropterygina. In: Franz H (Ed) Die Nordost-Alpen im Spiegel ihrer Landtierwelt. Universitätsverlag Wagner, Innsbruck, 481–789.

Klimesch J (1972) *Dechtiria nigrifasciata* (Wlsm.) (Lep., Nepticulidae). Ein Beitrag zur Genitalmorphologie und Ökologie. Mitteilungen der Münchner Entomologischen Gesellschaft 62: 1–12. http://biodiversitylibrary.org/page/29012537

Klimesch J (1975a) Ergebnisse von Untersuchungen einiger Nepticuliden-Typen der Sammlung des Muséum national d'Histoire naturelle, Paris (Lep., Nepticulidae). Bulletin du Muséum National d'Histoire Naturelle 3e Série, Zoologie 221: 861–866.

Klimesch J (1975b) Die an Ulmen lebenden europäischen Nepticuliden-Arten (Lepidoptera: Nepticulidae). Opuscula Entomologica 135: 1–15.

Klimesch J (1975c) Über neue mediterrane und kanarische Nepticuliden (Lep., Nepticulidae). Mitteilungen der Münchner Entomologischen Gesellschaft 65: 1–28. http://biostor.org/reference/98578

Klimesch J (1976) Zur Genitalmorphologie und Lebensweise der *Fedalmia* (*Stigmella*) *saturejae* (Parenti, 1963) (Lep., Nepticulidae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 28: 45–48. http://www.zobodat.at/pdf/ZAOE_28_0045-0048.pdf

Klimesch J (1977) Beiträge zur Kenntnis der Microlepidopteren-Fauna des Kanarischen Archipels. 1. Beitrag: Nepticulidae, Tischeriidae. Vieraea 6: 191–218.

Klimesch J (1978a) *Fomoria luisae* n. sp. (Lep., Nepticulidae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 29: 89–91. http://www.zobodat.at/pdf/ZAOE_29_0089-0091.pdf

Klimesch J (1978b) Beitrag zur Kenntnis der Nepticulidenfauna von Anatolien und der Insel Rhodos (Lepidoptera, Nepticulidae). Tijdschrift voor Entomologie 121: 239–278. http://biostor.org/reference/49889

Klimesch J (1979) Beiträge zur Kenntnis der Nepticuliden. Beschreibung zweier neuer Arten (*Stigmella abaiella* n. sp. und *Trifurcula* (*Fedalmia*) *sanctibenedicti* n. sp.) (Lepidoptera, Monotrysia). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 31: 21–27. http://www.zobodat.at/pdf/ZAOE_31_0021-0027.pdf

Klimesch J (1981) Beiträge zur Kenntris der Nepticulidae (Lep., Monotrysia) 1. Die Futterpflanzen der Nahrungsrassen der *Stigmella aurella* (F.) in Oberösterreich. 2. *Stigmella tormentillella* (H.S.) und verwandte Arten in den Ostalpen. Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 32: 113–128. http://www.zobodat.at/pdf/ZAOE_32_0113-0128.pdf

Koçak AÖ (1981) On the nomenclature of some genera of Lepidoptera. Priamus 1: 97–109.

Koçak AÖ, Kemal M (2007) Replacement names for some Microlepidoptera (Lepidoptera). Miscellaneous Papers Centre for Entomological Studies 105: 5–7. https://archive.org/stream/CentreForEntomologicalStudiesAnkaraMiscellaneousPapers105

Kollar V (1832) Systematisches Verzeichniss der Schmetterlinge im Erzherzogthume Oesterreich. Beiträge zur Landeskunde Oesterreich's unter der Enns 2: 1–101. https://books.google.nl/books?id=3_GAAAAAIAAJ&hl=nl&pg=PA1#v=onepage&q&f=false

Kozlov MV (1985) Novye i maloizvestnye Opostegidy (Lepidoptera, Opostegidae) iz Aziatskoj chasti SSSR. (New and little known opostegid moths (Lepidoptera, Opostegidae) from Asiatic parts of the USSR). Trudy Zoologicheskogo Instituta 134: 49–58. [In Russian]

Kozlov MV (1987) Morfologiya babochek i filogeneticheskie svyazi semejstv Opostegidae, Nepticulidae i Tischeriidae (Lepidoptera, Nepticulomorpha). Imaginal morphology and the phylogeny of the moth families Opostegidae, Nepticulidae and Tischeriidae (Lepidoptera, Nepticulomorpha). Zoologicheskii Zhurnal 66: 847–859. [In Russian]

Kozlov MV (1988) Paleontologiya Cheshuekrylykh i voprosy filogenii otryada Papilionida [Palaeontology of lepidopterans and problems in Papilionida order phylogeny]. In: Ponomarenko AG (Ed.) Melovoj biotsenoticheskij krizis i evolutsiya nasekomykh [Cretaceous biocoenotic crisis and insect evolution]. Nauka, Moscow, 16–69, references 216–228. [In Russian] http://nepticuloidea.info/content/paleontologiya-cheshuekrylykh-i-voprosy-filogenii-otryada-papilionida-palaeontology

Kraatz G (1875) Necrologie und Todesanzeigungen. Deutsche Entomologische Zeitschrift 19: 436–437, frontispiece.

Krassilov V (2008) Traumas on fossil leaves from the Cretaceous of Israel. In: Krassilov V, Rasnitsyn A (Eds) Plant-arthropod interactions in the early angiosperm history: evidence from the Cretaceous of Israel. Pensoft Publishers & Brill, Sofia & Leiden, 7–187. doi: 10.1163/ej.9789004170711.1-229.4

Krenek V (2000) Small moths of Europe. Ceský Tesin: 174 pp.

Krogerus H, Opheim M, Schantz M von, Svensson I, Wolff N (1971) Catalogus Lepidopterorum Fenniae et Scandinaviae, Microlepidoptera. Helsigin Hyönteisvaihtoyhdistys, Helsinki, 1–40.

Kumata T, Nakatani M (1995) Discovery of three lepidopterous leaf-miners on *Vaccinium vitis-idaea* from Hokkaido, Japan (Nepticulidae, Gracillariidae and Coleophoridae). Transactions of the Lepidopterological Society of Japan 46: 91–102.

Kuroko H (1978) A new species of Nepticulid moth from Japan (Lepidoptera: Nepticulidae). Akitu, new series 18: 1–5.

Kuroko H (1982a) Nepticuloidea: Nepticulidae, Opostegidae. In: Inoue H, Sugi S, Kuroko H, Moriuti S, Kawabe A (Eds) Moths of Japan 1 Text. Kodansha, Tokyo, 49–50. [In Japanese]

Kuroko H (1982b) Nepticulidae, Opostegidae. In: Inoue H, Sugi S, Kuroko H, Moriuti S, Kawabe A (Eds) Moths of Japan 2 Plates and synonymic catalogue. Kodansha, Tokyo, 155, pls 151, 230, 261–262. [In Japanese]

Kuroko H (1989) Lepidopterous Leaf Miners of Japan (1). Kita-Kyushu no Konchu 36: 73–74, 71 pl. [In Japanese]

Kuroko H (1990) Lepidopterous Leaf Miners of Japan (2). Kita-Kyushu no Konchu 37: 1–2, 1 pl. [In Japanese]

Kuroko H (1999) Lepidopterous Leaf Miners of Japan (3). Kita-Kyushu no Konchu 46: 79–88. [In Japanese]

Kuroko H (2004) A new species of the genus *Stigmella* Schrank (Lepidoptera, Nepticulidae) from Japan. Tinea 18: 238–240.

Larsen CS (1927) Upon some new and little-known species and aberrations of Microlepidoptera from Denmark. Entomologiske Meddelelser 17: 1–6, pl. 1.

Laštůvka A, Huemer P (2002) *Stigmella gutlebiella* sp. n., eine neue Schmetterlingsart aus dem Iran (Lepidoptera, Nepticulidae). Carinthia II 192/112: 603–606. http://www.zobodat.at/pdf/CAR_192_112_0603-0606.pdf

Laštůvka A, Laštůvka Z (1990) *Stigmella inopinata* sp. n. und *Parornix mixta* (Triberti) in Mitteleuropa (Lepidoptera). Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 38: 197–203.

Laštůvka A, Laštůvka Z (1993) *Acalyptris pyrenaica* sp. n. aus Spanien (Lepidoptera: Nepticulidae). SHILAP Revista de Lepidopterologia 21: 157–160.

Laštůvka A, Laštůvka Z (1997) Nepticulidae Mitteleuropas. Ein illustrierter Begleiter (Lepidoptera). Konvoj, Brno: 229 pp.

Laštůvka A, Laštůvka Z (2000a) Zwei neue *Globularia* minierende *Trifurcula*-Arten (Lepidoptera, Nepticulidae). Acta Musei Moraviae, Scientiae biologicae 85: 289–296.

Laštůvka A, Laštůvka Z (2000b) The *Ectoedemia* (*Fomoria*) *weaveri*-group of the southern Balkans with a description of two new species (Lepidoptera, Nepticulidae). Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 48: 19–26.

Laštůvka A, Laštůvka Z (2005a) Four new *Trifurcula* species and additional faunal data on Nepticulidae from Italy (Lepidoptera: Nepticulidae). Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 53: 7–14. doi: 10.11118/actaun200553010007

Laštůvka A, Laštůvka Z (2005b) *Parafomoria fumanae* sp. n., a new stem miner on *Fumana procumbens* (Lepidoptera: Nepticulidae). Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 53: 15–17. doi: 10.11118/actaun200553010015

Laštůvka Z, Laštůvka A (1994) Drei neue Arten der *Trifurcula pallidella*-Gruppe aus Mitteleuropa (Lepidoptera: Nepticulidae). Entomologia Generalis 18: 201–212.

Laštůvka Z, Laštůvka A (1998) Beitrag zur Kenntnis der Nepticulidenfauna Griechenlands (Lepidoptera, Nepticulidae). Stapfia 55: 313–326. http://www.zobodat.at/pdf/STAPFIA_0055_0313-0326.pdf

Laštůvka Z, Laštůvka A (2007) Seven new species of the subgenus *Glaucolepis* Braun from southern Europe (Lepidoptera: Nepticulidae, *Trifurcula*). Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 55: 101–110. doi: 10.11118/actaun200755050101

Laštůvka Z, Laštůvka A, Nieukerken EJ van (2013) The *Bupleurum* (Apiaceae) feeding species of *Trifurcula* (*Glaucolepis*): new species, biology and distribution (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 156: 191–210. doi: 10.1163/22119434-00002028

Le Marchand S (1937) *Nepticula alaternella* nova species, du midi de la France. Amateur de Papillons 8: 233–239.

Le Marchand S (1944) La biologie de *Stigmella* (*Nepticula*) *sericopeza* Z. pose un curieux problème. Revue française de Lépidoptérologie 9: 351–358.

Le Marchand S (1946a) Bonnes localités: Saint-Palais-sur-Mer (Charente-Maritime). Revue française de Lépidoptérologie 10: 200–217.

Le Marchand S (1946b) *Stigmella prinophyllella* n. sp. (Microlépidoptères, Stigmellidae). Revue française de Lépidoptérologie 10: 280–289.

Le Marchand S (1948) *Stigmella* (*Nepticula*) *prinophyllella* Le Md. ets synonyme de *Nepticula haraldi* Soffner. Revue française de Lépidoptérologie 11: 298–299.

Legrand H (1965) Lépidoptères des îles Seychelles et d'Aldabra. Mémoires du Musée National d'Histoire Naturelle, Série A, Zoologie 37: 1–210, pls. i-xvi.

Leraut P (1980) Liste systématique et synonymique des Lépidoptères de France, Belgique et Corse. Supplément à Alexanor et au Bulletin de la Société Entomologique de France, 1–334.

Leraut P (1997) Liste systématique et synonymique des Lépidoptères de France, Belgique et Corse (deuxième édition). Supplément à Alexanor, Paris, 526 pp.

Lewis JA, Sohn J-C (2015) Lepidoptera: Yponomeutoidea I (Argyresthiidae, Attevidae, Praydidae, Scythropiidae, and Yponomeutidae). Brill, Leiden, World Catalogue of Insects 12: xxiv, 253 pp.

Lhomme L (1945) Deux espèces nouvelles pour la faune française *Phlyctaenodes ustrinalis* Chris. (Pyraustinae) et *Stigmella (Nepticula) mahalebella* Klim. (Stigmellina). Revue française de Lépidoptérologie 10: 155–156.

Lhomme L ([1963]) Catalogue des Lépidoptères de France et de Belgique 2, Microlépidoptères. [Fascicule 8.]. Douelle (Lot), 2(8): 1081–1253.

Linnaeus C (1767) Systema naturae. Tom. I. Pars II. Editio duodecima reformata. Holmiae: 533–1328 +[1336] pp. doi: 10.5962/bhl.title.68927

Mallet J (2001) Species, concepts of. In: Levin SA (Ed.) Encyclopedia of Biodiversity 5: 427–440. Academic Press. http://www.ucl.ac.uk/taxome/jim/Sp/Sub-semi.pdf

Mann J (1855) Die Lepidopteren, gesammelt auf einer entomologischen Reise in Corsika im Jahre 1855. Verhandlungen der Zoologisch-Botanischen Vereins in Wien 5: 529–612. http://biodiversitylibrary.org/page/16402230

Mariani M (1937) Nuove specie e forme di Lepidotteri di Sicilia ed un nuovo parassita degli agrumi. Giornale di Scienze Naturali ed Economiche Palermo 34: 1–13, pl.

Mariani M (1939) Un nuove genere e due nuove specie di Lepidotteri di Sicilia. Giornale di Scienze Naturali ed Economiche di Palermo 40: 1–7.

Martins-Neto RG (1989) Novos insetos terciários do estado de São Paulo. Revista Brasileira de Geociências 19: 375–386.

Matsumura S (1931) 6000 Illustrated insects of Japan-Empire. Tokyo: x, ii, iii, iii, 23, 1497 + 191, pp., 10 pls.

McDunnough J (1939) Check list of the Lepidoptera of Canada and the United States of America, 2. Microlepidoptera. Memoirs of the Southern California Academy of Sciences 2: 1–171.

Meess A (1907) Nachtrag zu Reutti`s Lepidopteren-Fauna des Grossherzogtums Baden. Mitteilungen des Badischen Zoologischen Vereins 18: 121–130.

Mendes C (1910a) *Nepticula* et *Coleophora* novae ex Lusitania. Brotéria Serie Zoológica 9: 102–104, pl. VI. http://biodiversitylibrary.org/page/46580177

Mendes C (1910b) *Lithocolletes* et *Nepticulae* novae ex Lusitania. Brotéria Serie Zoológica 9: 163–166, pl. 167.

Mendes C (1918) Lepidopteros de Salamanca. Brotéria Serie Zoológica 16: 111–129.

Menken SBJ (1990) Biochemical systematics of the leaf mining moth family Nepticulidae (Lepidoptera). 3. Allozyme variation patterns in the *Ectoedemia subbimaculella* group. Bijdragen tot de Dierkunde 60: 189–197.

Menken SBJ, Brouwer J (1984) Biochemical systematics and speciation in the leaf-mining genus *Ectoedemia* Busck (Nepticulidae, Lepidoptera). International Congress of Entomology, Proceedings 17: 10.

Menken SBJ, Wiebosch Steeman M (1988) Clonal diversity, population structure, and dispersal in the parthenogenetic moth *Ectoedemia argyropeza*. Entomologia Experimentalis et Applicata 49: 141–152. doi: 10.1111/j.1570-7458.1988.tb02485.x

Menken SBJ, Boomsma JJ, Nieukerken EJ van (2010) Large-scale evolutionary patterns of host plant associations in the Lepidoptera. Evolution 64: 1098–1119. doi: 10.1111/j.1558-5646.2009.00889.x

Mey W (2004) Nepticulidae (Lepidoptera, Nepticuloidea). In: Mey W (Ed.) The Lepidoptera of the Brandberg Massif in Namibia Part 1. Esperiana Memoir 1: 27–38, plate 22.

Meyrick E (1880) Descriptions of Australian Micro-Lepidoptera. Proceedings of the Linnean Society of New South Wales 5: 132–182. doi: 10.5962/bhl.part.15874

Meyrick E (1889) Descriptions of New Zealand Micro-Lepidoptera. Transactions and Proceedings of the New Zealand Institute 21: 154–188. http://rsnz.natlib.govt.nz/volume/rsnz_21/rsnz_21_00_001620.pdf

Meyrick E (1893) Descriptions of Australian Microlepidoptera, XVI. Tineidae. Proceedings of the Linnean Society of New South Wales (2) 7: 477–612. doi: 10.5962/bhl.part.26071

Meyrick E (1895) A handbook of British Lepidoptera. Macmillan & Co, London, vi+843 pp. doi: 10.5962/bhl.title.8019

Meyrick E (1905) Descriptions of Indian Micro-Lepidoptera, 1. Journal of the Bombay Natural History Society 16: 580–619. http://www.biodiversitylibrary.org/part/13703

Meyrick E (1906a) Descriptions of Indian Micro-Lepidoptera, 2. Journal of the Bombay Natural History Society 17: 403–417. http://biodiversitylibrary.org/page/30119593

Meyrick E (1906b) Descriptions of Australian Tineina. Transactions of the Royal Society of South Australia 30: 33–66. http://www.biodiversitylibrary.org/part/79212

Meyrick E (1907) Descriptions of Indian Micro-Lepidoptera, 4. Journal of the Bombay Natural History Society 17: 976–994. http://www.biodiversitylibrary.org/part/68694

Meyrick E (1910) Notes and descriptions of Indian Micro-Lepidoptera. Records of the Indian Museum 5: 217–232. 10.5962/bhl.part.10499

Meyrick E (1911a) Descriptions of Transvaal Micro-Lepidoptera, 2. Annals of the Transvaal Museum 2: 218–240. http://biodiversitylibrary.org/page/50495141

Meyrick E (1911b) Descriptions of South African Micro-Lepidoptera, 3. Annals of the Transvaal Museum 3: 63–83. http://biodiversitylibrary.org/page/50476969

Meyrick E (1911c) Descriptions of Indian Micro-Lepidoptera, 14. Journal of the Bombay Natural History Society 21: 104–131. http://biodiversitylibrary.org/page/30150775

Meyrick E (1913) Descriptions of South African Micro-Lepidoptera, 4. Annals of the Transvaal Museum 3: 267–336. http://biodiversitylibrary.org/page/50477201

Meyrick E (1914) Descriptions of South African Micro-Lepidoptera, 5. Annals of the Transvaal Museum 4: 187–205. http://biodiversitylibrary.org/page/50439657

Meyrick E (1915a) Descriptions of South American Micro-Lepidoptera. Transactions of the Entomological Society of London 1915: 201–256. 10.1111/j.1365-2311.1915.tb02527.x

Meyrick E (1915b) Exotic Microlepidoptera 1(11): 321–352. http://biodiversitylibrary.org/page/10440213

Meyrick E (1916a) Exotic Microlepidoptera 1(20): 609–640. http://biodiversitylibrary.org/page/10440537

Meyrick E (1916b) Exotic Microlepidoptera 2(1): 1–32. http://biodiversitylibrary.org/page/9808402

Meyrick E (1916c) Descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 48: 414–419. http://rsnz.natlib.govt.nz/volume/rsnz_48/rsnz_48_00_004630.pdf

Meyrick E (1917a) Descriptions of South African Micro-Lepidoptera. Annals of the South African Museum 17: 1–21. http://biodiversitylibrary.org/page/1520692

Meyrick E (1917b) Descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 49: 245–247. http://rsnz.natlib.govt.nz/volume/rsnz_49/rsnz_49_00_004430.pdf

Meyrick E (1918a) Descriptions of South African Micro-Lepidoptera. Annals of the Transvaal Museum 6: 7–59. http://biodiversitylibrary.org/page/50361285

Meyrick E (1918b) Exotic Microlepidoptera 2(6): 161–192. http://biodiversitylibrary.org/page/9808551

Meyrick E (1919) Descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 51: 349–354. http://rsnz.natlib.govt.nz/volume/rsnz_51/rsnz_51_00_003990.pdf

Meyrick E (1920a) Descriptions of South African Micro-Lepidoptera. Annals of the South African Museum 17: 273–318. doi: 10.5962/bhl.part.22317

Meyrick E (1920b) Insectes Lépidoptères, 2. Microlepidoptera. Résultats Scientifiques Voyage de Ch Alluaud et R Jeannel en Afrique Orientale (1911–1912). A. Lhomme, Paris, 33–120.

Meyrick E (1920c) Exotic Microlepidoptera 2(12): 353–384. http://biodiversitylibrary.org/page/9808767

Meyrick E (1921a) Exotic Microlepidoptera 2(13): 385–416. http://biodiversitylibrary.org/page/9808803

Meyrick E (1921b) Descriptions of South African Micro-Lepidoptera. Annals of the Transvaal Museum 8: 49–148. http://biodiversitylibrary.org/page/50414570

Meyrick E (1921c) Notes and descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 53: 334–336. http://rsnz.natlib.govt.nz/volume/rsnz_53/rsnz_53_00_004360.pdf

Meyrick E (1921d) Exotic Microlepidoptera 2(15): 449–480. http://biodiversitylibrary.org/page/9808873

Meyrick E (1922) Exotic Microlepidoptera 2(18): 545–576. http://biodiversitylibrary.org/page/9808979

Meyrick E (1924a) Descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 55: 661–662. http://rsnz.natlib.govt.nz/volume/rsnz_55/rsnz_55_00_006360.pdf

Meyrick E (1924b) Exotic Microlepidoptera 3(3): 65–96. http://biodiversitylibrary.org/page/38654136

Meyrick E (1926) Exotic Microlepidoptera 3(10): 289–320. http://biodiversitylibrary.org/page/38654360

Meyrick E (1928a) Exotic Microlepidoptera 3(13): 385–416. http://biodiversitylibrary.org/page/38654456

Meyrick E (1928b) Exotic Microlepidoptera 3(15): 449–480. http://biodiversitylibrary.org/page/38654520

Meyrick E (1930) Exotic Microlepidoptera 4(1): 1–32.

Meyrick E (1931a) Micro-Lepidoptera from south Chile and Argentina. Anales del Museo Nacional de Historia Natural de Buenos Aires 36: 377–415.

Meyrick E (1931b) Exotic Microlepidoptera 4(6): 161–192.

Meyrick E (1932) Exotic Microlepidoptera 4(10): 289–320.

Meyrick E (1934a) Exotic Microlepidoptera 4(15): 449–480.

Meyrick E (1934b) Exotic Microlepidoptera 4(17): 513–544.

Meyrick E (1935) Exotic Microlepidoptera 4(18): 545–576.

Meyrick E (1937) Exotic Microlepidoptera 5(3): 65–96.

Miller SE, Hodges RW (1990) Primary types of microlepidoptera in the Museum of Comparative Zoology (with a discursion[sic] on V.T. Chambers' work). Bulletin of the Museum of Comparative Zoology 152: 45–87. http://www.biodiversitylibrary.org/page/4781870

Millière P (1876) Catalogue raisonné des Lépidoptères des Alpes-Maritimes. Troisième et dernière partie. Imprimerie H. Vidal, Cannes, pp. [249]-455, pls. I, II.

Millière P (1886) Catalogue raisonné des Lépidoptères des Alpes-Maritimes (2e supplément). Naturalista Siciliano 5: 195–204. http://biodiversitylibrary.org/page/30388424

Minet J (1990) Description préliminaire d'*Ectoedemia scoblei* n. sp., taxon representant une famille nouvelle pour la faune malgache (Lep. Nepticulidae). Bulletin de la Société Entomologique de France 94.

Minet J (2004) Un nom de remplacement pour le premier Nepticulidae signalé de Madagascar (Lep., Nepticulina). Bulletin de la Société Entomologique de France 109: 366.

Müller-Rutz J (1934a) Über Microlepidopteren. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 16: 118–128, 111 pl. doi: 10.5169/seals-400814

Müller-Rutz J (1934b) Berichtigung. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 16: slip. doi: 10.5169/seals-400814. [last page]

Mutanen M, Hausmann A, Hebert PDN, Landry J-F, de Waard JR, Huemer P (2012) Allopatry as a gordian knot for taxonomists: patterns of DNA barcode divergence in arctic-alpine Lepidoptera. PLoS ONE 7: e47214. doi: 10.1371/journal.pone.0047214

Mutanen M, Kivelä SM, Vos RA, Doorenweerd C, Ratnasingham S, Hausmann A, Huemer P, Dincă V, Nieukerken EJ van, Lopez-Vaamonde C, Vila R, Aarvik L, Decaëns T, Efetov KA, Hebert PDN, Johnsen A, Karsholt O, Pentinsaari M, Rougerie R, Segerer A, Tarmann G, Zahiri R, Godfray HCJ (2016) Species-level para- and polyphyly in DNA barcode gene trees: strong operational bias in European Lepidoptera. Systematic Biology early view. doi: 10.1093/sysbio/syw044

Navickaitė A, Diškus A, Stonis J (2011) Review of Nepticulidae (Insecta: Lepidoptera) occurring in the Curonian Spit (Baltic Coast of Lithuania). Acta Zoologica Lituanica 21: 221-231. doi: 10.2478/v10043-011-0023-1

Navickaitė A, Diškus A, Stonis JR (2014) An updated checklist of Nepticulidae (Lepidoptera) of the Crimea, Sub-Mediterranean SE Europe. Zootaxa 3847: 151–202. doi: 10.11646/zootaxa.3847.2.1

Nel J (2012) Une nouvelle espèce du sous-genre *Glaucolepis* Braun, 1917, *Trifurcula collinella* n. sp., dans le Var (Lep. Nepticulidae). Oreina 19: 24–25.

Nel J, Varenne T (2012) *Pseudopostega cyrneochalcopepla* n. sp., *Monochroa cyrneogonella* n. sp., *Syncopacma cinctelloides* n. sp., espèces nouvelles découvertes en Corse (Lep. Opostegidae, Gelechiidae). Oreina 17: 11–13.

Nemeş I (1970) *Stigmella* (*Fomoria*) *niculescui* nova species (Lep. Stigmellidae). Bulletin de la Société Entomologique de Mulhouse 1970: 33–35.

Nemeş I (1972) *Stigmella* (*Fomoria*) *peiuii* nova species (Lepidoptera, Stigmellidae). Studii şi Comunicări, Muzeul de Ştiinţele naturii Bacău 5: 153–156.

Newton PJ, Wilkinson C (1982) A taxonomic revision of the North American species of *Stigmella* (Lepidoptera: Nepticulidae). Systematic Entomology 7: 367–463. 10.1111/j.1365-3113.1982.tb00455.x

Nickerl O (1908) Die Motten Böhmens (Tineen). Beiträge zur Insekten- Fauna Böhmens VI. Verlag der Gesellschaft für Physiokratie, Prag, 161 pp. http://www.zobodat.at/pdf/Beitr-Insektenfauna-Boehmens_6_0001-0161.pdf

Nielsen ES (1980) Entomology. (The Danish scientific expedition to Patagonia and Tierra del Fuego 1978–1979). Geografisk Tidsskrift 80: 9–13.

Nielsen ES (1985) Primitive (non-ditrysian) Lepidoptera of the Andes: diversity, distribution, biology and phylogenetic relationships. Journal of Research on the Lepidoptera, Supplement 1: 1–16. http://lepidopteraresearchfoundation.org/journals/23/PDF23/2nd_symp_neot_lepid_1983_1_16.pdf

Nielsen ES (1996) Nepticulidae. In: Nielsen ES, Edwards ED, Rangsi TV (Eds) Checklist of the Lepidoptera of Australia. CSIRO, Canberra, 27.

Nieukerken EJ van (1982) New and rare Nepticulidae in the Netherlands (Lepidoptera). Entomologische Berichten, Amsterdam 42: 104–112. http://www.repository.naturalis.nl/document/50157

Nieukerken EJ van (1983a) The identity of *Stigmella zelleriella* (Snellen, 1875), a senior synonym of *S. repentiella* (Wolff, 1955) (Lepidoptera: Nepticulidae). Zoologische Mededelingen (Leiden) 57: 59–65. http://www.repository.naturalis.nl/document/150486

Nieukerken EJ van (1983b) The Cistaceae-feeding Nepticulidae (Lepidoptera) of the western Palaearctic region. Systematic Entomology 8: 453–478. doi: 10.1111/j.1365–3113.1983.tb00495.x

Nieukerken EJ van (1985a) A new species of *Parafomoria* van Nieukerken, and some additional notes on the genus (Lepidoptera: Nepticulidae). Entomologische Berichten, Amsterdam 45: 24–28. http://www.repository.naturalis.nl/document/50160

Nieukerken EJ van (1985b) A taxonomic revision of the western Palaearctic species of the subgenera *Zimmermannia* Hering and *Ectoedemia* Busck s. str. (Lepidoptera, Nepticulidae), with notes on their phylogeny. Tijdschrift voor Entomologie 128: 1–164. http://biodiversitylibrary.org/part/69300

Nieukerken EJ van (1986a) A provisional phylogenetic check-list of the western Palaearctic Nepticulidae, with data on hostplants (Lepidoptera). Entomologica Scandinavica 17: 1–27. doi: 10.1163/187631286X00099

Nieukerken EJ van (1986b) Systematics and phylogeny of Holarctic genera of Nepticulidae (Lepidoptera, Heteroneura: Monotrysia). Zoologische Verhandelingen 236: 1–93. http://www.repository.naturalis.nl/record/317599

Nieukerken EJ van (1987a) Identity of two Nepticulidae described by A. Caradja (Lepidoptera). Zoologische Mededelingen (Leiden) 61: 141–145. http://www.repository.naturalis.nl/record/227183

Nieukerken EJ van (1987b) Taxonomy and distribution of *Trifurcula squamatella* Stainton sp. rev., a senior synonym of *T. maxima* Klimesch (Lepidoptera: Nepticulidae). Entomologist's Gazette 38: 179–187. http://www.repository.naturalis.nl/document/50170

Nieukerken EJ van (1990a) Opostegidae. In: Johansson R, Nielsen ES, Nieukerken EJ van, Gustafsson B (Eds) The Nepticulidae and Opostegidae (Lepidoptera) of NW Europe. Fauna Entomologica Scandinavica 23: 357–372.

Nieukerken EJ van (1990b) The *Trifurcula subnitidella* group (Lepidoptera: Nepticulidae): taxonomy, distribution and biology. Tijdschrift voor Entomologie 133: 205–238. http://biodiversitylibrary.org/part/66032

Nieukerken EJ van (1990c) *Stigmella rolandi* sp. n.: a widespread southern European species on Rosa (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 133: 239–243. http://biodiversitylibrary.org/part/66033

Nieukerken EJ van (1996) Nepticulidae, Opostegidae. In: Karsholt O, Razowski J (Eds) The Lepidoptera of Europe A distributional checklist. Apollo Books, Stenstrup, 21–27, 300.

Nieukerken EJ van (2004) Identity of two *Nepticula* species named by Walsingham (Lepidoptera: Nepticulidae, Tischeriidae). Tijdschrift voor Entomologie 147: 111–112. doi: 10.1163/22119434-900000145

Nieukerken EJ van (2006) Records of mining Lepidoptera in Belgium with nine species new to the country (Nepticulidae, Opostegidae, Tischeriidae, Lyonetiidae). Phegea 34: 125–144. http://webh01.ua.ac.be/vve/Phegea/2006/Phegea34–4_125–144.pdf

Nieukerken EJ van (2007a) *Acalyptris* Meyrick: revision of the *platani* and *staticis* groups in Europe and the Mediterranean (Lepidoptera: Nepticulidae). Zootaxa 1436: 1–48. http://www.mapress.com/zootaxa/2007f/zt01436p048.pdf

Nieukerken EJ van (2007b) Review of the subgenus *Trifurcula* (*Levarchama*), with two new species (Lepidoptera: Nepticulidae). Acta Zoologica Academiae Scientiarum Hungaricae 53: 101–140. http://www.repository.naturalis.nl/record/261195

Nieukerken EJ van (2008) Two new species in the *Ectoedemia* (*Fomoria*) *weaveri*-group from Asia (Lepidoptera: Nepticulidae). Zoologische Mededelingen (Leiden) 82: 113–130. http://www.repository.naturalis.nl/document/93755.

Nieukerken EJ van (2010) Order Lepidoptera, family Nepticulidae. In: Harten A van (Ed) Arthropod Fauna of the United Arab Emirates, volume 3. Dar Al Ummah, Abu Dhabi, 492–514. http://repository.naturalis.nl/document/201468

Nieukerken EJ van (2013) Fauna Europaea: Crinopterygidae, Heliozelidae, Nepticulidae, Opostegidae, Schreckensteiniidae, Urodidae. In: Karsholt O, van Nieukerken EJ (Eds) 2013. Lepidoptera. Fauna Europaea version 2.6.2. Fauna Europaea, Copenhagen, Amsterdam, Paris. http://www.faunaeur.org/

Nieukerken EJ van (Ed) (2016) Nepticulidae and Opostegidae of the world. Scratchpads, biodiversity online. http://nepticuloidea.info/

Nieukerken EJ van, Berg C van den (2003) A new *Stigmella* feeding on Urticaceae from Guam: first records of Nepticulidae (Lepidoptera) from Micronesia and Polynesia. Invertebrate Systematics 17: 27–37. doi: 10.1071/IS02019

Nieukerken EJ van, Berggren K (2011) *Ectoedemia rosae*, a new species from the French Alps and Norway (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 154: 181–191. doi: 10.1163/22119434-900000316

Nieukerken EJ van, Dop H (1987) Antennal sensory structures in Nepticulidae (Lepidoptera) and their phylogenetic implications. Zeitschrift für Zoologische Systematik und Evolutionsforschung 25: 104–126. doi: 10.1111/j.1439-0469.1987.tb00595.x

Nieukerken EJ van, Johansson R (1987) Identity of hitherto doubtful specific names in European Nepticulidae (Lepidoptera). Zoologische Mededelingen (Leiden) 61: 459–474. http://www.repository.naturalis.nl/document/150254

Nieukerken EJ van, Johansson R (1990) Tribus Trifurculini. In: Johansson R, Nielsen ES, Nieukerken EJ van, Gustafsson B (Eds) The Nepticulidae and Opostegidae (Lepidoptera) of NW Europe. Fauna Entomologica Scandinavica 23: 239–321.

Nieukerken EJ van, Johansson R (2003) The *Quercus* feeding *Stigmella* species of the West Palaearctic: new species, key and distribution (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 146: 307–370. doi: 10.1163/22119434–900000129

Nieukerken EJ van, Kuroko H (2005) Redescription of *Stigmella kurilensis* Puplesis (Lepidoptera, Nepticulidae), found on Hokkaido. Transactions of the Lepidopterological Society of Japan 56: 31–35. http://www.repository.naturalis.nl/record/227765

Nieukerken EJ van, Laštůvka A (2002) *Ectoedemia* (*Etainia*) *obtusa* Puplesis & Diškus, new for Europe: taxonomy, distribution and biology (Lepidoptera: Nepticulidae). Nota Lepidopterologica 25: 87–95. http://biostor.org/reference/143089

Nieukerken EJ van, Liu Y (2000) Nepticulidae (Lepidoptera) in China, 1. Introduction and *Stigmella* Schrank feeding on Fagaceae. Tijdschrift voor Entomologie 143: 145–181. doi: 10.1163/22119434–99900042

Nieukerken EJ van, Puplesis R (1991) Taxonomy and distribution of the *Trifurcula* (*Glaucolepis*) *raikhonae* group (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 134: 201–210. http://biodiversitylibrary.org/part/66479

Nieukerken EJ van, Hoare RJB, Groenenberg DSJ (2004a) Evolution of Nepticulidae & host plant use. Abstracts XXII International Congress of Entomology, 15–21 August 2004. Brisbane, Australia.

Nieukerken EJ van, Laštůvka A, Laštůvka Z (2004b) Annotated catalogue of the Nepticulidae and Opostegidae (Lepidoptera: Nepticuloidea) of the Iberian Peninsula. SHILAP Revista de Lepidopterologia 32: 211–260. http://www.redalyc.org/pdf/455/45512710.pdf

Nieukerken EJ van, Zolotuhin VV, Mistchenko A (2004c) Nepticulidae from the Volga and Ural region. Nota Lepidopterologica 27: 125–157. http://www.biodiversitylibrary.org/part/147307

Nieukerken EJ van, Laštůvka A, Laštůvka Z (2006) Annotated catalogue of the Nepticulidae and Opostegidae (Lepidoptera: Nepticuloidea) of France and Corsica. Zootaxa 1216: 1–114. http://www.mapress.com/zootaxa/2006/zt01216p114.pdf

Nieukerken EJ van, Laštůvka A, Laštůvka Z (2010) Western Palaearctic *Ectoedemia* (*Zimmermannia*) Hering and *Ectoedemia* Busck s. str. (Lepidoptera: Nepticulidae): five new species and new data on distribution, hostplants and recognition. ZooKeys 32: 1–82. doi: 10.3897/zookeys.32.282

Nieukerken EJ van, Berg C van den, Hoare RJB (2011) A new species of the endemic Australian genus *Roscidotoga* Hoare from rainforests in southern Queensland (Lepidoptera: Nepticulidae). Tijdschrift voor Entomologie 154: 193–201. doi: 10.1163/22119434-900000318

Nieukerken EJ van, Mutanen M, Doorenweerd C (2012a) DNA barcoding resolves species complexes in *Stigmella salicis* and *S. aurella* species groups and shows additional cryptic speciation in *S. salicis* (Lepidoptera: Nepticulidae). Entomologisk Tidskrift 132: 235–255. http://www.repository.naturalis.nl/document/363525

Nieukerken EJ van, Doorenweerd C, Stokvis FR, Groenenberg DSJ (2012b) DNA barcoding of the leaf-mining moth subgenus *Ectoedemia* s. str. (Lepidoptera: Nepticulidae) with COI and EF1- α: two are better than one in recognising cryptic species. Contributions to Zoology 81: 1–24. http://www.contributionstozoology.nl/vol81/nr01/a01

Nieukerken EJ van, Doorenweerd C, Nishida K, Snyers C (2016) New taxa, including two new genera show uniqueness of Neotropic Nepticulidae (Lepidoptera). ZooKeys 628: 1–63. doi: 10.3897/zookeys.628.9805

Nolcken JHWB von (1871) Lepidopterologische Fauna von Estland, Livland und Kurland. Zweite Abtheilung, Microlepidoptera, 2. Heft. Arbeiten des Naturforscher-Vereins zu Riga 4: i-viii, 1–849. http://dspace.ut.ee/handle/10062/46698

Nolcken JHWB von (1882) Lepidopterologische Notizen. Stettiner Entomologische Zeitung 43: 173–201. http://biodiversitylibrary.org/page/34380163

Nowicki MS (1860) Enumeratio Lepidopterorum Haliciae Orientalis. sumptibus auctoris, Leopoli: i-xxx, i-xvi, 1–262 pp. doi: 10.5962/bhl.title.66986

Nuss M, Landry B, Mally R, Vegliante F, Tränkner A, Bauer F, Hayden J, Segerer A, Schouten R, Li H, Trofimova T, Solis MA, De Prins J, Speidel W (2003–2015) GlobiZ: Global Information System on Pyraloidea. Senckenberg Collection of Natural History, Museum of Zoology, Dresden (Germany). www.pyraloidea.org

Oberthür C (1915) Portraits de Lépidoptéristes, deuxième série. In: Oberthür C (Ed.) Études de Lépidoptérologie comparée. Oberthür, Rennes, 19 pls. http://biodiversitylibrary.org/page/10631606

Packard AS (1870) New or little known injurious insects. Annual Report of the Massachusetts Board of Agriculture 17: 235–238.

Parenti U (1963) *Stigmella saturejae*, nuova specie di microlepidottero, e la sua mina. Bollettino di Zoologia Agraria e Bachicoltura, Serie 2 5: 101–105.

Petersen W (1930a) Nachträge zur Lepidopterenfauna von Estland, mit besonderer Berücksichtigung der Gattungen *Lithocolletis* und *Nepticula*. Korrespondenzblatt des Naturforscher-Vereins zu Riga 60: 89–101.

Petersen W (1930b) Die Blattminierer-Gattungen *Lithocolletis* und *Nepticula* (Lep.). Teil II: *Nepticula* Z. Stettiner Entomologische Zeitung 91: 1–82, 3 pls.

Petry A (1904) *Nepticula thuringiaca* n. sp. Stettiner Entomologische Zeitung 65: 179–181. http://biodiversitylibrary.org/page/9010232

Peyerimhoff H de (1871) [no title.]. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 3: 409–415. http://biodiversitylibrary.org/page/49053833

Philpott A (1919) Descriptions of new species of Lepidoptera. Transactions and Proceedings of the New Zealand Institute 51: 224–225. http://rsnz.natlib.govt.nz/volume/rsnz_51/rsnz_51_00_003200.pdf

Philpott A (1927) Notes and descriptions of New Zealand Lepidoptera. Transactions and Proceedings of the New Zealand Institute 58: 80–90. http://rsnz.natlib.govt.nz/volume/rsnz_58/rsnz_58_00_000930.pdf

Pierce FN, Metcalfe JW (1935) The genitalia of the Tineid families of the Lepidoptera of the British islands. F.N. Pierce, Oundle, Northants.

Porritt GT (1883) List of Yorkshire Lepidoptera. Transactions of the Yorkshire Naturalist's Union, Series D Entomology 2: 1–198. doi: 10.5962/bhl.title.7900

Preissecker F (1942) Zwei neue *Nepticula*-Arten aus dem Gebiete des heutigen Reichsgaues Wien. Zeitschrift des Wiener Entomologen-Vereines 27: 208–211. http://www.zobodat.at/pdf/ZOEV_27_0208-0211.pdf

Pröse H (1984) Neue Ergebnisse zur Faunistik der Microlepidoptera in Bayern. 2. Beitrag. Nachrichtenblatt der Bayerischen Entomologen 33: 106–115. http://www.zobodat.at/pdf/NachBlBayEnt_033_0106-0115.pdf

Pullen KR, Jennings D, Oberprieler RG (2014) Annotated catalogue of Australian weevils (Coleoptera: Curculionoidea). Zootaxa 3896: 1–481. doi: 10.11646/zootaxa.3896.1.1

Puplesis RK (1984a) Novye vidy Molej-malyutok (Lepidoptera, Nepticulidae) iz juzhnogo Primor'ya. (New species of nepticulids (Lepidoptera, Nepticulidae) from southern Primorye). Entomologicheskoe Obozrenie 63: 111–125. [In Russian]

Puplesis RK (1984b) K sisteme molej-malyutok (Lepidoptera, Nepticulidae) palearkticeskoj fauny. (To the classification of nepticulids (Lepidoptera, Nepticulidae) of the Palaearctic fauna). Entomologicheskoe Obozrenie 63: 582–597. [In Russian]

Puplesis RK (1984c) Obzor vidov roda *Microcalyptris* (Lepidoptera, Nepticulidae) iz pustyn' Mongolii i SSSR (A review of nepticulid moths of the genus *Microcalyptris* (Lepidoptera, Nepticulidae) from the deserts of Mongolia and the USSR). Nasekomye Mongolii 9: 484–507. [In Russian]

Puplesis RK (1984d) Pervye nakhodki rodov *Stigmella* i *Ectoedemia* (Lepidoptera, Nepticulidae) v Mongolii. (First records of the genera *Stigmella* and *Ectoedemia* (Lepidoptera, Nepticulidae) in Mongolia). Nasekomye Mongolii 9: 508–510. [In Russian]

Puplesis RK (1984e) New species of (Lepidoptera) Nepticulidae. Entomological Review Washington 63: 96–111. [English translation of Puplesis 1984a]

Puplesis RK (1985a) Moli-malyutki (Lepidoptera, Nepticulidae) Pribaltiki i Dal'nego vostoka. Avtoreferat, Leningrad: Zoological Institute, Academy of Sciences USSR. [In Russian]

Puplesis RK (1985b) Contribution to the classification of the palaearctic Nepticulidae (Lepidoptera). Entomological Review Washington 63: 149–164. [English translation of Puplesis 1984b]

Puplesis RK (1985c) Novye vidy molej-malyutok (Lepidoptera, Nepticulidae) s yuga dal'nego vostoka i Tadzhikistana (New species of the nepticulid moths from Southern far east and Tadzhikistan). Trudy Zoologicheskogo Instituta Akademiya Nauk SSSR 134: 59–72. [In Russian]

Puplesis RK (1987) Kratkij zoogeograficheskij obzor molej-malyutok (Lepidoptera, Nepticulidae) yuga Dal'nego Vostoka s opisaniem novykh vidov [A short zoogeographical survey of Nepticulidae from the south of the Far East with descriptions of new species.]. In: Ler PA, Kirpichnikova VA, Kononenko VS (Eds) Cheshuekrylye Dal'nego Vostoka SSSR [Lepidoptera of the Far East of the USSR]. Far East Scientific Center, Vladivostok, 4–13. [In Russian]

Puplesis R (1988) New species of plant mining Lepidoptera (Nepticulidae, Tischeriidae) from central Asia. Stapfia 16: 273–290. http://www.zobodat.at/pdf/STAPFIA_0016_0273-0290.pdf

Puplesis R (1990) The genus *Acalyptris* Meyrick (Lepidoptera, Nepticulidae) in the USSR: distribution and taxonomy. Nota Lepidopterologica 13: 62–88. http://www.biodiversitylibrary.org/part/118095

Puplesis R (1991) The *Stigmella paradoxa* species-group (Lepidoptera: Nepticulidae) in the USSR. Entomologica Scandinavica 22: 123–127. doi: 10.1163/187631291X00345

Puplesis R (1994) The Nepticulidae of eastern Europe and Asia. Western, central and eastern parts. Backhuys Publishers, Leiden, 290 pp.

Puplesis RK, Arutyunova NV (1991) Dva novykh vida molej-malyutok (Lepidoptera, Nepticulidae), miniruyushchikh list'ya Yabloni, iz Tadzhikistana. (Two new species of nepticulid moths (Lepidoptera, Nepticulidae), mining apple-tree leaves from Tajikistan). Entomologicheskoe Obozrenie 70: 571–573.

Puplesis R, Diškus A (1995) *Acalyptris argyraspis* sp. n., a remarkable species from Tadzhikistan (Lepidoptera: Nepticulidae). Phegea 23: 51–54. http://www.phegea.org/Phegea/1995/Phegea23-1_51-54.pdf

Puplesis R, Diškus A (1996a) First record of the genus *Etainia* Beirne from Central Asia with descriptions of two new species and some provisional notes on the world fauna (Lepidoptera: Nepticulidae). Phegea 24: 41–48. http://www.phegea.org/Phegea/1996/Phegea24-1_41-48.pdf

Puplesis R, Diškus A (1996b) A review of the *Stigmella sorbi* species-group with descriptions of two new species from Turkmenistan and Tadzhikistan (Lepidoptera: Nepticulidae). Phegea 24: 171–182. http://www.phegea.org/Phegea/1996/Phegea24-4_171-182.pdf

Puplesis R, Diškus A (1996c) Five new mining Lepidoptera (Nepticulidae, Bucculatricidae) from central Asia. Tijdschrift voor Entomologie 139: 181–190. http://www.biodiversitylibrary.org/part/65974

Puplesis R, Diškus A (2003a) Strateginiu regionu taksonomines revizijos ir nauju rusiu aprasai. Strategic regional revisions with description of new species. In: Puplesis R, Diškus A (Eds) Nepticuloidea ir Tischerioidea (Lepidoptera) pasaulio ir lietuvos faunoje. The Nepticuloidea & Tischerioidea (Lepidoptera) - a global review, with strategic regional revisions. Lututе publishers, Kaunas, 176–289.

Puplesis R, Diškus A (2003b) Nepticuloidea ir Tischerioidea (Lepidoptera) pasaulio ir lietuvos faunoje. The Nepticuloidea & Tischerioidea (Lepidoptera) - a global review, with strategic regional revisions. Lututе publishers, Kaunas, 512 pp.

Puplesis RK, Ivinskis PP (1985) Obzor fauny moleï-malyutok roda *Obrussa* Braun SSSR i opisanie 4 novykh vidov *Obrussa capesella* sp. n., *O. tigrinella* sp. n., *O. peterseni* sp. n., *O. sabina* sp. n. [Review of nepticulid moths fauna of the Obrussa Braun genus in the USSR and a description of 4 new species - *Obrussa capesella* sp. n., *O. tigrinella* sp. n., *O. peterseni* sp. n., *O. sabina* sp. n]. Trudy Akademii Nauk Litovskoi SSR Seriya C 92: 36–46. [In Russian]

Puplesis R, Robinson GS (1999) Revision of the Oriental Opostegidae (Lepidoptera) with general comments on phylogeny within the family. Bulletin of the Natural History Museum London Entomology 68: 1–92. http://biodiversitylibrary.org/part/76471

Puplesis R, Robinson GS (2000) A review of the Central and South American Nepticulidae (Lepidoptera) with special reference to Belize. Bulletin of the Natural History Museum London Entomology 69: 3–114. http://www.biodiversitylibrary.org/part/76459

Puplesis R, Seksjaeva S, Sruoga V (1991) Leaf-mining Lepidoptera (Nepticulidae, Bucculatricidae, Gracillariidae) from *Ulmus* in northern Caspiya (Kaspia). Tijdschrift voor Entomologie 134: 69–73. http://biodiversitylibrary.org/page/28239905

Puplesis R, Seksjaeva S, Noreika R, Puplesiene J (1992) Some leaf-mining Lepidoptera from the Aksu Dzhabagly Reserve (western Tian Shan) with the descriptions of four new species (Lepidoptera: Nepticulidae, Bucculatricidae). Nota Lepidopterologica 15: 47–64. http://www.biodiversitylibrary.org/part/81848

Puplesis R, Diškus A, Noreika R, Saparmamedova N (1996) Revised check-list of mining Lepidoptera (Nepticuloidea, Tischerioidea and Gracillarioidea) from central Asia. Tijdschrift voor Entomologie 139: 191–200. http://www.biodiversitylibrary.org/part/65975

Puplesis R, Diškus A, Nieukerken EJ van (1997) *Stigmella divina* sp. n., a remarkable species from Turkmenistan and Turkey (Lepidoptera, Nepticulidae). Tijdschrift voor Entomologie 140: 55–58. http://www.biodiversitylibrary.org/part/66284

Puplesis R, Diškus A, Robinson GS (2002a) New Neotropical Nepticulidae (Lepidoptera) from the western Amazonian rainforest and the Andes of Ecuador. Bulletin of the Natural History Museum London Entomology 71: 19–58. doi: 10.1017/s0968045402000020

Puplesis R, Diškus A, Robinson GS, Onore G (2002b) A review and check list of the Neotropical Nepticulidae (Lepidoptera). Bulletin of the Natural History Museum London Entomology 71: 59–76. http://www.biodiversitylibrary.org/part/76406

Ratnasingham S, Hebert PDN (2007) BOLD: The Barcode of Life Data System (www.barcodinglife.org). Molecular Ecology Notes 7: 355–364. doi: 10.1111/j.1471–8286.2006.01678.x

Rebel H (1901) Catalog der Lepidopteren des Palaearctischen Faunengebietes. II. Theil: Famil. Pyralidae - Micropterygidae. Friedländer & Sohn, Berlin: xxx, 368 pp.

Rebel H (1909) [no title]. In: Versammlung am 4. Dezember 1908. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 58: (269). http://biodiversitylibrary.org/page/13250061

Rebel H (1910) Sechster Beitrag zur Lepidopterenfauna der Kanaren. Annalen des Kaiserlich-Königlichen Naturhistorischen Hofmuseums, Wien 24: 327–374. http://www.zobodat.at/pdf/ANNA_24_0327-0374.pdf

Rebel H (1926) Beitrag zur Microlepidopterenfauna Dalmatiens. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 74: (106)-(110). http://www.zobodat.at/pdf/VZBG_74-75_0001-0204.pdf

Rebel H (1933) Zwei neue Microlepidopteren aus Dalmatien. Zeitschrift des Österreichischen Entomologen-Vereines 18: 81–82. http://www.zobodat.at/pdf/ZOEV_18_0077-0080.pdf

Regier JC, Mitter C, Kristensen NP, Davis DR, Nieukerken EJ van, Rota J, Simonsen TJ, Mitter KT, Kawahara AY, Yen S-H, Cummings MP, Zwick A (2015) A molecular phylogeny for the oldest (nonditrysian) lineages of extant Lepidoptera, with implications for classification, comparative morphology and life-history evolution. Systematic Entomology 40: 671–704. doi: 10.1111/syen.12129

Remeikis A, Stonis JR (2015) First discovery of *Quercus* -feeding Nepticulidae (Lepidoptera) in South America, with description of new species and designation of the *S. nigriverticella* complex in the *S. saginella* group. Zootaxa 4057: 409–429. doi: 10.11646/zootaxa.4057.3.7

Remeikis A, Stonis JR, Diškus A, Davis DR (2009) Contribution to the Opostegidae fauna of Central America, with an updated checklist and description of new species from Costa Rica and Mexico (Insecta: Lepidoptera). Acta Zoologica Lituanica 19: 268–277. doi: 10.2478/v10043-009-0037-0

Retzius AI (1783) Genera et Species Insectorum (e generosissimi auctori scriptis). Siegfried Lebrecht Crusium, Lipsiae, i-vi, 1–220 pp. doi: 10.5962/bhl.title.12419

Reutti C (1853) Übersicht der Lepidopteren-Fauna des Grossherzogtum´s Baden. Beiträge zur Rheinischen Naturgeschichte 3: 1–216. http://biodiversitylibrary.org/page/33803749

Richardson NM (1890) Description of a *Nepticula* (*auromarginella*) new to science from near Weymouth. Entomologist's Monthly Magazine 26: 30–31. http://biodiversitylibrary.org/page/33378802

Robinson GS, Nielsen ES (1983) The Microlepidoptera described by Linnaeus and Clerck. Systematic Entomology 8: 191–242. doi: 10.1111/j.1365–3113.1983.tb00479.x

Rocienė A, Stonis JR (2013) Nepticulidae (Lepidoptera) of East Asia (2). Study of a collection sample deposited at the Russian Academy of Sciences, with descriptions of new species and a checklist. Zootaxa 3652: 75–116. doi: 10.11646/zootaxa.3652.1.3

Rocienė A, Stonis JR (2014) Revised fauna of the Nepticulidae (Lepidoptera) of continental East Asia: lots of effort to elucidate the little-known diversity of pygmy moths. In: Stonis JR (Ed.) Selected abstracts and papers of the First Baltic International Conference on Field Entomology and Faunistics. Edukologija Publishers, Vilnius, 51–62 pp.

Roskov Y, Abucay L, Orrell T, Nicolson D, Kunze T, Flann C, Bailly N, Kirk P, Bourgoin T, DeWalt RE, Decock W, De Wever A (2015) Species 2000 & ITIS Catalogue of Life. Species 2000: Naturalis. http://www.catalogueoflife.org/col

Rössler A (1867) Verzeichnis der Schmetterlinge des Herzogthums Nassau, mit besonderer Berücksichtigung der biologischen Verhältnisse und der Entwicklungsgeschichte. Jahrbuch des Nassauischen Vereins für Naturkunde 19+20: 99–442 (1–342).

Rössler A (1881) Die Schuppenflügler (Lepidopteren) des Kgl. Regierungsbezirks Wiesbaden und ihre Entwicklungsgeschichte. Jahrbuch des Nassauischen Vereins für Naturkunde 33 + 34: 1–392. http://www.entomologische-literatur.de/uploads/media/R_Roessler__Jahrb._Nass.Ver._1881__Schuppenfluegler_S.1-200.pdf

Rothschild NC (1912) Adatok Magyarország lepkefaunájához. [Data to the Lepidoptera fauna of Hungary]. Rovartani Lapok 19: 21–29. http://biodiversitylibrary.org/page/32992423

Rungs CEE (1979) Catalogue raisonné des Lepidoptères du Maroc. Inventaire faunistique et observations écologiques, 1. Travaux de l'Institut Scientifique Chérifien, Série Zoologique 39: 1–222, A221–A225.

Sand M (1879) Catalogue raisonné des Lépidoptères du Berry & de l'Auvergne (Cher, Indre, Creuse, Puy-de-Dôme) (France centrale). E. Deyrolle, Paris, vi+206 pp.

Sattler K (1973) A catalogue of the family-group and genus-group names of the Gelechiidae, Holcopogonidae, Lecithoceridae and Symmocidae (Lepidoptera). Bulletin of the British Museum (Natural History) Entomology 28: 153–282. http://www.biodiversitylibrary.org/page/2252332

Sauber A (1904) Die Kleinschmetterlinge Hamburgs und der Umgegend. Verhandlungen des Vereins für Naturwissenschaftliche Unterhaltung zu Hamburg 12: 1–60. http://biodiversitylibrary.org/page/27366980

Schintlmeister A (2013) Notodontidae & Oenosandridae (Lepidoptera). Brill, Leiden, World Catalogue of Insects 11: 608 pp.

Schleich CL (1867) Einige microlepidopterologische Beobachtungen über eine neue *Nepticula*, die Raupe von *Gelech. micella* und über *Gracil. imperialella*. Stettiner Entomologische Zeitung 28: 449–455. http://biodiversitylibrary.org/page/33831516.

Schoorl JW, Wilkinson C (1986) The *Stigmella betulicola* species-group in Europe (Lepidoptera: Nepticulidae). Journal of Natural History 20: 225–244. doi: 10.1080/00222938600770181

Schoorl JW, Nieukerken EJ van, Wilkinson C (1985) The *Stigmella oxyacanthella* species-group in Europe (Nepticulidae: Lepidoptera). Systematic Entomology 10: 65–103. doi: 10.1111/j.1365–3113.1985.tb00565.x

Schrank F von Paula (1802) Fauna Boica. Durchgedachte Geschichte der in Baiern einheimischen und zahmen Thiere. Zweyter Band, zweyte Abtheilung. J. W. Krüll, Ingolstadt, 2(2): 412 pp. doi: 10.5962/bhl.title.44923

Schütze KT (1905) Mitteilungen über einige Kleinschmetterlinge. Deutsche Entomologische Zeitschrift Iris 17: 192–208. http://biodiversitylibrary.org/page/12683191.

Scoble MJ (1978a) Nepticulidae (Lepidoptera) of southern Africa: the genus *Ectoedemia* Busck. Journal of the Entomological Society of Southern Africa 41: 81–86.

Scoble MJ (1978b) Nepticulidae of southern Africa: a taxonomic revision of the genus *Stigmella* Schrank (Lepidoptera: Monotrysia). Annals of the Transvaal Museum 31: 1–129. http://hdl.handle.net/10499/AJ1837

Scoble MJ (1979) A new species of *Ectoedemia* Busck from Botswana with observations on its imaginal skeletal anatomy (Lepidoptera: Nepticulidae). Annals of the Transvaal Museum 32: 35–54. http://hdl.handle.net/10499/AJ1828

Scoble MJ (1980a) *Trifurcula* Zeller: a critical analysis of the genus, cladistic relationships and descriptions of two new species from South Africa (Lepidoptera: Nepticulidae). Journal of the Entomological Society of Southern Africa 43: 139–150.

Scoble MJ (1980b) The genus *Niepeltia* Strand: taxonomy, and comments on structure and relationships (Lepidoptera: Nepticulidae). Annals of the Transvaal Museum 32: 197–229. http://hdl.handle.net/10499/AJ1823

Scoble MJ (1982) A pectinifer in the Nepticulidae (Lepidoptera) and its phylogenetic implications. Annals of the Transvaal Museum 33: 123–129. http://hdl.handle.net/10499/AJ10150

Scoble MJ (1983) A revised cladistic classification of the Nepticulidae (Lepidoptera) with descriptions of new taxa mainly from South Africa. Transvaal Museum Monograph 2: 1–105. http://hdl.handle.net/10499/AJ10150

Segerer AH (1997) Verifikation älterer und fraglicher Regensburger Lepidopterenmeldungen (Insecta: Lepidoptera). Beiträge zur bayerischen Entomofaunistik 2: 177–265.

Shield R (1853) List of Micro-Lepidoptera taken in the vicinity of Dublin during the year 1853. Zoologist 11: 4152–4153. http://biodiversitylibrary.org/page/39806334

Shinozaki Y, Yamamoto A, Oishi M, Sato H (2012) Early leaf abscission has little effect on larval mortality of *Ectoedemia cerviparadisicola* (Lepidoptera, Nepticulidae) associated with *Quercus gilva*. Annals of the Entomological Society of America 105: 572–581. doi: 10.1603/an11185

Šimkevičiūtė A, Stonis JR, Diškus A (2009) Taxonomic checklist of Nepticulidae of Mexico, with the description of three new species from the Pacific Coast (Insecta, Lepidoptera). Acta Zoologica Lituanica 19: 268–277. doi: 10.2478/v10043-009-0037-0

Sinev SY (1990) Novye dannye po sistematike Molej-Opostegid (Lepidoptera, Opostegidae) Dalnego Vostoka SSSR. [New data on the systematics of opostegid moths (Lepidoptera,

Opostegidae) of Far East SSSR]. Novosti faunistiki i sistematiki: sbornik nauchn'ykh trudov [New data on faunistics and systematics : collected scientific papers]. Naukova Dimka, Kiev, 100–106. [In Russian]

Sircom J (1849) Descriptions of three new British Tineidae. Zoologist 7: xlii. http://biodiversitylibrary.org/page/39815380

Skala H (1933a) Neue Neptikel. Zeitschrift des Österreichischen Entomologen-Vereines 18: 30–32. http://www.zobodat.at/pdf/ZOEV_18_0030-0032.pdf

Skala H (1933b) Die Nepticulidae Oberösterreichs. Entomologischer Anzeiger 13: 149–152. http://www.zobodat.at/pdf/EntAnz_13_0149-0152.pdf

Skala H (1934a) *Neptic. ulmi* spec. nov. und andere wenig bekannte Neptikel. Zeitschrift des Österreichischen Entomologen-Vereines 19: 51. http://www.zobodat.at/pdf/ZOEV_19_0051.pdf

Skala H (1934b) Die Nepticulidae Oberösterreichs. Entomologischer Anzeiger 14: 5–8. http://www.zobodat.at/pdf/EntAnz_14_0005-0008.pdf

Skala H (1934c) Die Nepticulidae Oberösterreichs. Entomologischer Anzeiger 14: 29–32. http://www.zobodat.at/pdf/EntAnz_14_0029-0032.pdf

Skala H (1935) Zur Minenfauna Oberösterreichs. Zeitschrift des Österreichischen Entomologen-Vereines 20: 77–80. http://www.zobodat.at/pdf/ZOEV_20_0077-0080.pdf

Skala H (1936) Minen aus Mittel- und Südeuropa. Zeitschrift des Österreichischen Entomologen-Vereins 21: 78–79. http://www.zobodat.at/pdf/ZOEV_21_0078-0079.pdf

Skala H (1938a) Einiges über Falter-Minen aus dem Mediterrangebiet. Zeitschrift des Österreichischen Entomologen-Vereins 23: 8–10. http://www.zobodat.at/pdf/ZOEV_23_0008-0010.pdf

Skala H (1938b) Einiges über Falter-Minen aus dem Mediterrangebiet. Zeitschrift des Österreichischen Entomologen-Vereins 23: 43–46. http://www.zobodat.at/pdf/ZOEV_23_0043-0046.pdf

Skala H (1939a) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 27–30. http://www.zobodat.at/pdf/ZOEV_24_0027-0030.pdf

Skala H (1939b) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 43–45, pls 5–6. http://www.zobodat.at/pdf/ZOEV_24_0043-0045.pdf

Skala H (1939c) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 62–64, pls 7–9. http://www.zobodat.at/pdf/ZOEV_24_0062-0064.pdf

Skala H (1939d) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 91–95. http://www.zobodat.at/pdf/ZOEV_24_0091-0095.pdf

Skala H (1939e) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 111–112. http://www.zobodat.at/pdf/ZOEV_24_0111-0112.pdf

Skala H (1939f) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 125–128. http://www.zobodat.at/pdf/ZOEV_24_0125-0128.pdf

Skala H (1939g) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 138–144. http://www.zobodat.at/pdf/ZOEV_24_0138-0144.pdf

Skala H (1939h) Miner in deutschen Landen. Zeitschrift des Österreichischen Entomologen-Vereins 24: 149–152, pl. 110. http://www.zobodat.at/pdf/ZOEV_24_0149-0152.pdf

Skala H (1940) In der Welser Heide beobachtete Falter. Zeitschrift des Wiener Entomologen-Vereines 25: 143–144. http://www.zobodat.at/pdf/ZOEV_25_0143-0144.pdf

Skala H (1941a) Neues über Miner. Zeitschrift des Wiener Entomologen-Vereines 26: 55–57. http://www.zobodat.at/pdf/ZOEV_26_0055-0057.pdf

Skala H (1941b) Neues über Miner. Zeitschrift des Wiener Entomologen-Vereines 26: 77–80. http://www.zobodat.at/pdf/ZOEV_26_0077-0080.pdf

Skala H (1942) Falter von Haid und anderes. Zeitschrift des Wiener Entomologen-Vereines 27: 5–7. http://www.zobodat.at/pdf/ZOEV_27_0005-0007.pdf

Skala H (1943) *Nepticula groschkei* spec. nov. Zeitschrift der Wiener Entomologischen Gesellschaft 28: 86. http://www.zobodat.at/pdf/ZOEV_28_0086.pdf

Skala H (1948) Zur Familie der Miner (Nepticulidae). Zeitschrift der Wiener Entomologischen Gesellschaft 32: 121–122. http://www.zobodat.at/pdf/ZOEV_32_0081-0090.pdf

Skala H (1949) Wenig und nicht bekannte Minen der Gattung *Nepticula*. Zeitschrift der Wiener Entomologischen Gesellschaft 34: 129–131, pl. 112, 113. http://www.zobodat.at/pdf/ZOEV_34_0129-0131.pdf

Snellen PCT (1873) Eenige aanteekeningen op Catalog der Lepidopteren des Europäischen Faunengebietes bearbeitet von Dr O Staudinger en Dr M Wocke. Tijdschrift voor Entomologie 16: 36–60. http://www.biodiversitylibrary.org/part/29466

Snellen PCT (1875) *Nepticula zelleriella* nov. sp. Tijdschrift voor Entomologie 18: 113–117, pl. 117. http://biostor.org/reference/13327

Snellen PCT (1882) De vlinders van Nederland. Microlepidoptera, systematisch beschreven, 2 volumes. E.J. Brill, Leiden, 1–2: 1–536, 537–1196, 14 pls. pp.

Sobczyk T (2011) Psychidae (Lepidoptera). Brill, Leiden, World Catalogue of Insects 10: 467 pp.

Soffner J (1942) Eine neue *Nepticula*-Art. Mitteilungen der Deutschen Entomologischen Gesellschaft 11: 56–61.

Sohn J-C, Labandeira CC, Davis DR, Mitter C (2012) An annotated catalog of fossil and subfossil Lepidoptera (Insecta: Holometabola) of the world. Zootaxa 3286: 1–132. http://www.mapress.com/zootaxa/2012/f/zt03286p132.pdf

Sokolov SJ, Svjazeva OA, Kubli VA (1980) Arealy derev'ev i kustarnikov SSSR I. Tissovye-Kirkazonovye. Areographia arborum fruticumque URSS 1. Taxaceae-Aristolochiaceae. Nauka (Leningrad) 1: 1–164, 91 maps. [In Russian]

Sommerer M (2002) To agree or not to agree – the question of gender agreement in the International Code of Zoological Nomenclature. Nota Lepidopterologica 25: 191–204. http://biodiversitylibrary.org/page/41371966

Sorhagen L (1885) Die Microlepidopteren der Hamburger Torfmoore. Entomologische Nachrichten, Berlin 11: 257–264. http://biostor.org/reference/111630

Sorhagen L (1922) Beiträge zur Biologie Europäischer *Nepticula*-Arten. Archiv für Naturgeschichte 88A: 9–60. http://biodiversitylibrary.org/page/13259635

Spuler A, Meess A (1910) Superfam. Tineides Aculeatae. In: Spuler A (Ed.) Die Schmetterlinge Europas II Band. Schweizerbart, Stuttgart, 464–482, pl. 491. doi: 10.5962/bhl.title.9477

Stainton HT (1848) A monograph on the British *Argyromiges*. Zoologist 6: 2152-2164. http://biodiversitylibrary.org/page/39803972

Stainton HT (1849) An attempt at a systematic catalogue of the British Tineidae & Pterophoridae. John van Voorst, London: iv+ 32 pp. http://sammlungen.ub.uni-frankfurt.de/varia/id/5427655

Stainton HT (1851) A supplementary catalogue of the British Tineidae & Pterophoridae. John van Voorst, London: iv+28 pp. https://books.google.nl/books?id=tfRhAAAAcAAJ&hl=nl&pg=PP1#v=onepage&q&f=false

Stainton HT (1853) An introduction to the study of Nepticulidae. Zoologist 11: 3952–3960. http://biodiversitylibrary.org/page/39806134

Stainton HT (1854) Insecta Brittanica. Lepidoptera: Tineina. Lovell Reeve, London, 313 pp. doi: 10.5962/bhl.title.9558

Stainton HT (1855) Lepidoptera. The new British species in 1854. Entomologist's Annual 1855: 62–72. http://biodiversitylibrary.org/page/10427580

Stainton HT (1856) Lepidoptera. New British species in 1855. Entomologist's Annual 1856: 26–45. http://biodiversitylibrary.org/page/10427732

Stainton HT (1857a) Lepidoptera. New British species in 1856. Entomologist's Annual 1857: 97–112. http://biodiversitylibrary.org/page/10427982

Stainton HT (1857b) *Nepticula Myrtillella*. Entomologist's Weekly Intelligencer 2: 44. http://biodiversitylibrary.org/page/50829919

Stainton HT (1857c) *Nepticula Poterii*. Entomologist's Weekly Intelligencer 2: 116. http://biodiversitylibrary.org/page/50829991

Stainton HT (1858) Lepidoptera. New British species in 1857. Entomologist's Annual 1858: 85–98. http://biodiversitylibrary.org/page/10429446

Stainton HT (1859a) Manual of British butterflies and moths, 2. Cooke & Son, London: xi, 480 pp. https://books.google.nl/books?id=xv0GAQAAIAAJ&hl=nl&pg=PR1#v=onepage&q&f=false

Stainton HT (1859b) A new *Nepticula*. Entomologist's Weekly Intelligencer 5: 123. http://biodiversitylibrary.org/page/50830218

Stainton HT (1861) Lepidoptera. New British species in 1860. Entomologist's Annual 1861: 82–92. http://biodiversitylibrary.org/page/10428910

Stainton HT (Ed) (1869a) The Tineina of Southern Europe. John van Voorst, London, 370 pp. doi: 10.5962/bhl.title.26568

Stainton HT (1869b) Tineina observed at Cannes and Mentone in February and March 1867. In: Stainton HT (Ed) The Tineina of Southern Europe. John van Voorst, London, 216–232. http://biodiversitylibrary.org/page/21067701

Stainton HT (1883) Philipp Christoph Zeller. Entomologist's Monthly Magazine 20: 1–89, frontispiece. http://biodiversitylibrary.org/page/9267666

Stainton HT (1884) Note on a new *Nepticula* bred from rose in Lancashire by Mr Hodgkinson. Entomologist's Monthly Magazine 21: 103. http://biodiversitylibrary.org/page/9276447

Stainton HT (1886) *Nepticula argyropeza* and *apicella*. Entomologist's Monthly Magazine 22: 237–238. http://biodiversitylibrary.org/page/9276885

Stainton HT (1887) A new species of *Nepticula* bred from birch, from Herefordshire (*N. Woolhopiella*). Entomologist's Monthly Magazine 24: 62. http://biodiversitylibrary.org/page/9267104

Stainton HT (1888a) *Nepticula serella*, n. sp. Entomologist's Monthly Magazine 24: 260. http://biodiversitylibrary.org/page/9267302

Stainton HT (1888b) Description of a new *Nepticula* from Beech. Entomologist's Monthly Magazine 25: 12–13. http://biodiversitylibrary.org/page/30320955

Stainton HT, Zeller PC, Douglas JW (1855) *Nepticula*, part 1; *Cemiostoma*, part 1. John van Voorst, London, The natural history of the Tineina 1: i-xiii, 1–338, 7+1 pls. https://books.google.nl/books?id=0jsDAAAAQAAJ&hl=nl&pg=PR1#v=onepage&q&f=false

Stainton HT, Zeller PC, Douglas JW, Frey H (1862) *Bucculatrix*; part 1; *Nepticula*, part 2. John van Voorst, London, The natural history of the Tineina 7: i–vii,1– 251, 4 + 4 pls. http://biodiversitylibrary.org/page/20379482

Staudinger O (1871) Beschreibung neuer Lepidopteren des europäischen Faunanegebiets. Berliner Entomologische Zeitschrift 14: 273–330. http://biodiversitylibrary.org/page/8340899

Stephens JF (1829) A systematic catalogue of British Insects, being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities: containing also the references to every English writer on entomology, and to the principal foreign authors : with all the published British genera to the present time. Baldwin and Cradock, London: xxxiv, 388 pp. doi: 10.5962/bhl.title.8987

Stephens JF (1834) Illustrations of British Entomology, or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food and economy as far as practicable. Haustellata 4. Baldwin and Cradock, London: 433 pp., pls 33–41. http://biodiversitylibrary.org/page/8185399

Steudel W, Hofmann E (1882) Verzeichniss württembergischer Kleinschmetterlinge. Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 38: 143–262. http://biodiversitylibrary.org/page/11313547

Stonis JR, Remeikis A (2015) The first records on the genus *Acalyptris* from the Caribbean (Lepidoptera: Nepticulidae). Zootaxa 4057: 79–90. doi: 10.11646/zootaxa.4057.1.4

Stonis JR, Remeikis A (2016) Southern Andean *Stigmella sinuosa* complex (Lepidoptera, Nepticulidae): unraveling problematic taxonomy with a pictorial key of adults? Zootaxa 4136: 309–322. doi: 10.11646/zootaxa.4136.2.3

Stonis JR, Rocienė A (2013) Nepticulidae (Lepidoptera) of East Asia (1). Re-examination of the male genitalia of types deposited at the Russian Academy of Sciences. Zootaxa 3652: 1–59. doi: 10.11646/zootaxa.3652.1.1

Stonis JR, Rocienė A (2014) Additions to the Nepticulidae (Lepidoptera) of East Asia, with descriptions of three new species from Primorskiy Kray. Zootaxa 3846: 17. doi: 10.11646/zootaxa.3846.2.2

Stonis JR, Remeikis A, Sruoga V (2013a) An annotated list of the Opostegidae of the Himalaya, with a description of *Pseudopostega brevicaudata* sp. nov. (Lepidoptera: Nepticuloidea). Zootaxa 3609: 182–194. doi: 10.11646/zootaxa.3609.2.4

Stonis JR, Remeikis A, Diškus A, Noreika R (2013b) New Nepticulidae species (Insecta: Lepidoptera) from the Yucatán Peninsula. Zootaxa 3609: 223–230. doi: 10.11646/zootaxa.3609.2.8

Stonis JR, Diškus A, Remeikis A, Schuster J (2013c) First discovery of *Quercus* feeding Nepticulidae (Lepidoptera) in Central America. Zootaxa 3737: 1–23. doi: 10.11646/zootaxa.3737.1.1

Stonis JR, Diškus A, Remeikis A, Noreika R, Schuster J (2013d) Four new leaf-mining *Acalyptris* species from Guatemala and Belize, with new data on bionomics of *Stigmella pruinosa* (Lepidoptera: Nepticulidae). Zootaxa 3737: 101–117. doi: 10.11646/zootaxa.3737.2.1

Stonis JR, Diškus A, Remeikis A, Navickaite A, Rociene A (2013e) Description of new species of oak leaf-miners (Lepidoptera: Nepticulidae), with notes on the species groups of *Stigmella* Schrank associated with *Quercus* as a host-plant. Zootaxa 3737: 201–222. doi: 10.11646/zootaxa.3737.3.1

Stonis JR, Remeikis A, Davis DR (2014) Ten new species from the Patagonian Andes (Argentina and Chile), mostly belonging to a newly designated *Stigmella purpurimaculae* group (Lepidoptera: Nepticulidae). Zootaxa 3887: 321–353. doi: 10.11646/zootaxa.3887.3.3

Stonis JR, Diškus A, Remeikis A (2015a) The first description of the leaf-mining Nepticulidae (Lepidoptera) feeding on the South American plant genus *Liabum*, Asteraceae. Zootaxa 4040: 576–582. doi: 10.11646/zootaxa.4040.5.6

Stonis JR, Navickaite A, Diškus A (2015b) A puzzle regarding the Siberian *Ectoedemia jacutica* (Lepidoptera: Nepticulidae): re-examination and the first photographic documentation of the type series. Biologija 61: 116–122. doi: 10.6001/biologija.v61i3-4.3203

Stonis JR, Diškus A, Remeikis A, Torres NC (2016a) First description of leaf-mining Nepticulidae and Tischeriidae (Insecta, Lepidoptera) feeding on the Chilean endemic plant genus *Podanthus* Lag. (Asteraceae). Zootaxa 4061: 119–130. doi: 10.11646/zootaxa.4061.2.2

Stonis JR, Diškus A, Remeikis A, Davis DR, Solis MA, Torres NC (2016b) The first record of *Baccharis* L. (Asteraceae) as a host-plant genus for Nepticulidae (Lepidoptera), with description of new *Stigmella* species from South America. Zootaxa 4136: 101–128. doi: 10.11646/zootaxa.4136.1.4

Strand E (1934) *Niepeltia* novum genus Lepidopterorum. Internationale Entomologische Zeitschrift 28: 241.

Straus A (1976) Eine Gangmine im Eocän von Messel. Der Aufschluss 27: 445–446.

Straus A (1977) Gallen, Minen und andere Frassspuren im Pliokän von Willershausen am Harz. Verhandlungen des Botanischen Vereins der Provinz Brandenburg 113: 43–80.

Suire J (1951) Microlépidoptères des plantes caractérisant les zones naturelles de la Costière du Gard. Mémoires de la Société des Études et des Sciences Naturelles de Nîmes 8: 1–150, pls. i-vi.

Suire L (1928) Sur une *Nepticula* Française inédite. Encyclopédie Entomologique B, Lepidoptera 3: 127–129.

Svensson I (1966) New and confused species of Microlepidoptera. Opuscula Entomologica 31: 183–202, pls 1–4.

Svensson I (1976) Six new species of Microlepidoptera from northern Europe. Entomologica Scandinavica 7: 195–206. doi: 10.1163/187631276X00324

Svensson I (1982) Four new species of Microlepidoptera from northern Europe. Entomologica Scandinavica 13: 293–300. doi: 10.1163/187631282X00336

Svensson I (1985) Anmärkningsvärda fynd av Microlepidoptera i Sverige 1984. Entomologisk Tidskrift 106: 71–82.

Swezey OH (1921) *Opostega* in the Hawaiian Islands (Lep.). Proceedings of the Hawaiian Entomological Society 4: 531–536, pl. 539. http://biodiversitylibrary.org/page/25417459

Szőcs J (1957) New *Nepticula* species from Hungary. Annales Historico-Naturales Musei Nationalis Hungarici 49 (sn 8): 321–323. http://annales.nhmus.hu/cikkreszletes.php?idhoz=752

Szőcs J (1959) A new *Nepticula* species from Hungary (Lepidopt.). Acta Zoologica Academiae Scientiarum Hungariae 4: 417–419.

Szőcs J (1965) 7. család: Nepticulidae-Törpemolyok. In: Gozmány LA, Szőcs J (Eds) Lepidoptera 2 Molylepkék I, Microlepidoptera I. Akadémia Kiadó, Budapest, 48–104. [In Hungarian]

Szőcs J (1968) Some unknown data concerning miners (Lepidoptera). Acta Zoologica Academiae Scientiarum Hungariae 14: 225–231.

Szőcs J (1978) Adatok a Pili-hegység aknázómoly faunájához. (Data to the mining moths fauna from the Pilis Mts.). Folia Entomologica Hungarica, sn 31: 265–271. [In Hungarian]

Szőcs J (1981) Angaben über minierenden Motten aus Budapest und Umgebung. Folia Entomologica Hungarica, sn 42: 209–220.

Tengström JMJ (1848) Bidrag till Finlands Fjäril-Fauna. Notiser ur Sällskapets pro Fauna et Flora Fennica Förhandlingar 1: 69–164. http://biodiversitylibrary.org/page/34482255

Threlfall JH (1884) Notes on certain Tineae. The Entomologist 17: 113–114. http://biodiversitylibrary.org/page/16076900

Thunberg CP (1794) Dissertatio entomologica sistens insecta Svecica. Quorum partem septimam, Venia Exp. Ord. Med. Upsal. præside Carol. Pet. Thunberg, (...). Publico examini subjicit Gustavus Magnus Wenner, Holmiensis. In Audit. Botanico die 22 Dec. 1794. Joh. Fred. Edman, Upsala, 7: 83–98. http://gallica.bnf.fr/document?O=N099238

Toll S (1934a) *Nepticula heringi* sp. nova. Annales Musei Zoologici Polonici 11: 1–3.

Toll S (1934b) Jeszcze o krajowych gatunkach rozaju *Nepticula* Zell. (Noch über die heimischen Arten der Gattung *Nepticula* Zell.). Polskie Pismo Entomologiczne 13: 61–84.

Toll S (1936) Untersuchung der Genitalien bei *Pyrausta purpuralis* L. und *P. ostrinalis* Hb., nebst Beschreibung 11 neuer Microlepidopteren-Arten. Annales Musei Zoologici Polonici 11: 401–413, pls. 47–49.

Toll S (1957) *Nepticula nowakowskii* spec. nova aus Polen. Zeitschrift der Wiener Entomologischen Gesellschaft 42: 199–201. http://www.zobodat.at/pdf/ZOEV_42_0199-0201.pdf

Treitschke F (1833) Die Schmetterlinge von Europa, 9 (2). Schaben, Geistchen, G. *Hypsolopha-Orneodes*. Ernst Fleischer, Leipzig, 9 (2): 294 pp. http://biodiversitylibrary.org/page/34444079

Turati E (1923) Cinque anni di ricerche nell'Appennino Modenese (Note di Lepidotterologia). Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano 62: 4–74.

Turner AJ (1900) New Micro-Lepidoptera - mostly from Queensland. Transactions of the Royal Society of South Australia 24: 6–23. http://biodiversitylibrary.org/page/37025639

Turner AJ (1923) New Australian Micro-Lepidoptera. Transactions and Proceedings of the Royal Society of South Australia 47: 165–194. http://biostor.org/reference/114437

Tutt JW (1899) Natural History of the British Lepidoptera. A textbook for Students and Collectors, 1. London-Berlin, iv, 560 pp. doi: 10.5962/bhl.title.59327

Vári L (1944a) Eenige voor Nederland nieuwe Lepidoptera (3e Faunistische Mededeling). Tijdschrift voor Entomologie 86: xxv-xxvi. http://biodiversitylibrary.org/page/38811869

Vári L (1944b) Nederlandse Lepidoptera. 4e Faunistische Mededeling. Entomologische Berichten, Amsterdam 11: 213–216.

Vári L (1947) Biologische aantekeningen over *Fomoria weaveri* Stt. (Stigmellidae). Tijdschrift voor Entomologie 88: 521–523. http://www.biodiversitylibrary.org/part/65764

Vári L (1950) Nederlandse Lepidoptera. 5e Faunistische Mededeling. Entomologische Berichten, Amsterdam 13: 180–184.

Vári L (1951) Nederlandse Lepidoptera. 5e Faunistische Mededeling (slot). Entomologische Berichten, Amsterdam 13: 193–198.

Vári L (1955) South African Lepidoptera 1. Descriptions of new leafmining Tineina. Annals of the Transvaal Museum 22: 331–351.

Vári L (1961) South African Lepidoptera, 1. Lithocolletidae. Transvaal Museum Memoir 12: i-xix, 1–238, plates 231–112.

Vári L (1963) South African Lepidoptera, 3. Descriptions of new Stigmellidae. Koedoe 6: 66–75. doi: 10.4102/koedoe.v6i1.813

Vári L, Kroon D (1986) Southern African Lepidoptera. A series of cross-referenced indices. Lepidopterists Society of Southern Africa & The Transvaal Museum, Pretoria: x, 198 pp.

Vári L, Kroon DM, Krüger M (2002) Classification and checklist of the species of Lepidoptera recorded in Southern Africa. Simple Solutions, Chatswood, Australia: xxi, 384 pp.

Vaughan PH (1858) A new *Nepticula*. Entomologist's Weekly Intelligencer 5: 43–44. http://biodiversitylibrary.org/page/50830138

Vorbrodt K, Müller-Rutz J (1914) Die Schmetterlinge der Schweiz, II. Band. Wyss, Bern, 2: 726 pp. doi: 10.5962/bhl.title.37847

Wagner DL (1987) A new *Microcalyptris* species from California (Lepidoptera: Nepticulidae). Pan-Pacific Entomologist 63: 278–283.

Walker F (1864) List of the specimens of lepidopterous insects in the collection of the British Museum, 30. Tineites. Trustees of the British Museum (Natural History), London, 30: 837–1096. http://biodiversitylibrary.org/page/38948425

Walsingham L (1891) Micro-Lepidoptera collected near Cannes, 1890. Entomologist's Monthly Magazine 27: 137–152. http://biodiversitylibrary.org/page/9013552

Walsingham L (1897) Revision of the West-Indian Micro-Lepidoptera, with descriptions of new species. Proceedings of the General Meetings for Scientific Business of the Zoological Society of London 1897: 54–183. http://biodiversitylibrary.org/page/30987175

Walsingham L (1904) Spanish and Moorish Micro-Lepidoptera. Entomologist's Monthly Magazine 40: 7–8. doi: 10.5962/bhl.part.5486

Walsingham L (1907) Microlepidoptera. Fauna Hawaiiensis 1: 469–759. http://biodiversitylibrary.org/page/38854839

Walsingham L (1908a) Microlepidoptera of Tenerife. Proceedings of the Zoological Society of London 1907: 911–1034, pls. 51–53. http://biostor.org/reference/58131

Walsingham L (1908b) Spanish and Moorish Micro-Lepidoptera. Entomologist's Monthly Magazine 44: 226–229. doi: 10.5962/bhl.part.1630

Walsingham L (1911) Algerian Microlepidoptera. Entomologist's Monthly Magazine 47: 187–191. http://biodiversitylibrary.org/page/9016760

Walsingham L (1914) Insecta Lepidoptera-Heterocera, 4. Tineina, Pterophorina, Orneodina and Pyralidina and Hepialina (part.). In: Godman FD, Salvin O (Eds) Biologia Centrali-Americana. Dulau & co, London, 281–392, pl. 10. http://biodiversitylibrary.org/page/9016760

Walsingham L (1916) *Stigmella speciosa* Frey, an addition to the British list (Lep-Tin.). Entomologist's Monthly Magazine 52: 159–160. http://biodiversitylibrary.org/page/8958775

Waters EGR (1928) A new *Nepticula* from North Wales. Entomologist's Monthly Magazine 64: 105–106.

Watt MN (1921) The leaf-mining insects of New Zealand: Part II. The genus *Nepticula* (Lepidoptera). Transactions and Proceedings of the New Zealand Institute 53: 197–219, pls xl-xliii. http://rsnz.natlib.govt.nz/volume/rsnz_53/rsnz_53_00_002790.pdf

Watt MN (1924) The leaf-mining insects of New Zealand: Part 5. The genus *Nepticula* (Lepidoptera), and the Agromyzidae (Diptera) continued, and *Gracilaria selenites* Meyr. (Lepidoptera). Transactions and Proceedings of the New Zealand Institute 55: 674–687. http://rsnz.natlib.govt.nz/volume/rsnz_55/rsnz_55_00_006570.pdf

Weber P (1937) Über Mikrolepidopteren. Beschreibung neuer Arten und Formen aus der Schweiz, sowie Angaben über weniger bekannte Arten. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 16: 666–672. http://www.e-periodica.ch/cntmng?pid=seg-001:1934-1936:16::873

Weber P (1938) Zwei neue Arten der Mikrolepidopterengattung *Nepticula* und ein Beitrag zur Kenntnis von *Nepticula stelviana* Wck. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 17: 211–215. http://www.e-periodica.ch/cntmng?pid=seg-001:1937-1939:17::690

Weber P (1945) Die Schmetterlinge der Schweiz. 7. Nachtrag. Mikrolepidoptera. Mit Neubeschreibung von 5 Arten und 13 Formen. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 19: 347–407. http://www.e-periodica.ch/cntmng?pid=seg-001:1943-1946:19::819

Wilkinson C (1978) On the *Stigmella* - *Nepticula* controversy (Lepidoptera). Tijdschrift voor Entomologie 121: 13–22. http://biostor.org/reference/49887

Wilkinson C (1979) A taxonomic study of the micro-lepidopteran genera *Microcalyptris* Braun and *Fomoria* Beirne occurring in the United States of America (Lepidoptera, Nepticulidae). Tijdschrift voor Entomologie 122: 59–90. http://biostor.org/reference/49894

Wilkinson C (1981) A supplement to the genus *Ectoedemia* Busck (Nepticulidae: Lepidoptera) in North America, dealing with some difficult species and also some new ones. Tijdschrift voor Entomologie 124: 93–110. http://biostor.org/reference/49900

Wilkinson C (1982) Systematic entomology in a Dutch university. Antenna 6: 158–161.

Wilkinson C, Scoble MJ (1979) The Nepticulidae (Lepidoptera) of Canada. Memoirs of the Entomological Society of Canada 111: 1–129. doi: 10.4039/entm111107fv

Wilkinson C, Newton PJ (1981) The micro-lepidopteran genus *Ectoedemia* Busck (Nepticulidae) in North America. Tijdschrift voor Entomologie 124: 27–92. http://biostor.org/reference/49899

Wilkinson C, Bryan G, Menken SBJ, Nieukerken EJ van (1983) A clarification of the status of four taxa in the *Ectoedemia angulifasciella* group (Nepticulidae: Lepidoptera). Netherlands Journal of Zoology 33: 211–224. doi: 10.1163/002829683X00093

Wilson JO (1939) A new species of the family Nepticulidae (Lepidoptera). Transactions of the Royal Society of South Australia 63: 238–239. http://www.samuseum.sa.gov.au/Journals/TRSSA/TRSSA_V063/trssa_v063_p238p239.pdf

Wocke MF (1860) Mittheilungen über einige für die schlesische Fauna neue Arten. Jahresbericht der Schlesischen Gesellschaft für Vaterländische Kultur 38: 132–133. http://biodiversitylibrary.org/page/37119512

Wocke M (1861) II. Microlepidoptera. In: Staudinger O, Wocke M (Eds) Catalog der Lepidopteren Europas und der angrenzenden Länder Catalogue des Lépidopteres d'Europe et des pays limitrophes II Microlepidoptera. O. Staudinger & Hermann Burdach, Dresden, 85–130. https://archive.org/details/bub_gb_1S9DAAAAYAAJ

Wocke MF (1862) Reise nach Finmarken. II. Microlepidoptera. Stettiner Entomologische Zeitung 23: 233–257. http://biodiversitylibrary.org/page/8931313

Wocke MF (1865) Zwei neue Nepticulen. Stettiner Entomologische Zeitung 26: 269–270. http://biodiversitylibrary.org/page/8933761

Wocke M (1871) II. Microlepidoptera. In: Staudinger O, Wocke M (Eds) Catalog der Lepidopteren des europäischen Faunengebiets, II Microlepidoptera Catalogue ou enumération méthodique des lépidoptères qui habitent le terroire de la faune Européenne, II Microlepidoptera. O. Staudinger & Hermann Burdach, Dresden, 201–346, 383–414, 424–426. doi: 10.5962/bhl.title.9475

Wocke MF (1874) Verzeichniss der Falter Schlesiens, 2. Microlepidoptera. Zeitschrift für Entomologie (Breslau), N F 4: 1–107. http://biodiversitylibrary.org/page/31556630

Wocke MF (1877) Über einige wenig bekannte oder neue Falter der deutschen Fauna. Zeitschrift für Entomologie (Breslau), N F 6: 42–52. http://biodiversitylibrary.org/page/31571342

Wocke MF (1879) Lepidopterologische Mitteilungen. Zeitschrift für Entomologie (Breslau), N F 7: 70–80. http://biodiversitylibrary.org/page/43220148

Wocke MF (1881) Die Lepidopterenfauna des Stilfser Jochs. Fortsetzung II. Jahresbericht der Schlesischen Gesellschaft für Vaterländische Kultur 58: 198–205. http://biodiversitylibrary.org/page/46524364

Wolff NL (1955a) *Nepticula repentiella* n. sp. (Lepidoptera, Nepticulidae). Entomologiske Meddelelser 27: 82–90, pls. 1–4.

Wolff NL (1955b) *Nepticula benanderella* n. sp. (Lep., Nepticulidae). Opuscula Entomologica 20: 49–53.

Wolff NL (1957) *Trifurcula griseella* nov. spec. (Lepidoptera, Nepticulidae). Entomologiske Meddelelser 28: 19–22.

Wood JH (1890) *Nepticula torminalis*, a species new to science. Entomologist's Monthly Magazine 26: 209–210. http://biodiversitylibrary.org/page/33378985

Wood JH, Walsingham L (1894) *Nepticula confusella*, a new birch-mining species. Entomologist's Monthly Magazine 30: 272–273. http://biodiversitylibrary.org/page/32438162

Wörz A (1937) *Nepticula hahniella* spec. nov. m. Entomologische Rundschau 54: 290–292.

Yang C-K (1989) *Sinopticula sinica* (Lepidoptera: Nepticulidae) a new genus and species from China. Entomotaxonomia 11: 79–82.

Zangheri S (1969) Regno Animale. Sottord. Heterocera. In: Zangheri S (Ed) Repertorio sistematico e topografico della Flora e Fauna vivente e fossile della Romagna, 3. Museo Civico di Storia Naturale, Verona, 855–1015.

Zeller PC (1839) Versuch einer naturgemässen Eintheilung der Schaben. Isis von Oken 1839: 167–220. http://www.biodiversitylibrary.org/part/9422

Zeller PC (1843) [no title]. In: Vierter Bericht des Schlesischen Tauschvereins für Schmetterlinge : 21.

Zeller PC (1847) Bemerkungen über die auf einer Reise nach Italien und Sicilien beobachteten Schmetterlingsarten X. Isis von Oken 1847: 881-904. http://biodiversitylibrary.org/page/13234103

Zeller PC (1848) Die Gattungen der mit Augendeckeln versehenen blattminirenden Schaben. Linnaea Entomologica 3: 248–344, pl. 242. http://biodiversitylibrary.org/page/44017229

Zeller PC (1873) Beiträge zur Kenntnis der nordamericanischen Nachtfalter, besonders der Microlepidopteren. Zweite Abtheilung. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien 23: 201–334, 2 pls. http://biodiversitylibrary.org/page/16411868

Zeller PC (1877) Exotische Microlepidoptera. Horae Societatis Entomologicae Rossicae 13: 3–491, 6 pls. http://biodiversitylibrary.org/page/12834243

Zetterstedt JW (1838–1840) Insecta Lapponica. Lipsiae: vi, 1140 pp. doi: 10.5962/bhl.title.8242

Zimmerman EC (1978) Microlepidoptera. Part 1, Monotrysia, Tineoidea, Tortricoidea, Gracillarioidea, Yponomeutoidea, and Alucitoidea. The University Press of Hawaii, Honolulu, Insects of Hawaii 9 (1): xviii, 881 pp.

Zimmermann F (1940) Eine neue Nepticulide aus Deutschland (Lep.), *Ectoedemia liebwerdella* spec. nov. (Lep.). Mitteilungen aus dem Zoologischen Museum in Berlin 24: 264–265. doi: 10.1002/mmnz.19390240204

Zimmermann F (1944) Zur Kenntnis der Verbreitung der Nepticuliden in den Reichsgauen Wien und Niederdonau (Lepidopt.). Zeitschrift der Wiener Entomologischen Gesellschaft 29: 107–122. http://www.zobodat.at/pdf/ZOEV_29_0107-0122.pdf

Note added in proof

Just before publication, another new species of Nepticulidae was published, and two more described, but not named (Stonis et al. 2016c). The following species should be added to page 115, the *Stigmella epicosma* group. The species has also been added to the online databases.

Stigmella polylepiella Diškus & Stonis in Stonis et al., 2016c: 86 NEO

Stonis JR, Diškus A, Remeikis A, Karsholt O (2016c) Do leaf-mining Nepticulidae occur in the natural but so threatened Andean *Polylepis* forests? Biologija 62: 83–97. doi: 10.6001/biologija.v62i2.3334.

Supplementary material 1

Specimen data for Catalogue of Nepticuloidea with BOLD data

Authors: Erik J. van Nieukerken, Camiel Doorenweerd, Robert J. B. Hoare, Donald R. Davis

Data type: Microsoft Excel (xls).

Explanation note: The 213 records comprise specimens that have either been cited briefly or completely in the Notes, or have been used more implicitly for the texts and figures. Records are listed alphabetically by species, country, stateProvince, AdminDivision2, AdminDivision3, Locality.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Supplementary material 2

BOLD TaxonID Tree of the dataset Lepidoptera - Nepticuloidea of the World 2016 [DS-NEPCAT]

Authors: Erik J. van Nieukerken, Camiel Doorenweerd, Robert J. B. Hoare, Donald R. Davis

Data type: PDF file

Explanation note: A Neighbor joining tree of COI sequences, with Kimura 2 Parameter as distance model for sequences linger than 200 bp, 3203 records.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons.org/licenses/odbl/1.0/). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.